Interstellar Bow Shocks around Fast Stars Passing through the Local Interstellar Medium

J. Michael Shull

Department of Astrophysical and Planetary Sciences
University of Colorado, Boulder, CO 80309

S. R. Kulkarni

Owens Valley Radio Observatory, 249-17
California Institute of Technology, Pasadena, CA 91125

`michael.shull@colorado.edu, srk@astro.caltech.edu`

ABSTRACT

Bow-shocks are produced in the local interstellar medium by the passage of fast stars from the Galactic thin-disk and thick-disk populations with velocities $V_* = 40$–80 km s^{-1}. Stellar transits of local H I clouds occur every 3500–7000 yr on average and last between 10^4 and 10^5 yr. There could be 10–20 active bow shocks around low-mass stars inside clouds within 15 pc of the Sun. At local cloud distances of 3–10 pc, their turbulent wakes have transverse radial extents $R_{\rm wake} \approx 100$–300 AU , angular sizes 10–100 arcsec, and Lyα surface brightnesses of 2-8 Rayleighs in gas with total hydrogen density $n_{\rm H} \approx 0.1$ cm^{-3} and $V_* =$ 40-80 km s^{-1}. These transit wakes may cover an area fraction $f_A \approx (R_{\rm wake}/R_{\rm cl}) \approx 10^{-3}$ of local H I clouds and be detectable in IR (dust), UV (Lyα, two-photon), or non-thermal radio emission. Turbulent heating in these wakes could produce the observed elevated rotational populations of H$_2$ ($J \geq 2$) and influence the endothermic formation of CH$^+$ in diffuse interstellar gas at $T > 10^3$ K.

1. Introduction

Recent observations have advanced our knowledge of gas clouds in the interstellar medium (ISM) and the velocity dispersions of stars in the thin-disk and thick-disk populations. This is an opportune moment for a new look at the frequency and physical effects of bow shocks produced by stars passing through low-density H I clouds near the Sun. Most of these transits will be by low-mass stars, although more distant examples of bow shocks around runaway O-type stars have been found in infrared emission (Van Buren & McCray 1988; Peri et al. 2015; Kobulnicki et al. 2016). The *Herschel* far-infrared survey (Cox et al. 2012) identified numerous bow shocks around AGB stars and red supergiants, including Betelgeuse (Decin et al. 2012). Theoretical studies of intermediate-velocity shocks (Kulkarni & Shull 2023) suggest that the far-UV background radiation may include contributions from Lyα and two-photon continuum emission from the metastable (2s) state of H I. To excite the *2s* and *2p* levels (10.2 eV) requires shock velocities $V_s > 40$ km s^{-1} with post-shock temperatures $T_s \approx (3\mu V_s^2/16k) \approx (36,300 \text{ K}) V_{40}^2$, where $V_{40} = V_s/40$ km s^{-1}. Magnetic fields will reduce this post-shock temperature.

In addition to their astronomical importance, fast stars passing through local clouds contribute to the thermal energy budget of interstellar gas in their transit wakes and extended regions. Observations of interstellar H_2 in rotationally excited states with angular momentum quantum number $J \geq 2$ suggest additional excitation mechanisms besides UV-pumping (Gry et al. 2002; Nehmé et al. 2008; Ingalls et al. 2011; Shull, Danforth, & Anderson 2021). Elevated rotational populations could be produced by collisional excitation in warm molecular gas ($T > 500$ K) heated by MHD turbulent dissipation (Moseley et al. 2021). The observed correlation of CH^+ with rotationally-excited H_2 (Jensen et al. 2010) has led to similar production schemes for CH^+ involving warm H_2 (Falgarone & Puget 1995; Myers et al. 2015). Local clouds of diffuse interstellar gas exhibit significant turbulent line widths of 2–3 km s^{-1} (Linsky et al. 2022) derived by comparing the widths of absorption lines from atomic species of different masses. They found that the LISM is inhomogeneous on length scales less than 5100 AU. The inferred temperature distribution appears Gaussian, with some sight lines reaching $T \approx 12,000$ K. Table 1 lists a number of local "fast stars", together with their velocities and properties of the LISM (gas temperatures and turbulent velocities).

Strong shocks will be produced by stars moving at velocities (40–70 km s^{-1}) through the warm neutral medium of local clouds, in excess of the sound speed c_s (or magnetosonic wave speed v_m). For magnetized gas, magnetosonic waves are characterized by a combination of adiabatic sound speed, $c_s = (5kT/3\mu)^{1/2} = (9.30 \text{ km s}^{-1}) T_{8000}^{1/2}$, and Alfvén velocity, $v_A = (5.53 \text{ km s}^{-1})(B/3\,\mu G) n_{\rm H}^{-1/2}$. Here, we scaled to gas temperature $T \approx (8000 \text{ K}) T_{8000}$, total hydrogen density $n_{\rm H}$ (in cm^{-3}), mean particle mass $\mu = 1.27 m_H$ for $y \equiv \text{He/H} = 0.1$, and magnetic field strength B. Absorption-line studies (Frisch et al. 2011; Linsky et al. 2022) of clouds in the local interstellar medium (LISM) suggest $n_{\rm H} \approx 0.1$ cm^{-3} and $B \approx$ 3–4 μG (McComas et al. 2012; Zank et al. 2013). For wave propagation perpendicular to the magnetic field, the fast-mode magnetosonic wave speed, $v_m = (v_A^2 + c_s^2)^{1/2}$. Considering spatial variations in LISM parameters (n_H, T, B), magnetosonic wave speeds in the LISM could range from 20 to 30 km s^{-1}, with further uncertainties arising from propagation angle and ionization fraction[1].

Observations of the kinematics of the Milky Way disk suggest that populations of older stars in the Galactic disk have supersonic and super-Alfvénic velocities relative to the LISM. The large stellar velocity dispersions measured in the Galactic thin disk (Nordstrom et al. 2004; Anguiano

---

[1] Ionization fractions are important at the shock interface, owing to charge exchange reactions with neutrals. Scherer & Fichtner (2014) also pointed out the importance of including both He^+ and H^+ in determining the speed of Alfvén waves. For wave propagation (vector $\mathbf{k}$ at angle ϕ to the magnetic field, the fast-mode (Kulsrud 2005) dispersion relation $(\omega/k)^2 = (v_A^2 + c_s^2)/2 + \frac{1}{2}[(v_A^2 - c_s^2)^2 + 4 v_A^2 c_s^2 \sin^2\phi]^{1/2}$ defines the phase and group velocities of the wave packets. Near the front of the bow shock, we expect that $\phi \approx 90^\circ$ ($\mathbf{k} \perp \mathbf{B}$). Away from the forward region, the fast-mode wave speed could be somewhat smaller than $(v_A^2 + c_s^2)^{1/2}$.

et al. 2020) are likely the result of dynamical heating processes that increase the stellar velocities. Massive Galactic objects (spiral arms, giant molecular clouds, bars, infalling dwarf satellites) perturb the motions of the newborn stars, causing their random velocities to increase with time. A recent study (Viera et al. 2022) using *Gaia*-DR3 proper motions found two kinematically distinct Gaussian components: a thin disk with velocity dispersions $(\sigma_R, \sigma_\phi, \sigma_Z) = 30$, 21, and 11 km s^{-1} and a thick disk with dispersions of 49, 35, and 22 km s^{-1}. These correspond to 2D (in-plane) dispersions of $\sigma_{\rm 2D} = 37$ km s^{-1} (thin disk) and 60 km s^{-1} (thick disk). The vertical (out-of-plane) dispersions (11 and 22 km s^{-1}) and total 3D dispersions (38 to 64 km s^{-1}) also differ between the components. These velocities are considerably larger than those of stars forming inside dense molecular gas, seen in stellar associations and star clusters with radial velocity dispersions $\sigma_{\rm 1D}$ =1–3 km s^{-1}. Proper motion measurements with *Gaia* (Kuhn et al. 2019; Shull, Darling, & Danforth 2021) show that young star clusters are expanding at velocities ~ 0.5 km s^{-1} and will disperse in several Myr after the gas is expelled.

2. Stellar Bow Shocks in the Local ISM

2.1. Star-Cloud Encounters

Because of the large velocity dispersions for stars in Galactic disk populations, H I clouds in the LISM will have frequent encounters with stars at velocities 40–70 km s^{-1}. Absorption-line studies along 157 stellar sight lines (Redfield & Linsky 2008; Linsky et al. 2019) identified numerous warm H I clouds within 15 pc of the Sun. The nearest structures within $d \sim 4$ pc are labeled the Local Interstellar Cloud (LIC), the G Cloud, and the Blue Cloud. Redfield & Linsky (2008) estimated that between 6 and 19% of the LISM volume is filled with warm H I gas. An updated analysis (Linsky et al. 2022) found that clouds within 4 pc are more tightly packed, with mean hydrogen density $\langle n_{\rm HI} \rangle \approx 0.1$ cm^{-3} and typical volume factors $f_V \approx 0.38 - 62$. For local clouds beyond 4 pc, they find $f_V < 0.4$.

We now estimate the rate at which passing stars encounter warm H I gas clouds in the LISM from cloud cross sections, stellar space densities, and velocity dispersions. If the region within distance $R_{\rm LISM}$ of the Sun is filled by $N_{\rm cl}$ spherical clouds of radius $R_{\rm cl}$, the volume filling factor is $f_V = N_{\rm cl}(R_{\rm cl}/R_{\rm LISM})^3$, and the area covering factor is $f_A = N_{\rm cl}(R_{\rm cl}/R_{\rm LISM})^2$. These factors are related by $f_A = f_V(R_{\rm LISM}/R_{\rm cl})$ for spheres. More generally, f_A and f_V depend on cloud geometry, although the non-spherical or filamentary shapes of some local clouds do not change these estimates substantially. For 15 clouds within 10 pc, the above approximation suggests mean cloud radii, $R_{\rm cl} = (10~{\rm pc})(f_V/N_{\rm cl})^{1/3} \approx 2$–3 pc and area covering factors $f_A \approx 0.7$–1.0 for $f_V =$ 0.15–0.30.

The star-cloud encounter rate depends on the local density of stars, both those currently seen within the local clouds and the stars in the disk populations that transit the LISM at high velocity. The recent Gaia-DR3 survey of the local volume within 10 pc (Reylé et al. 2022) found 541 stars in 336 systems, of which 286 were luminous stars (we do not include the 85 brown dwarfs in the survey). This corresponds to a mean density of 0.065 pc^{-3}, similar to that quoted in Table 4.5 of Mihalas & Binney (1981). The normalized densities of stars in the thin-disk and thick-disk populations should sum to similar values, although there is some disagreement as to their relative local densities (Anguiano et al. 2020; Everall et al. 2022; Viera et al. 2023). Not all of these stars will have velocities sufficient to produce the type of shocks discussed here. Only a fraction will have the high velocities needed to exceed magnetosonic speeds. The radial extent of the astrospheres, bow shocks, and turbulent wakes depend on the presence of a moderately strong stellar wind (Herbst et al. 2020).

To be conservative, we scale our encounter rate to a local density $n_* = 0.01\ \text{pc}^{-3}$ of stars of mean velocity $V_* = (40\ \ \text{km s}^{-1})V_{40}$, passing through an extended LISM with $R_{\text{LISM}} = (15\ \text{pc})R_{15}$. The collision rate with local clouds is

$$k_{\text{coll}} = n_*(\pi R_{\text{LISM}}^2) f_A V_* \approx (3 \times 10^{-4}\ \text{yr}^{-1})\, V_{40} R_{15}^2 f_A\ . \tag{1}$$

For $f_A \approx 0.5$–1.0, the encounter rate is one collision every 3500–7000 yr. A typical stellar crossing time through a cloud of mean length $\ell \approx 4R_{\text{cl}}/3 \approx 3$ pc is

$$t_{\text{cross}} \approx \ell/V_* \approx (70{,}000\ \text{yr})(\ell/3\ \text{pc})V_{40}^{-1}\ , \tag{2}$$

and the number of active bow-shock encounters within 15 pc is

$$N_{\text{bow}} = k_{\text{coll}}\, t_{\text{cross}} \approx 20\,(\ell/3\ \text{pc}) f_A R_{15}^2\ . \tag{3}$$

The chances of detection could be significantly higher for larger cloud complexes beyond the LISM (Frisch et al. 2011). However, the local clouds lie inside a low-density cavity, and the nearest star-forming clouds are more distant, including gas in the Taurus molecular cloud (140 pc) and the Scorpius-Centaurus association (130 pc).

2.2. Estimated Bow Shock Sizes

The two *Voyager* missions passed through the solar-wind termination shock at distances of 94 and 84 AU (Stone et al. 2005; Richardson et al. 2008; Stone et al. 2008). Low-mass stars like the Sun, with weak winds and low mass-loss rates, are expected to produce astrospheres preceded by bow shocks with transverse radial extents R_{wake} ≈100–300 AU, similar to that of the Sun's heliosheath and termination shock[2] of the solar wind. These estimates are consistent with results

---

[2] Observations from the *Interstellar Boundary Explorer* (McComas et al. 2012; Zank et al. 2013; McComas et al. 2019) suggest that the solar wind termination may be in a "bow wave" rather than a strong shock, owing to the influence of inclined magnetic fields in the heliosphere and LISM. This interpretation has been questioned by Scherer & Fichtner (2014), who noted the importance of including both He^+ and H^+ in the speeds of Alfvén and magnetosonic waves.

from an HST survey (Wood et al. 2021) of astrospheric Lyα absorption toward nearby M-dwarf stars. For stars with super-magnetosonic motions through the ISM, the hydrogen wall and bow shock in MHD models (e.g., Zank et al. 2013) precede the wind termination shock by a substantial factor that depends on the stellar speed and magnetosonic Mach number relative to the surrounding medium. This warm, extended gas is influenced in density, temperature, and wave speeds. The trailing wakes typically have transverse radial extent 2–3 times the size of the forward radius of the termination shock. Most low-mass stars have mass-loss rates comparable to (and sometimes less than) that of the Sun, $\dot{M} \approx 4 \times 10^{-14}\ M_\odot\ \mathrm{yr}^{-1}$. At the typical 3–10 pc distances to the local interstellar clouds, the bow-shock angular sizes could range from 10–100 arcsec. More distant bow shocks would be smaller (0.1 arcsec at 1 kpc) and probably undetectable.

A few, more distant fast-moving OB stars with much stronger winds have observed arc-like features, attributed to bow shocks of parsec size (Van Buren & McCray 1988; Mackey et al. 2016). An O-star moving at velocity $V_* \approx 40$ km s^{-1} would produce a wind-termination shock of approximate size,

$$r_b \approx \left[\frac{\dot{M}_w V_w}{4\pi \mu n_{\rm H}} \right]^{1/2} V_*^{-1} \approx (1.0\ {\rm pc})\, \dot{M}_{-6}^{1/2} n_{\rm H}^{-1/2} \left(\frac{V_w}{1000\ {\rm km\ s}^{-1}} \right)^{1/2} \left(\frac{V_*}{40\ {\rm km\ s}^{-1}} \right)^{-1} . \tag{4}$$

This termination shock in O-star winds will be preceded a wind-driven bubble of high-pressure gas (Weaver et al. 1977; Mackey et al. 2016), whose shape can be distorted by supersonic stellar motion. Here, we express the stellar mass-loss rate as $\dot{M} = (10^{-6}\ M_\odot\ \mathrm{yr}^{-1})\dot{M}_{-6}$ and equate the wind ram pressure, $\dot{M}_w V_w / 4\pi r^2$, with the pressure, $\rho_{\rm ISM} V_*^2$, of the ISM of mass density $\rho_{\rm ISM} = 1.4 m_{\rm H} n_{\rm H}$. At $d \approx 1$ kpc, the angular sizes would be $\theta_b = r_b/d \approx 10^{-3}$ rad $(3.4'\ d_{\rm kpc}^{-1})$. The above estimate of r_b is uncertain by factors of 1.5–2.0 when one includes the effects of magnetic fields.

2.3. Estimated Luminosity and Surface Brightness

Intermediate velocity shocks moving into H I gas will produce optical and UV emission lines that could be signatures of fast-star passages through local clouds. These include Lyα, metal-ion emission lines, and two-photon continuum from the metastable (2s) state of H I (Shull & McKee 1979). An analysis of the bow-shock luminosity and surface brightness requires calculations of the photon flux, Φ_0, which is proportional to the flux, $n_{\rm H} v_s$, of hydrogen atoms entering the shock front. The radiative shock models (Shull & Silk 1979) of primeval material (H and He) found photon fluxes that can be fitted to expressions rising steeply for V_s between 15 and 100 km s^{-1},

$$\Phi_0(2\gamma) = (2.2 \times 10^4\ {\rm phot\ cm^{-2}\ s^{-1}})(n_{\rm H}/0.1\ {\rm cm^{-3}}) V_{40}^{2.18} , \tag{5}$$

$$\Phi_0({\rm Ly}\alpha) = (2.7 \times 10^5\ {\rm phot\ cm^{-2}\ s^{-1}})(n_{\rm H}/0.1\ {\rm cm^{-3}}) V_{40}^{2.12} . \tag{6}$$

We use these expressions to estimate the luminosities of Lyα and two-photon emission produced in a bow shock of radius r_b from a low-mass star located at distance d. Because of the approximate nature of these estimates, we consider plane-parallel shocks at small angles from the forward direction of the star. More careful 3D models with MHD codes would clearly be warranted in the future. Most of the emission comes from a polar cap with opening angle $\theta_{\rm cap} \approx 30^\circ$ relative to the stellar velocity vector. This cap occupies a fraction $(1 - \cos\theta_{\rm cap})$ of the $2\pi r_b^2$ area of the forward hemisphere, with a photon luminosity,

$$S_{\rm phot} = 2\pi r_b^2 \Phi_0 (1 - \cos\theta_{\rm cap}) \ , \tag{7}$$

and its projected solid angle on the sky is approximately

$$\Omega_{\rm arc} \approx \frac{2 r_b \theta_{\rm cap} (\Delta r)}{d^2} = \left(\frac{2 r_b^2}{d^2} \right) \theta_{\rm cap} (1 - \cos\theta_{\rm cap}) \ . \tag{8}$$

Here, we adopt an angular length, $2\theta_{\rm arc} = 2\theta_{\rm cap}$, for the arc (angles in radians) and a projected arc width $\Delta r = r_b(1 - \cos\theta_{\rm cap})$. Using eq. (7), the arc surface brightness is

$$B_{\rm arc} = \frac{S_{\rm phot}/4\pi d^2}{\Omega_{\rm arc}} = \frac{\Phi_0}{4\theta_{\rm cap}} \ . \tag{9}$$

As expected for optically thin emission, the brightness is independent of both d and r_b.

For solar-type stars passing through the LISM at $V_* \approx (40 \text{ km s}^{-1}) V_{40}$, we adopt $r_b = 100$ AU, $n_{\rm H} \approx 0.1 \text{ cm}^{-3}$, and $\theta_{\rm cap} \approx 30^\circ$ to find

$$\begin{aligned} S_{\rm phot}(2\gamma) &= (4.2 \times 10^{34} \text{ phot s}^{-1}) \left(\frac{n_{\rm H}}{0.1 \text{ cm}^{-3}} \right) V_{40}^{2.18} \ , & (10) \\ S_{\rm phot}({\rm Ly}\alpha) &= (5.1 \times 10^{35} \text{ phot s}^{-1}) \left(\frac{n_{\rm H}}{0.1 \text{ cm}^{-3}} \right) V_{40}^{2.12} \ . & (11) \end{aligned}$$

With the 10.2 eV energies of the upper (*2s*, *2p*) levels, these luminosity coefficients sum to 9×10^{24} erg s^{-1}, comparable to the kinetic energy of interstellar gas entering the bow shock over the cap portion of the front,

$$\frac{\rho_{\rm ISM} V_s^3}{2} (2\pi r_b^2)(1 - \cos\theta_{\rm cap}) \approx (1.4 \times 10^{25} \text{ erg s}^{-1}) \left(\frac{n_{\rm H}}{0.1 \text{ cm}^{-3}} \right) V_{40}^3 \ . \tag{12}$$

Additional heating of the dust in the gas assembled by the bow front will come from stellar radiation, particularly in the case of OB stars. The Lyα photons will scatter in the low-density local H I, but owing to low H I column densities only a small fraction will be absorbed by dust grains. However, some of these bow shocks may also be seen in infrared emission, as observed for OB-stars and young-star outflows with the *Spitzer Space Telescope* (Winston et al. 2012; Kobulnicki et al. 2016) and the *Wide-field Infrared Survey Explorer* (Peri et al. 2015).

Most of the surface brightness of the bow shock comes from the material directly in front of the star, visible on the sky as a small arc of emission (Kobulnicky et al. 2016). Adopting $\theta_{\rm cap} = 30^\circ$ ($\pi/6$ radians) and $\Delta r / r_b \approx 0.13$, and using equations (5) and (6) for Φ_0, we estimate photon surface brightnesses, $B_{\rm arc} \approx (3\Phi_0/2\pi)$ of the arc,

$$\begin{aligned} B_{\rm cap}(2\gamma) &\approx (10{,}500 \text{ phot cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}) V_{40}^{2.18} \left(\frac{n_{\rm H}}{0.1 \text{ cm}^{-3}} \right) \ , & (13) \\ B_{\rm cap}({\rm Ly}\alpha) &\approx (129{,}000 \text{ phot cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}) V_{40}^{2.25} \left(\frac{n_{\rm H}}{0.1 \text{ cm}^{-3}} \right) \ . & (14) \end{aligned}$$

The above brightness coefficients (in units photons cm^{-2} s^{-1} sr^{-1}) can also be expressed as 0.13 and 1.6 Rayleighs. Thus, for stars in the thin-disk and thick-disk populations, with rms velocities of 37 km s^{-1} and 60 km s^{-1} respectively, the Lyα surface brightnesses could range from 2–4 Rayleighs in gas with $n_{\rm H} \approx 0.1$ cm^{-3}. Even greater brightnesses might appear around several of the fast local stars listed in Table 1.

3. Turbulent Heating of the LISM gas

The LISM clouds penetrated by transiting low-mass stars will be affected by turbulence generated by transiting fast stars at various shocks and in the wake. In clouds of pathlength $\ell = 2$–3 pc, stars moving at 40–80 km s^{-1} will spend 30,000–100,000 years, leaving wakes of dynamically disturbed gas. Owing to downstream flows from the bow shock, the transverse sizes of the wakes will be larger than the forward termination shock and astrosheath (e.g., Mac Low et al. 1991; Mackey et al. 2016). The fractional volume of the cloud affected by the turbulent wake is

$$f_V \approx \left(\frac{R_{\rm wake}}{R_{\rm cl}}\right)^2 \approx (2\times10^{-6})\left[\frac{R_{\rm wake}/300\ {\rm AU}}{R_{\rm cl}/{\rm pc}}\right]^2 \ . \tag{15}$$

Note that $R_{\rm wake}$ is the transverse size of the post-bowshock flow, not its length. The total affected portion of local clouds includes both the direct "bore holes" and extended regions influenced by heat generated by MHD wave damping. As described by Spitzer (1982) and Ferriére et al. (1988), the waves arise from reflected shocks produced by interactions with density irregularities encountered by the bow shock. After a nonlinear cascade to smaller eddies, much of the turbulence will decay into heat, on an eddy-turnover time,

$$t_{\rm eddy} \approx R_{\rm eddy}/V_{\rm eddy} \approx (10^3\ {\rm yr})\left(\frac{r_b}{100\ {\rm AU}}\right)\left(\frac{3\ {\rm km/s}}{V_{\rm eddy}}\right) \ . \tag{16}$$

Here, we assumed the initial eddy to have a characteristic size $R_{\rm eddy} \approx r_b = 100$ AU at the termination shock scale and an initial turbulent velocity $V_{\rm eddy} \approx 3$ km s^{-1}. After cascading to smaller scales through nonlinear effects, the turbulent energy is dissipated. The turbulent heating rate can be estimated by dividing the kinetic energy density ($\rho_{\rm gas} V_{\rm eddy}^2$) by the eddy turnover time,

$$\epsilon_{\rm turb} \approx \rho_{\rm gas} V_{\rm eddy}^3/R_{\rm eddy} \approx (4\times10^{-23}\ {\rm erg\ s^{-1}}) n_{\rm H}\left(\frac{r_b}{100\ {\rm AU}}\right)\left(\frac{3\ {\rm km\ s^{-1}}}{V_{\rm eddy}}\right)^3 \ , \tag{17}$$

where we assumed $\rho_{\rm gas} = 1.4 n_{\rm H} m_{\rm H}$. The turbulent heating time would be

$$t_{\rm heat} \approx \frac{3nkT/2}{\epsilon_{\rm turb}} \approx (10^3\ {\rm yr})\left(\frac{r_b}{100\ {\rm AU}}\right)^{-1}\left(\frac{V_{\rm eddy}}{3\ {\rm km\ s^{-1}}}\right)^{-3}\left(\frac{T}{6000\ {\rm K}}\right) \ . \tag{18}$$

Because this time is much shorter than the stellar crossing time through the cloud, the thermal state of gas in the bow-shock wake will be time dependent. The situation will involve transient heating over multiple time scales, with $t_{\rm heat} < t_{\rm cool}$ and $t_{\rm cool} \approx t_{\rm cross}$, The post-shock wake will cool gradually, as the turbulent eddies quickly cascade down to smaller scales. Gas temperatures will rise transiently to $T \approx 20,000 - 30,000$ K, where H I can be collisionally ionized. Radiative cooling by H I (Lyα) and metal forbidden lines will reduce the temperature to $T \leq 10^4$ K on an isobaric cooling timescale,

$$t_{\rm cool} = \frac{5 n_{\rm tot} kT/2}{n_e n_H \mathcal{L}(T)} \approx (50,000 \text{ yr}) \left(\frac{n_{\rm H}}{0.1 \text{ cm}^{-3}} \right)^{-1} \left(\frac{T}{20,000 \text{ K}} \right) \mathcal{L}_{-22}^{-1} \ , \tag{19}$$

comparable to the stellar crossing time. We assume a partially ionized medium with $n_e \approx 0.5 n_{\rm H}$ and $n_{\rm tot} \approx 1.6 n_{\rm H}$, and a cooling rate coefficient $\mathcal{L} \approx (10^{-22} \text{ erg cm}^3 \text{ s}^{-1}) \mathcal{L}_{-22}$ at 20,000 K. Analyzing the time-dependent evolution of ionization and temperature behind bow shocks is a problem in hydrodynamics and thermodynamics worth further study. The effects of magnetic fields in the turbulent wakes (Baalmann et al. 2021 add to the complexities, including reconnection and dissipation of magnetic field energy.

In our model we have ignored cooling through all elements other than hydrogen and ignored external heat inputs. By 10^4 K the cooling power from hydrogen decreases with temperature and will compete with cooling by forbidden and fine structure lines of metals. As cooling proceeds, the internal heat is proportionally reduced and heating by external agents (e.g., ionization by cosmic rays, photoelectric dust heating, ionizing soft X-ray radiation) will become important. The recombination time scale is about $10^5 n_{\rm H}^{-1}$ yr. The ionization time per atom from the diffuse EUV field which ionizes the warm ionized medium is 10^6 yr.

4. Conclusion and Future Prospects

This paper analyzes the interstellar bow-shocks in the LISM generated around passing stars from the Galactic thin disk and thick disk with velocities of 40–70 km s^{-1}. We estimate that bow shocks transit the local H I clouds every 3500–7000 yr. On average, there could be 10–20 active bow shocks inside clouds within 15 pc of the Sun, with stellar transit times varying from 10^4–10^5 yr. At typical local cloud distances of 3–10 pc, the bow shocks and astrospheres around low-mass stars likely have 100–300 AU transverse radial extent and angular sizes of 10–100 arcsec. Their turbulent transit wakes of passing stars might cover an area fraction $f_A \approx (R_{\rm wake}/R_{\rm cl})^2 \approx 10^{-3}$ of nearby clouds. They could explain the 2–3 km s^{-1} turbulent widths and temperature variations on scales less than 5100 AU in LISM clouds (Linsky et al. 2022), as well as elevated rotational populations of H_2 and endothermic production of CH$^+$. Bow shocks around rarer massive stars and AGB stars would produce shocks of parsec scale and angular sizes of several arcmin (at ~ 1 kpc).

Detecting the low-mass bow shocks will require wide-field surveys, either in the infrared (dust emission) or in the ultraviolet (for Lyα and two-photon emission). The IR detections could be a task for the *WISE* data base, and Lyα emission could be a project for the Alice spectrograph aboard the New Horizons (NH) spacecraft during its extended mission. During six great-circle scans (at 40-47 AU) the NH spectrograph observed interplanetary Lyα (Gladstone et al. 2021) with a uniform background of 43 ± 3 Rayleighs over the sky, interpreted as the Galactic Lyα background. There is also resonantly scattered solar Lyα, which at the current NH distance (56 AU from the Sun) has fallen to about the same level. Thus, the typical brightness in any direction (away from the Sun) would be 80–100 R. Our calculations of far-UV radiation from the forward caps of 40–80 km s^{-1} bow shocks suggest that arc-like structures around low-mass stars transiting local clouds with $n_{\rm H} \approx 0.1$ cm^{-3} should have surface brightnesses of 2–8 Rayleigh in Lyα and 0.1–0.4 Rayleigh in the UV two-photon continuum. These could possibly be seen with NH and with slit spectroscopy on the proposed UVEX mission (Kulkarni et al. 2021).

Examples of bow shocks around runaway O-type stars and AGB stars have already been found in infrared emission (Kobulnicki et al. 2016; Peri et al. 2015; Cox et al. 2012). Wide-angle searches in Lyα, far-UV continua, and infrared emission for lower luminosity examples could detect the more numerous cases of low-mass stars transiting the LISM. Their signatures might also be detectable in non-thermal radio emission from particles accelerated at the termination shock and bow-shock vicinity. Although the predicted stellar transit wakes have small covering fractions of LISM clouds, both by volume and on the sky, the effects of down-stream turbulent dissipation, MHD wave heating, and energetic particle acceleration could be more widespread throughout the local clouds.

We thank Jeffrey Linsky and Seth Redfield for discussions on gas structures in the local interstellar medium, Randy Gladstone for insights on Lyα scanning observations with the New Horizons spacecraft, and Brian Wood for providing characteristics of the winds and astrospheric absorption around nearby solar type stars. We also thank the anonymous referee for introducing us to important papers on MHD models of astrospheres.

REFERENCES

Anguiano, B., Majewski, S. R., Hayes, C. R., et al. 2020, AJ, 160, 43

Baalmann, L. R., Scherer, K., Kleimann, J., et al. 2021, A&A, 650, A36

Cox, N. L. J., Kerschbaum, F., van Marle, A.-J., et al. 2012, A&A, 537, A35

Decin, L., Cox, N. L. J., Royer, P., et al. 2012, A&A, 548, A113

Everall, A., Belokurov, V., Evans, N. W., et al. 2022, MNRAS, 511, 3863

Falgarone, E., & Puget, J.-L. 1995, A&A, 293, 840

Ferrière, K. M., Zweibel, E. G., & Shull, J. M. 1988, ApJ, 332, 984

Frisch, P. C., Redfield, S., & Slavin, J. D. 2011, ARA&A, 49, 237

Gladstone, G. R., Pryor, W. R., Hall, D. T., et al. 2021, AJ, 162, 241

Gry, C., Boulanger, F., Nehmé, C., et al. 2002, A&A, 391, 675

Herbst, K., Scherer, K., Ferreira, S. E. S., et al. 2020, ApJ, 897, L26

Ingalls, J. G., Bania, T.M., Boulanger, F., et al. 2011, ApJ, 743, 174

Jensen, A. G., Snow, T. P., Sonneborn, G., & Rachford, B. L. 2010, ApJ, 711, 1236

Kobulnicky, H. A., Chick, W. T., Schurhammer, D. P., et al. 2016, ApJS, 227, 18

Kuhn, M. A., Hillenbrand, L. A., Sills, A., et al. 2019, ApJ, 870, 32

Kulkarni, S. R., Harrison, F. A., Grefenstette, B. W., et al. 2021, Science with the Ultraviolet Explorer (UVEX), arXiv:2111.15608

Kulkarni, S. R., & Shull, J. M. 2023, PASP, submitted (arXiv: 2302.13963)

Kulsrud, R. M. 2005, Plasma Physics for Astrophysics (Princeton: Princeton Univ. Press), p 108

Linsky, J. L., Redfield, S., & Tilipman, D. 2019, ApJ, 886, 41

Linsky, J. L., Redfield, S., Ryder, D., & Chasan-Taber, A. 2022, AJ, 164, 106

Mackey, J., Haworth, T. J., Gvaramadze, V. V., et al. 2016, A&A, 586, A114

Mac Low, M.-M., Van Buren, D., Wood, D. O. S., & Churchwell, E. 1991, ApJ, 369, 395

McComas, D. J., Alexashov, D., Bzowski, M., et al. 2012, Science, 336, 1291

McComas, D. J., Rankin, J. S., Schwadron, N. A., & Swaczyna, P. 2019, ApJ, 884, 145

Mihalas, D., & Binney, J. 2005, Galactic Astronomy, Chapter 4 (San Francisco: W. H. Freeman)

Moseley, E. R., Draine, B. T., Tomida, K., & Stone, J. M. 2021, MNRAS, 500, 3290

Myers, A. F., McKee, C. F., & Li, P. S. 2015, MNRAS, 453, 2747

Nehmé, C., Le Bourlot, J., Boulanger, F., et al. 2008, A&A, 483, 485

Nordström, B., Major, M., Andersen, J., et al. 2004, A&A, 418, 989

Peri, C. S., Benaglia, P., & Isequilla, N.L. 2015, A&A, 578, A45

Redfield, S., & Linsky, J. L. 2008, ApJ, 673, 283

Reylé, C., Jardine, K., Fouqué, P., et al. 2022, in 21st Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun, ed. A. S. Brun, J. Bouvier, P. Petit, arXiv:2302.02810

Richardson, J. D., Kasper, J. C., Wang, C., et al. 2008, Nature, 454, 63

Scherer, K., & Fichtner, H. 2014, ApJ, 782, 25

Shull, J. M., & McKee, C. F. 1979, ApJ, 227, 131

Shull, J. M., & Silk, J. 1979, ApJ, 234, 427

Shull, J. M., Danforth, C. W., & Anderson, K. L. 2021, ApJ, 911, 55

Shull, J. M., Darling, J., & Danforth, C. W. 2021, ApJ, 914, 18

Slavin, J. D., & Frisch, P. C. 2008, A&A, 491, 53

Spitzer, L. 1982, ApJ, 262, 315

Stone, E. C., Cummings, A. C., McDonald, F. B., et al. 2005, Science, 309, 2017

Stone, E. C., Cummings, A. C., McDonald, F. B., et al. 2008, Nature, 454, 71

van Buren, D., & McCray, R. 1988, ApJ, 329, L93

Vieira, K., Carraro, G., Korchagin, V., et al. 2022, ApJ, 932, 28

Weaver, R., McCray, R., Castor, J., Shapiro, P., & Moore, R. 1977, ApJ, 218, 377

Winston, E., Wolk, S.J., Bourke, T.L., et al. 2012, ApJ, 744, 126

Wood, B. E., Müller, H.-R., Redfield, S., et al. 2021, ApJ, 915, 37

Zank, G. P., Heerikhuisen, J., Wood, B. E., et al. 2013, ApJ, 763, 20

This preprint was prepared with the AAS LaTeX macros v5.2.

Table 1. Nearby Fast Stars[a]

Star	SpT	$V_{\rm ISM}$ (km/s)	$V_{\rm rad}$ (km/s)	d (pc)	$T_{\rm gas}$ (K)	$V_{\rm turb}$ (km/s)
61 Cyg A	K5 V	86	-65.2	3.50	6500	2.08
GJ 887	M2 V	85	+8.8	3.29	6000	1.65
GJ 436	M3 V	79	+9.6	9.78	...	...
GJ 205	M1.5 V	70	+8.6	5.70	4500	2.37
ϵ Ind	K5 V	68	-40.0	3.64	8340	1.97
61 Vir	G6.5 V	51	-7.8	8.53	6600	1.60
GJ 892	K3 V	49	-18.5	6.54	...	...
GJ 860A	M3 V	47	-24.0	4.01	4780	2.24
EV Lac	M3.5 V	45	+0.3	5.05	3030	2.21
π^1 UMa	G1.5 V	43	-12.8	14.4	2450	2.47
36 Oph A	K2 V	40	+0.4	5.95	5870	2.33
GJ 173	M1 V	38	-6.8	11.2	...	...
δ Eri	K0 IV	37	-6.2	9.09	3650	2.25
ξ Boo A	G7 V	32	+1.6	6.71	5310	1.68
GJ 338 A	M0 V	29	+11.5	6.33	6650	2.29
δ Pav	G8 IV	29	-21.5	6.10	9100	2.44
GJ 15 A	M2 V	28	+11.7	3.56	7800	2.81
ϵ Eri	K2 V	27	+16.4	3.22	7410	2.03
α Cen B	K1 V	25	-22.6	1.35	5500	1.37
Sun	G2 V	23	...	...	6400	...
Means					6638	2.54

[a] List of local "fast stars" with spectral types and Gaia parallax distances. $V_{\rm ISM}$ is the ISM flow speed seen by the star in the stellar rest frame (Wood et al. 2021), and $V_{\rm rad}$ is the star's heliocentric radial velocity. The last two columns list gas temperatures and turbulent velocities, inferred from an ensemble of absorption-line widths of hydrogen (both H I and D I) and various metal ions (Linsky et al. 2022).