Torsion-free nilpotent groups of small Hirsch length with isomorphic finite quotients

Alexander Cant and Bettina Eick

May 5, 2023

Abstract

Let $\mathcal{T}$ denote the class of finitely generated torsion-free nilpotent groups. For a group G let $F(G)$ be the set of isomorphism classes of finite quotients of G. Pickel proved that if $G \in \mathcal{T}$, then the set $\mathfrak{g}(G)$ of isomorphism classes of groups $H \in \mathcal{T}$ with $F(G) = F(H)$ is finite. We give an explicit description of the sets $\mathfrak{g}(G)$ for the $\mathcal{T}$-groups G of Hirsch length at most 5. Based on this, we show that for each Hirsch length $n \geq 4$ and for each $m \in \mathbb{N}$ there is a $\mathcal{T}$-group G of Hirsch length n with $|\mathfrak{g}(G)| \geq m$.

1 Introduction

Let G be a finitely generated torsion-free nilpotent group, $\mathcal{T}$-group for short, and write $F(G)$ for the set of isomorphism classes of finite quotients of G. The *genus* $\mathfrak{g}(G)$ of G (in the class of all $\mathcal{T}$-groups) is the set of isomorphism classes of $\mathcal{T}$-groups H with $F(G) = F(H)$. A purpose of this paper is to determine explicitly the genus $\mathfrak{g}(G)$ for each $\mathcal{T}$-group G of Hirsch length at most 5. This addresses the "Basic Problems" as introduced by Grunewald & Zalesskii [6] for $\mathcal{T}$-groups.

Pickel [15, 16] proved the interesting theorem that if G is a $\mathcal{T}$-group, then its genus $\mathfrak{g}(G)$ is finite. He obtains this result by showing that the elements in $\mathfrak{g}(G)$ correspond to a subset of the double cosets in a subgroup of $\prod_{p \text{ prime}} \mathrm{Aut}(\mathbb{Q}_p G)$, where $\mathbb{Q}_p G$ is the rational p-adic completion of G. While this allows to prove the desired finiteness, it does not yield many insights into the nature of the groups in $\mathfrak{g}(G)$.

Our results here yield an explicit description of the genera of the $\mathcal{T}$-groups of Hirsch length at most 5. This provides an illustration of Pickel's theorem [15, 16] and gives some insight into the nature of the groups in $\mathfrak{g}(G)$ for a $\mathcal{T}$-group G of Hirsch length at most 5. Our methods rely on the classification up to isomorphisms of the $\mathcal{T}$-groups of Hirsch length at most 5 by Eick & Engel [4].

Grunewald & Scharlau [8] proved that the genus of a $\mathcal{T}$-group of Hirsch length at most 5 and class at most 2 is trivial; that is, it consists of a single isomorphism class only. Thus the genus of a $\mathcal{T}$-group of Hirsch length at most 3 is trivial. Our results here imply the following, see Corollary 3 and Section 4.1 below.

1 Theorem: *For each $n \geq 4$ and each $m \in \mathbb{N}$ there exists a $\mathcal{T}$-group G of Hirsch length n whose genus $\mathfrak{g}(G)$ has more than m elements.*

Grunewald & Scharlau [8] proved a that for each $m \in \mathbb{N}$ there exists a $\mathcal{T}$-group G of Hirsch length 6 whose genus $\mathfrak{g}(G)$ has more than m elements. Theorem 1 extends this result to all possible Hirsch lengths. We also note that the groups used in the proof of Theorem 1 have class 3, while the groups in [8] have class 2.

Based on work by Pickel [17], Grunewald, Pickel & Segal [7] proved that the genus in the class of polycyclic-by-finite groups is finite. In contrast, the work by Nekrashevych [13] as well as by Nikolov & Segal [14] shows that there are classes of groups that allow uncountably large genera.

2 Results

Let G be a $\mathcal{T}$-group of Hirsch length n and class c. We denote with $G = \gamma_1(G) > \ldots > \gamma_c(G) > \gamma_{c+1}(G) = \{1\}$ the lower central series of G and for $1 \leq i \leq c+1$ we write $I_i(G)/\gamma_i(G)$ for the torsion subgroup of $G/\gamma_i(G)$. Then

$$G = I_1(G) > I_2(G) > \ldots > I_c(G) > I_{c+1}(G) = \{1\}$$

is the *isolator series* of G. It is a central series with torsion-free abelian quotients. The sequence $(d_1, \ldots, d_c)$ of ranks of these quotients is the *type* of G.

We refine the isolator series of G to a central series $G = G_1 > G_2 > \ldots > G_n > G_{n+1} = \{1\}$ whose quotients are infinite cyclic. For all i, let $g_i G_{i+1}$ be a generator of G_i/G_{i+1}. Then $(g_1, \ldots, g_n)$ is known as a *basis* of G. Note that every group element has a unique representation of the form $g_1^{x_1} \cdots g_n^{x_n}$ with integer exponents $x_1, \ldots, x_n$. Throughout the article, let $\mathbb{Z}(n)$ denote the set of all sequences $(t_{i,j,k} \in \mathbb{Z} \mid 1 \leq i < j < k \leq n)$. Each basis of G induces a finite presentation for G of the form

$$G(t) = \langle\, g_1, \ldots, g_n \mid [g_j, g_i] = g_{j+1}^{t_{i,j,j+1}} \cdots g_n^{t_{i,j,n}} \text{ for } 1 \leq i < j \leq n \,\rangle$$

for some $t \in \mathbb{Z}(n)$. As $n \leq 5$ in all our applications, we write t_{ijk} for $t_{i,j,k}$.

2.1 Type $(2,1,1)$

For $t \in \mathbb{Z}(4)$ define $d(t) = \gcd(t_{123}, t_{134})$. Further, we write

$$T(2,1,1) = \{\, t \in \mathbb{Z}(4) \mid 0 < t_{123}, t_{134};\ 0 \leq t_{124} \leq \tfrac{d(t)}{2};\ t_{234} = 0 \,\}.$$

Part (a) of the following theorem follows from the work of Eick & Engel [4] and part (b) is our first main result here.

2 Theorem:

(a) *For each $\mathcal{T}$-group G of type $(2,1,1)$ there exists a unique $t \in T(2,1,1)$ with $G \cong G(t)$.*

(b) *Let $s, t \in T(2,1,1)$. Then $F(G(s)) = F(G(t))$ if and only if*

- $s_{123} = t_{123}$ *and* $s_{134} = t_{134}$*, and*
- *for each maximal prime power p^k dividing $d(s)$ there exists $w_p \in \mathbb{Z}$ with $p \nmid w_p$ and $p^k \mid (t_{124} w_p - s_{124})$.*

For a prime p let $t_i \in T(2,1,1)$ be defined by $t_{i,123} = t_{i,134} = p$ and $t_{i,124} = i$. Then $\{G(t_i) \mid 1 \leq i \leq \lfloor \frac{p}{2} \rfloor\}$ consists of pairwise non-isomorphic groups with the same finite quotients and hence $|\mathfrak{g}(G(t_i))| \geq \lfloor \frac{p}{2} \rfloor$. In Section 4 we observe that if G is of type $(2,1,1)$ and A is free abelian of finite rank, then $|\mathfrak{g}(G \times A)| \geq |\mathfrak{g}(G)|$ follows. Hence we obtain the following which, in turn, proves Theorem 1.

3 Corollary: *For each $n \geq 2$, there is no finite upper bound to the size of the genus of a $\mathcal{T}$-group of type $(n,1,1)$.*

2.2 Type $(3,1,1)$

For $t \in \mathbb{Z}(5)$ define $d_1(t) = \gcd(t_{145}, t_{235})$ and $d_2(t) = \gcd(t_{124}, t_{135}, d_1(t))$. Further, we write

$$T(3,1,1) = \{\, t \in \mathbb{Z}(5) \mid 0 < t_{124}, t_{145};\ 0 \leq t_{235};\ 0 \leq t_{135} \leq \tfrac{d_1(t)}{2};\ 0 \leq t_{125} \leq \tfrac{d_2(t)}{2}; \\ t_{ijk} = 0 \text{ otherwise} \,\}.$$

Part (a) of the following theorem follows from the work of Eick & Engel [4] and part (b) is our second main result here.

4 Theorem:

(a) *For each $\mathcal{T}$-group G of type $(3,1,1)$ there exists a unique $t \in T(3,1,1)$ with $G \cong G(t)$.*

(b) *Let $s, t \in T(3,1,1)$. Then $F(G(s)) = F(G(t))$ if and only if*

- *$s_{ijk} = t_{ijk}$ for $(ijk) \in \{(124), (145), (235)\}$,*
- *for each maximal prime power p^k dividing $d_1(s)$ there exists $w_p \in \mathbb{Z}$ with $p \nmid w_p$ and $p^k \mid (t_{135} w_p - s_{135})$, and*
- *for each maximal prime power p^k dividing $d_2(s)$ there exists $v_p \in \mathbb{Z}$ with $p \nmid v_p$ and $p^k \mid (t_{125} v_p - s_{125})$.*

2.3 Type $(2,1,1,1)$

For $t \in \mathbb{Z}(5)$ define $d_3(t) = \gcd(t_{123}, t_{145}, t_{235})$. Consider the set

$$T_0(2,1,1,1) = \{\, t \in \mathbb{Z}(5) \mid 0 < t_{123}, t_{134}, t_{145};\ 0 \leq t_{124}, t_{125}, t_{135}, t_{235}; \\ t_{ijk} = 0 \text{ otherwise} \,\}.$$

We define a relation on $T_0(2,1,1,1)$ over a ring R that is either $\mathbb{Z}$ or $\mathbb{Z}_p$ via $s \sim_R t$ if $s_{ijk} = t_{ijk}$ for $(ijk) \in \{(123), (134), (145), (235)\}$ and there exist $u \in R^*$ and $v, w, x, y, z \in R$ so that

$$\begin{aligned} t_{124} u &= s_{124} - s_{123} w + s_{134} v, \\ t_{135} u &= s_{135} + s_{145} w + s_{134} x + s_{235} y, \text{ and} \\ t_{125} u^2 &= s_{125} + s_{135} v + s_{124} x + s_{134} v x + d_3(s) z. \end{aligned}$$

We note that this is an equivalence relation that corresponds to an action of the group $R^5 \rtimes R^*$ on $T_0(2,1,1,1)$. Let $T(2,1,1,1)$ be a set of representatives for $\sim_{\mathbb{Z}}$. Part (a) of the following theorem follows from the work of Eick & Engel [4] and part (b) is our third main result here.

5 Theorem:

(a) *For each $\mathcal{T}$-group G of type $(2,1,1,1)$ there exists a unique $t \in T(2,1,1,1)$ with $G \cong G(t)$.*

(b) *Let $s, t \in T(2,1,1,1)$. Then $F(G(s)) = F(G(t))$ if and only if $s \sim_{\mathbb{Z}_p} t$ for all primes p, or, equivalently, if and only if*

- *$s_{ijk} = t_{ijk}$ for $(ijk) \in \{(123),(134),(145),(235)\}$, and*
- *for each prime p there are $u, v, w, x, y, z \in \mathbb{Z}$ with $p \nmid u$ so that with p^k the maximal p-power dividing the product $s_{134}d_3(s)$ it follows that*

$$\begin{aligned} t_{124}u &\equiv s_{124} - s_{123}w + s_{134}v && \bmod p^k, \\ t_{135}u &\equiv s_{135} + s_{145}w + s_{134}x + s_{235}y && \bmod p^k, \quad \textit{and} \\ t_{125}u^2 &\equiv s_{125} + s_{135}v + s_{124}x + s_{134}vx + d_3(s)z && \bmod p^k. \end{aligned}$$

2.4 Type $(2,1,2)$

For $k \in \mathbb{N}$ and $R \in \{\mathbb{Z}, \mathbb{Z}_p\}$ let $D_k(R)$ denote the congruence subgroup

$$D_k(R) = \left\{ \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \,\middle|\, k \mid a_{12} \right\} \leq \mathrm{GL}(2, R).$$

For $a, b, c \in \mathbb{N}$ with $a \mid b$ define $L(a,b,c) = \{ (x,y) \in \mathbb{N}_0^2 \mid x < \gcd(a,c),\ y < \gcd(b,c) \}$. Then $L(a,b,c)$ is a finite set and the group $\mathrm{GL}(2,\mathbb{Z}_p)$ acts on this set via

$$(x,y) \mapsto (x,y)M \mod (\gcd(a,c), \gcd(b,c)),$$

where the mod-reduction is carried out independently in both components. Let $O(a,b,c)$ denote a set of orbit representatives for the action of $D_{b/a}(\mathbb{Z})$ on $L(a,b,c)$. We write

$$\begin{aligned} T(2,1,2) = \{ t \in \mathbb{Z}(5) \mid\ & 0 < t_{123}, t_{134}, t_{235};\ t_{134} \mid t_{235};\ (t_{124}, t_{125}) \in O(t_{134}, t_{235}, t_{123}); \\ & t_{ijk} = 0 \text{ otherwise} \}. \end{aligned}$$

Part (a) of the following theorem follows from the work of Eick & Engel [4] and part (b) is our fourth main result here.

6 Theorem:

(a) *For each $\mathcal{T}$-group G of type $(2,1,2)$ there exists a unique $t \in T(2,1,2)$ with $G \cong G(t)$.*

(b) *Let $s, t \in T(2,1,2)$ and $k = s_{235}/s_{134}$. Then $F(G(t)) = F(G(s))$ if and only if*

- *$t_{ijk} = s_{ijk}$ for $(ijk) \in \{(123),(134),(235)\}$, and*
- *for every prime p the tuples (s_{124}, s_{125}) and (t_{124}, t_{125}) are in the same orbit under the action of $D_k(\mathbb{Z}_p)$ on $L(t_{134}, t_{235}, t_{123})$.*

An explicit description of the orbits in Theorem 6(b) is given in Remark 14.

3 Methods

This section describes the principal ideas used to prove the above results. Let G be a $\mathcal{T}$-group with basis $(g_1, \ldots, g_n)$. For $x = (x_1, \ldots, x_n) \in \mathbb{Z}^n$ we write g^x as a short form of $g_1^{x_1} \cdots g_n^{x_n}$. The multiplication, inversion and powering in G can be represented by functions $f_j \colon \mathbb{Z}^n \times \mathbb{Z}^n \to \mathbb{Z}$ and $k_j \colon \mathbb{Z} \times \mathbb{Z}^n \to \mathbb{Z}$ defined via

$$g_1^{f_1(x,y)} \cdots g_n^{f_n(x,y)} = g^x g^y \quad \text{and} \quad g_1^{k_1(\ell,x)} \cdots g_n^{k_n(\ell,x)} = (g^x)^\ell.$$

Hall [9] proved that the functions $f_1, \ldots, f_n$ and $k_1, \ldots, k_n$ can be described by rational polynomials, nowadays called Hall polynomials. We note the following.

7 Lemma: *The Hall polynomials of a $\mathcal{T}$-group with respect to a fixed basis are unique.*

Proof: This follows directly from the observation that a multivariate rational polynomial $h \in \mathbb{Q}[z_1, \ldots, z_m]$ with $h(z) = 0$ for all $z = (z_1, \ldots, z_m) \in \mathbb{Z}^m$ is the zero polynomial. •

As a next step we extend the ring of exponents of group elements from $\mathbb{Z}$ to the rational numbers $\mathbb{Q}$, the p-adic integers $\mathbb{Z}_p$ or the p-adic rationals $\mathbb{Q}_p$. The following lemma provides a starting point for this.

8 Lemma: *Let $f \in \mathbb{Q}[z_1, \ldots, z_m]$ so that $f(a) \in \mathbb{Z}$ for all $a = (a_1, \ldots, a_m) \in \mathbb{Z}^m$ and let $R = \mathbb{Q}$, $\mathbb{Z}_p$ or $\mathbb{Q}_p$. Then $f(a) \in R$ for all $a \in R^m$.*

Proof: The cases $R = \mathbb{Q}$ and $R = \mathbb{Q}_p$ are obvious, since f has rational coefficients. It remains to consider the case $R = \mathbb{Z}_p$. It is well-known that a rational polynomial which takes integer values on integer inputs is a $\mathbb{Z}$-linear combination of polynomials of the form

$$f_{s_1,\ldots,s_m}(z_1, \ldots, z_m) := \binom{z_1}{s_1} \cdots \binom{z_m}{s_m},$$

where $s_1, \ldots, s_m \in \mathbb{N}_0$, $z_1, \ldots, z_m$ are indeterminates and

$$\binom{z_i}{s_i} = \frac{z_i(z_i - 1) \cdots (z_i - s_i + 1)}{s_i!}.$$

Clearly, $\binom{a}{s} \in \mathbb{Q}_p$ holds for $a \in \mathbb{Z}_p$ and $s \in \mathbb{N}_0$. By definition, a is the limit of a Cauchy sequence $(a_i)_{i \in \mathbb{N}}$ of non-negative integers with respect to the p-adic absolute value. Further $\binom{a_i}{s} \in \mathbb{Z}$ for all i. As polynomial functions are continuous and $\mathbb{Z}_p$ is closed and therefore complete, we obtain that $\binom{a}{s} \in \mathbb{Z}_p$. •

3.1 Completions

Let G be a $\mathcal{T}$-group with basis $(g_1, \ldots, g_n)$. Then $G = \{g^a \mid a \in \mathbb{Z}^n\}$. For $R = \mathbb{Q}$, $\mathbb{Z}_p$ or $\mathbb{Q}_p$ we define the set RG via

$$RG = \{g^a \mid a \in R^n\}.$$

The Hall polynomials $f_1, \ldots, f_n$ of G facilitate a multiplication on RG: For $g^a, g^b \in RG$ we define $g^a \cdot g^b := g^{f(a,b)} \in RG$, where $f(a,b) = (f_1(a,b), \ldots, f_n(a,b))$.

9 Lemma: *Let G be a $\mathcal{T}$-group and $R = \mathbb{Q}$, $\mathbb{Z}_p$ or $\mathbb{Q}_p$.*

(a) *RG equipped with the multiplication via Hall polynomials is a group.*

(b) *The isomorphism type of the group RG is independent of the chosen basis.*

Proof: (a) The multiplication is well-defined by Lemmas 7 and 8. It is not difficult to observe that an identity element and inverses are contained in RG. The proof of Lemma 7 implies that the multiplication is associative, as the multiplication in G is associative.
(b) Let $(g_1, \ldots, g_n)$ be a basis for G and let $(h_1, \ldots, h_n)$ be a basis for another $\mathcal{T}$-group H. Suppose that $\varphi \colon G \to H$ is an isomorphism. We show that φ extends to an isomorphism $RG \to RH$. Applying this with $G = H$ and two different bases then yields the desired result.

For $1 \leq i \leq n$ let $w_i = \varphi(g_i) = h^{r_i}$ with $r_i \in \mathbb{Z}^n$. The Hall polynomials $\overline{f}_1, \ldots, \overline{f}_n$ of H and the values $r_1, \ldots, r_n$ can be used to determine rational polynomials $m_1, \ldots, m_n$ satisfying that $\varphi(g^x) = w_1^{x_1} \cdots w_n^{x_n} = h_1^{m_1(x)} \cdots h_n^{m_n(x)}$ holds for all $x \in \mathbb{Z}^n$. Since φ is a group homomorphism, it follows that

$$h^{m(f(x,y))} = \varphi(g^{f(x,y)}) = \varphi(g^x g^y) = \varphi(g^x)\varphi(g^y) = h^{m(x)}h^{m(y)} = h^{\overline{f}(m(x),m(y))}$$

and thus $m(f(x,y)) = \overline{f}(m(x), m(y))$ for all $x, y \in \mathbb{Z}^n$. This equation extends to $x, y \in R^n$ and thus φ extends to a group homomorphism $\tilde{\varphi} \colon RG \to RH$. Similarly, the inverse of φ extends to a homomorphism $RH \to RG$, and this extension inverts $\tilde{\varphi}$. Thus $\tilde{\varphi}$ is an isomorphism. •

A group H is called $\mathbb{Q}$-*powered* if for every $h \in H$ and every $m \in \mathbb{N}$ there exists a unique $k \in H$ with $k^m = h$. A $\mathbb{Q}$-powered hull of a $\mathcal{T}$-group G is a $\mathbb{Q}$-powered group H containing G with the property that for every $h \in H$ there exists $m \in \mathbb{N}$ with $h^m \in G$. Malcev [12] observed that a $\mathbb{Q}$-powered hull of a $\mathcal{T}$-group exists and that it is unique up to isomorphism. The construction of $\mathbb{Q}G$ implies the following; its elementary proof is left to the reader.

10 Theorem: *Let G be a $\mathcal{T}$-group. Then $\mathbb{Q}G$ is a $\mathbb{Q}$-powered hull of G.*

The following theorem summarizes the relationship between properties of pairs of $\mathcal{T}$-groups and isomorphisms of their completions. Recall that groups G and H are *commensurable* if there are subgroups of finite index $\tilde{G} \leq G$ and $\tilde{H} \leq H$ with $\tilde{G} \cong \tilde{H}$. The Malcev-correspondence associates a rational Lie algebra $\Lambda(G)$ with a $\mathcal{T}$-group G. This can be constructed explicitly via the inverse Baker-Campbell-Hausdorff formula, see [15, Sec. 2] or [1] for details.

11 Theorem: *Let G and H be $\mathcal{T}$-groups.*

(a) $F(G) = F(H)$ *if and only if* $\mathbb{Z}_pG \cong \mathbb{Z}_pH$ *for all primes* p.

(b) *G and H are commensurable if and only if* $\mathbb{Q}G \cong \mathbb{Q}H$ *or, equivalently,* $\Lambda(G) \cong \Lambda(H)$.

(c) *If G and H are commensurable, then* $\mathbb{Z}_pG \cong \mathbb{Z}_pH$ *for almost all primes* p.

(d) *If* $\mathbb{Z}_pG \cong \mathbb{Z}_pH$ *for a prime p, then G and H have the same type.*

Proof: Pickel [15, 16] proved similar theorems, but using a different definition for $\mathbb{Z}_pG$: For a finitely generated nilpotent group G, Pickel defines $\mathbb{Z}_pG$ as the completion of G with respect to the uniform topology whose neighborhood basis of the identity consists of the groups $\{G^{p^i} \mid i \in \mathbb{N}\}$. This completion coincides with the inverse limit of the quotients $\{G/G^{p^i} \mid i \in \mathbb{N}\}$. If G is a $\mathcal{T}$-group, then Pickel's definition for $\mathbb{Z}_pG$ coincides with our definition, see Lemma 1.3 in [15]. Hence Pickel's results apply in our setting.
(a) This is Lemma 1.2 in [15] and it is attributed to Borel.
(b) This is often attributed to Malcev [12]. Alternatively, see [15, p. 330 & Sec. 2].
(c) Suppose that G and H are commensurable and let $\tilde{G} \leq G$ and $\tilde{H} \leq H$ both of finite index in G and H, respectively. If p is a prime with $p \nmid [G : \tilde{G}]$ and $p \nmid [H : \tilde{H}]$, then Lemma 1.8 in [15] implies that $\mathbb{Z}_pG = \mathbb{Z}_p\tilde{G} \cong \mathbb{Z}_p\tilde{H} = \mathbb{Z}_pH$.
(d) Suppose that $\varphi\colon \mathbb{Z}_pG \to \mathbb{Z}_pH$ is an isomorphism. Then $\varphi(\mathbb{Z}_pI_i(G)) = \mathbb{Z}_pI_i(H)$, since $\mathbb{Z}_p\gamma_i(G) = \gamma_i(\mathbb{Z}_pG)$ is invariant under isomorphisms and further $\mathbb{Z}_pI_i(G)/\mathbb{Z}_p\gamma_i(G) \cong \mathbb{Z}_p(I_i(G)/\gamma_i(G))$ is the torsion subgroup of $\mathbb{Z}_p(G/\gamma_i(G))$. Hence $\mathbb{Z}_p(I_i(G)/I_{i+1}(G)) \cong \mathbb{Z}_p(I_i(H)/I_{i+1}(H))$ for each $i \in \mathbb{N}$, and G and H have the same type. •

3.2 Isomorphisms

The following result is the foundation to our approach towards testing for isomorphic finite quotients. Eick & Engel [4, Lemma 7] used a similar result to test isomorphisms of $\mathcal{T}$-groups.

12 Theorem: *Let $t, s \in \mathbb{Z}(n)$ so that $G(t)$ and $G(s)$ are $\mathcal{T}$-groups of the same type $(d_1, \ldots, d_c)$ with bases $(g_1, \ldots, g_n)$ and $(h_1, \ldots, h_n)$, respectively.*

(a) *Let p be a prime and* $\varphi\colon \mathbb{Z}_pG(t) \to \mathbb{Z}_pG(s)$ *an isomorphism. Then* $\varphi(\mathbb{Z}_pI_i(G(t)) = \mathbb{Z}_pI_i(G(s))$ *for each* $i \in \mathbb{N}$.

(b) *Let p be a prime and* $\varphi\colon \mathbb{Z}_pG(t) \to \mathbb{Z}_pG(s)$ *an isomorphism. Then for* $1 \leq j \leq n$ *it follows that* $\varphi(g_j) = h^{m_j}$ *for some* $m_j = (m_{11}, \ldots, m_{1n}) \in \mathbb{Z}_p^n$ *and the matrix* $M = (m_{ij})_{1 \leq i,j \leq n}$ *is a block upper triangular matrix with diagonal blocks* $M_1, \ldots, M_c$ *satisfying* $M_i \in \mathrm{GL}(d_i, \mathbb{Z}_p)$.

(c) $F(G(t)) = F(G(s))$ *if and only if for all primes p there are* $m_1, \ldots, m_n \in \mathbb{Z}_p^n$ *so that the elements* $h^{m_1}, \ldots, h^{m_n} \in G(s)$ *satisfy the relations of* $G(t)$ *and the matrix* $M = (m_{ij})_{1 \leq i,j \leq n}$ *is invertible over* $\mathbb{Z}_p$.

Proof: (a) This is part of the proof of Theorem 11 (d).
(b) Let $\varphi\colon \mathbb{Z}_pG(t) \to \mathbb{Z}_pG(s)$ be an isomorphism. If $g_j \in I_i(G(t))$, then $\varphi(g_j) \in \mathbb{Z}_pI_i(G(s))$

by (a). Hence M is in block upper triangular form and the blocks on the diagonal correspond to the isomorphisms $\varphi_i\colon \mathbb{Z}_pI_i(G(t))/\mathbb{Z}_pI_{i+1}(G(t)) \to \mathbb{Z}_pI_i(G(s))/\mathbb{Z}_pI_{i+1}(G(s))$ induced by φ. Since $\mathbb{Z}_pI_i(G(t))/\mathbb{Z}_pI_{i+1}(G(t)) \cong \mathbb{Z}_p^{d_i} \cong \mathbb{Z}_pI_i(G(s))/\mathbb{Z}_pI_{i+1}(G(s))$, each isomorphism φ_i corresponds to an invertible $d_i \times d_i$-matrix over $\mathbb{Z}_p$.
(c) By (b) and Theorem 11 (a) it suffices to prove the following for an arbitrary prime p: If there are $m_1, \dots, m_n \in \mathbb{Z}_p^n$ so that the elements $h^{m_1}, \dots, h^{m_n}$ satisfy the relations of $G(t)$ and the matrix of exponents $M = (m_{ij})_{1 \leq i,j \leq n}$ is invertible over $\mathbb{Z}_p$, then $\mathbb{Z}_pG(t) \cong \mathbb{Z}_pG(s)$ follows.

Fix a prime p and assume that $m_1, \dots, m_n \in \mathbb{Z}_p^n$ exist. Let $\varphi\colon \{g_1, \dots, g_n\} \to \mathbb{Z}_pG(s)$, $g_i \mapsto h^{m_i}$. As the images $\varphi(g_i)$ satisfy the relations of $G(t)$, it follows that φ extends to an epimorphism $\tilde{\varphi}\colon G(t) \to H \leq \mathbb{Z}_pG(s)$, where H is the subgroup generated by $h_1^{m_1}, \dots, h_n^{m_n}$. As M is invertible, it induces isomorphisms $I_i(G(t))/I_{i+1}(G(t))$ onto $I_i(H)/I_{i+1}(H)$. This implies that $\tilde{\varphi}$ is injective. Lemma 9(b) now yields that $\tilde{\varphi}$ extends to an isomorphism $\mathbb{Z}_pG(t) \to \mathbb{Z}_pH$. Finally, M is invertible, hence $\mathbb{Z}_pH = \mathbb{Z}_pG(s)$. •

Our general approach to classify $\mathcal{T}$-groups up to isomorphic finite quotients is the following: Fix a type and consider indeterminates for s and t. Theorem 12(c) translates the condition $F(G(s)) = F(G(t))$ into a system of equations in s, t and the entries of a block upper triangular matrix M. It now remains to determine for which integral values of s and t this system of equations is solvable over $\mathbb{Z}_p$ for all primes p.

13 Lemma: *Let G be a $\mathcal{T}$-group and let $C_i(G) = C_G(I_i(G)/I_{i+2}(G))$ be a two-step centralizer. Then $C_i(G)$ is characteristic in G and $\mathbb{Z}_pC_i(G)$ is characteristic in $\mathbb{Z}_pG$.*

Proof: We note that $\mathbb{Z}_pC_i(G) = C_{\mathbb{Z}_pG}(\mathbb{Z}_pI_i(G)/\mathbb{Z}_pI_{i+2}(G))$. Now the result follows, since centralizers of characteristic subgroups are characteristic, see also Theorem 12(a). •

4 Proofs

This section contains the proofs for our results in Section 2. We use the parameterized Hall polynomials for $\mathcal{T}$-groups of Hirsch length at most 5, see [2], the solution to the isomorphism problem for these groups as described in [4] and the computer algebra system GAP [5] as starting points. Our GAP code is publically available at [3]. Further, for a prime p let $\nu_p\colon \mathbb{Z}_p \to \mathbb{N}_0 \cup \{\infty\}$ denote the p-adic valuation.

4.1 Proof of Theorem 1 and Corollary 3

Let G be a $\mathcal{T}$-group of type $(2,1,1)$ so that the genus $\mathfrak{g}(G)$ contains m pairwise non-isomorphic groups $H_1, \dots, H_m$ of type $(2,1,1)$, see Section 2.1, and let A be free abelian of rank ℓ. We show that $H_1 \times A, \dots, H_m \times A$ are pairwise non-isomorphic groups in $\mathfrak{g}(G \times A)$.

Consider $H \in \mathfrak{g}(G)$. Then $\mathbb{Z}_p(H \times A) = \mathbb{Z}_pH \times \mathbb{Z}_pA$ by construction for all primes p. This implies that $F(H \times A) = F(G \times A)$ by Theorem 11. Hence $H \times A \in \mathfrak{g}(G \times A)$.

The result that $H_i \times A \not\cong H_j \times A$ for $i \neq j$ follows from Hirshon [10], since H_i and H_j both are of type $(2,1,1)$ and thus have cyclic centers. •

4.2 $\mathcal{T}$-groups of type $(2, 1, 1)$

Proof of Theorem 2(b): Let $s, t \in T(2,1,1)$ and write $(g_1, \ldots, g_4)$ for a basis of $G(t)$ and $(h_1, \ldots, h_4)$ for a basis of $G(s)$. Then $C_2(G(t))$ satisfies $G(t) > C_2(G(t)) > I_2(G(t))$ with $G(t)/C_2(G(t))$ infinite cyclic and $C_2(G(t))/I_2(G(t))$ infinite cyclic, see [4, Lemma 8]. Using Theorem 12 and Lemma 13, we assume that an isomorphism $\mathbb{Z}_pG(t) \to \mathbb{Z}_pG(s)$ has the form

$$\begin{aligned} g_1 &\mapsto h_1^{m_{11}} h_2^{m_{12}} h_3^{m_{13}} h_4^{m_{14}} \\ g_2 &\mapsto h_2^{m_{22}} h_3^{m_{23}} h_4^{m_{24}} \\ g_3 &\mapsto h_3^{m_{33}} h_4^{m_{34}} \\ g_4 &\mapsto h_4^{m_{44}} \end{aligned}$$

where m_{ii} are invertible over $\mathbb{Z}_p$ and $m_{ij} \in \mathbb{Z}_p$. Evaluating the relations of $G(t)$ in $G(s)$ translates to the following equations. (This translation can best be done by computer, see [3] for GAP code for this purpose.)

(1) $t_{123}m_{11}^{-1}m_{22}^{-1}m_{33} - s_{123} = 0$

(2) $t_{134}m_{11}^{-1}m_{33}^{-1}m_{44} - s_{134} = 0$

(3) $t_{124}m_{11}^{-1}m_{22}^{-1}m_{44} - s_{124} = s_{123}(s_{134}(m_{11} - 1)/2 - m_{33}^{-1}m_{34}) + s_{134}m_{22}^{-1}m_{23}$

Equations (1–2) have solutions over $\mathbb{Z}_p$ for all primes p if and only if $s_{123} = t_{123}$ and $s_{134} = t_{134}$, since $m_{ii} \in \mathbb{Z}_p^*$. In this case $m_{33} = m_{11}m_{22}$ and $m_{44} = m_{11}^2 m_{22}$ follow and it remains to consider Equation (3). This has a solution over $\mathbb{Z}_p$ if and only if there are $x, y \in \mathbb{Z}_p$ and $u \in \mathbb{Z}_p^*$ with

$$t_{124}u - s_{124} = s_{123}x + s_{134}y.$$

We note that if x and y range over $\mathbb{Z}_p$, then $s_{123}x + s_{134}y$ ranges precisely over the $\mathbb{Z}_p$-multiples of $d(s) = \gcd(s_{123}, s_{134})$. Let $k = \nu_p(d(s))$. Then the $\mathbb{Z}_p$-multiples of $d(s)$ coincide with the $\mathbb{Z}_p$-multiples of p^k, since $d(s)/p^k$ is a unit in $\mathbb{Z}_p$.

Hence a solution of Equation (3) implies the existence of a $w_p \in \mathbb{Z}$ with $p \nmid w_p$ so that

$$t_{124}w_p - s_{124} \equiv 0 \bmod p^k.$$

Vice versa, such an integer w_p is a unit in $\mathbb{Z}_p$ and its existence shows that Equation (3) can be solved. •

4.3 $\mathcal{T}$-groups of type $(3, 1, 1)$

Proof of Theorem 4(b): Let $s, t \in T(3,1,1)$ and write $(g_1, \ldots, g_5)$ for the basis of $G(t)$ and $(h_1, \ldots, h_5)$ for the basis of $G(s)$. The presentation of $G(t)$ implies $C_1(G(t)) = \langle g_3, \ldots, g_5 \rangle$ and $C_2(G(t)) = \langle g_2, \ldots, g_5 \rangle$. Theorem 12 and Lemma 13 show that an isomorphism

$\mathbb{Z}_pG(t) \to \mathbb{Z}_pG(s)$ has the form

$$\begin{array}{rl}
g_1 \mapsto & h_1^{m_{11}} h_2^{m_{12}} h_3^{m_{13}} h_4^{m_{14}} h_5^{m_{15}} \\
g_2 \mapsto & h_2^{m_{22}} h_3^{m_{23}} h_4^{m_{24}} h_5^{m_{25}} \\
g_3 \mapsto & h_3^{m_{33}} h_4^{m_{34}} h_5^{m_{35}} \\
g_4 \mapsto & h_4^{m_{44}} h_5^{m_{45}} \\
g_5 \mapsto & h_5^{m_{55}}
\end{array}$$

where m_{ii} are invertible over $\mathbb{Z}_p$ and $m_{ij} \in \mathbb{Z}_p$. Evaluating the relations of $G(t)$ in $G(s)$ translates to the following equations.

$$\begin{array}{ll}
(1) & t_{124}m_{11}^{-1}m_{22}^{-1}m_{44} - s_{124} = 0 \\
(2) & t_{145}m_{11}^{-1}m_{44}^{-1}m_{55} - s_{145} = 0 \\
(3) & t_{235}m_{22}^{-1}m_{33}^{-1}m_{55} - s_{235} = 0 \\
(4) & t_{135}m_{11}^{-1}m_{33}^{-1}m_{55} - s_{135} = s_{145}m_{33}^{-1}m_{34} + s_{235}m_{11}^{-1}m_{12} \\
(5) & t_{125}m_{11}^{-1}m_{22}^{-1}m_{55} - s_{125} = s_{124}(s_{145}(m_{11} - 1)/2 - m_{44}^{-1}m_{45}) + s_{135}m_{22}^{-1}m_{23} \\
 & \qquad + s_{145}m_{22}^{-1}m_{24} + s_{235}(m_{11}^{-1}m_{22}^{-1}m_{12}m_{23} - m_{11}^{-1}m_{13})
\end{array}$$

Equations (1–3) have solutions over $\mathbb{Z}_p$ for all primes p if and only if $s_{ijk} = t_{ijk}$ for $(ijk) \in \{(124), (145), (235)\}$. In this case $m_{44} = m_{11}m_{22}$ and $m_{55} = m_{11}^2 m_{22}$ follow. If $s_{235} > 0$, then $m_{33} = m_{11}^2$ follows additionally.

It remains to consider Equations (4–5). Note that $m_{11}^{-1}m_{33}^{-1}m_{55} = m_{11}m_{22}m_{33}^{-1}$ and $m_{11}^{-1}m_{22}^{-1}m_{55} = m_{11}$ can take arbitrary values in $\mathbb{Z}_p^*$ independently. Further note that the right hand sides of Equations (4–5) independently range over all $\mathbb{Z}_p$-multiples of $d_1(s) = \gcd(s_{145}, s_{235})$ and $d_2(s) = \gcd(s_{124}, s_{135}, d_1(s))$, respectively.

Fix a prime p and let $k_1 = \nu_p(d_1(s))$ and $k_2 = \nu_p(d_2(s))$. Similar to our proof of Theorem 2 (b), we deduce that Equations (4–5) have a solution if and only if there are $w_p, v_p \in \mathbb{Z} \setminus p\mathbb{Z}$ so that $p^{k_1} \mid (t_{135}w_p - s_{135})$ and $p^{k_2} \mid (t_{125}v_p - s_{125})$ hold. •

4.4 $\mathcal{T}$-groups of type $(2, 1, 1, 1)$

Proof of Theorem 5(b): Let $s, t \in T(2, 1, 1, 1)$ and write $(g_1, \ldots, g_5)$ for the basis of $G(t)$ and $(h_1, \ldots, h_5)$ for the basis of $G(s)$. The presentation of $G(t)$ implies $C_3(G(t)) = \langle g_2, \ldots, g_5\rangle$. Thus an isomorphism $\mathbb{Z}_pG(t) \to \mathbb{Z}_pG(s)$ has the same general form as for type $(3, 1, 1)$, see also Theorem 12 and Lemma 13. Evaluating the relations of $G(t)$ in the images of an isomorphism translates to 7 equations. We define

$$\begin{aligned}
v &= s_{123}(m_{11}-1) + m_{22}^{-1}m_{23}\\
w &= s_{134}(m_{11}-1)/2 + m_{33}^{-1}m_{34}\\
x &= -m_{44}^{-1}m_{45}\\
y &= m_{11}^{-1}m_{12}\\
z_1 &= s_{134}s_{145}(m_{11}-1)(m_{11}-2)/6 + s_{134}m_{44}^{-1}m_{45}(m_{11}-1)/2 - s_{135}(m_{11}-1)/2\\
&\quad + s_{235}m_{12} + s_{235}(m_{22}-1)/2 + m_{33}^{-1}(m_{44}^{-1}m_{34}m_{45} - m_{35})\\
z_2 &= s_{124}(m_{11}-1)/2 + s_{134}m_{22}^{-1}m_{23}(m_{11}-1)/2 + m_{22}^{-1}m_{24}\\
z_3 &= m_{11}^{-1}(m_{22}^{-1}m_{12}m_{23} - m_{13})
\end{aligned}$$

and, based on this, obtain the equations

$$\begin{aligned}
&(1)\quad t_{123}m_{11}^{-1}m_{22}^{-1}m_{33} - s_{123} = 0\\
&(2)\quad t_{134}m_{11}^{-1}m_{33}^{-1}m_{44} - s_{134} = 0\\
&(3)\quad t_{145}m_{11}^{-1}m_{44}^{-1}m_{55} - s_{145} = 0\\
&(4)\quad t_{235}m_{22}^{-1}m_{33}^{-1}m_{55} - s_{235} = 0\\
&(5)\quad t_{124}m_{11}^{-1}m_{22}^{-1}m_{44} - s_{124} = s_{134}v - s_{123}w\\
&(6)\quad t_{135}m_{11}^{-1}m_{33}^{-1}m_{55} - s_{135} = s_{134}x + s_{145}w + s_{235}y\\
&(7)\quad t_{125}m_{11}^{-1}m_{22}^{-1}m_{55} - s_{125} = s_{124}x + s_{134}vx + s_{135}v + s_{123}z_1 + s_{145}z_2 + s_{235}z_3
\end{aligned}$$

Equations (1–4) have solutions over $\mathbb{Z}_p$ for all primes p if and only if $s_{ijk} = t_{ijk}$ holds for $(ijk) \in \{(123), (134), (145), (235)\}$. In this case $m_{33} = m_{11}m_{22}$, $m_{44} = m_{11}^2 m_{22}$ and $m_{55} = m_{11}^3 m_{22}$ follow. If $s_{235} > 0$, then $m_{22} = m_{11}^2$ follows additionally.

It remains to consider Equations (5–7). Write $u = m_{11}$ to shorten notation and note that $u = m_{11}^{-1}m_{22}^{-1}m_{44} = m_{11}^{-1}m_{33}^{-1}m_{55}$ and $u^2 = m_{11}^{-1}m_{22}^{-1}m_{55}$. Next, note that if all m_{ij} range independently over $\mathbb{Z}_p$, then so do v, w, x, y, z_1, z_2 and z_3. Further, $s_{123}z_1 + s_{145}z_2 + s_{235}z_3$ ranges over all $\mathbb{Z}_p$-multiples of $d_3(s) = \gcd(s_{123}, s_{145}, s_{235})$.

Fix a prime p. Then Equations (5–7) have a solution over $\mathbb{Z}_p$ if and only if there are $u \in \mathbb{Z}_p^*$ and $v, w, x, y, z \in \mathbb{Z}_p$ with

$$\begin{aligned}
&\text{(i)} &\quad s_{134}v &= t_{124}u - s_{124} + s_{123}w,\\
&\text{(ii)} &\quad s_{134}x &= t_{135}u - s_{135} - s_{145}w - s_{235}y, \quad\text{and}\\
&\text{(iii)} &\quad d_3(s)z &= t_{125}u^2 - s_{125} - s_{124}x - s_{134}vx - s_{135}v.
\end{aligned}$$

If Equations (i–iii) have a solution, then they can be solved modulo any power of p by $u, v, w, x, y, z \in \mathbb{Z}$ with $p \nmid u$. Vice versa, assume that $u, v, w, x, y, z \in \mathbb{Z}$ with $p \nmid u$ solve Equations (i–iii) modulo $p^k = p^{\alpha+\beta}$, where $\alpha = \nu_p(d_3(s))$ and $\beta = \nu_p(s_{134})$. Then $u \in \mathbb{Z}_p^*$ and there are $A, B, C, D \in \mathbb{Z}$ with

$$\begin{aligned}
s_{134}v + Ap^{\alpha+\beta} &= t_{124}u - s_{124} + s_{123}w,\\
s_{134}x + Bp^{\alpha+\beta} &= t_{135}u - s_{135} - s_{145}w - s_{235}y \quad\text{and}\\
Dp^{\alpha} = d_3(s)z + Cp^{\alpha+\beta} &= t_{125}u^2 - s_{125} - s_{124}x - s_{134}vx - s_{135}v.
\end{aligned}$$

It follows that $\tilde{u} = u$, $\tilde{w} = w$, $\tilde{y} = y$, $\tilde{v} = v + Ap^{\alpha}(s_{134}/p^{\beta})^{-1}$ and $\tilde{x} = x + Bp^{\alpha}(s_{134}/p^{\beta})^{-1}$ solve Equations (i–ii). If we insert these values in Equation (iii), then the right hand side translates to

$$\Big(D - \big(s_{124}B + s_{134}(vB + xA + ABp^{\alpha}(s_{134}/p^{\beta})^{-1}) + s_{135}A\big)(s_{134}/p^{\beta})^{-1}\Big)p^{\alpha} =: Ep^{\alpha},$$

so $\tilde{z} = E(d_3(s)/p^{\alpha})^{-1}$ yields a solution to Equation (iii). •

4.5 $\mathcal{T}$-groups of type $(2, 1, 2)$

Proof of Theorem 6(b): Let $s, t \in T(2, 1, 2)$ and $k = s_{235}/s_{134} \in \mathbb{Z}$. Write $(g_1, \ldots, g_5)$ for the basis of $G(t)$ and $(h_1, \ldots, h_5)$ for the basis of $G(s)$. Theorem 12 and Lemma 13 show that an isomorphism $\mathbb{Z}_pG(t) \to \mathbb{Z}_pG(s)$ has the form

$$\begin{aligned}
g_1 &\mapsto h_1^{m_{11}} h_2^{m_{12}} h_3^{m_{13}} h_4^{m_{14}} h_5^{m_{15}} \\
g_2 &\mapsto h_1^{m_{21}} h_2^{m_{22}} h_3^{m_{23}} h_4^{m_{24}} h_5^{m_{25}} \\
g_3 &\mapsto h_3^{m_{33}} h_4^{m_{34}} h_5^{m_{35}} \\
g_4 &\mapsto h_4^{m_{44}} h_5^{m_{45}} \\
g_5 &\mapsto h_4^{m_{54}} h_5^{m_{55}}
\end{aligned}$$

where $m_{ij} \in \mathbb{Z}_p$ with $m_{33} \in \mathbb{Z}_p^*$ and

$$M_1 = \begin{pmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{pmatrix}, \; M_3 = \begin{pmatrix} m_{44} & m_{45} \\ m_{54} & m_{55} \end{pmatrix} \in \mathrm{GL}(2, \mathbb{Z}_p).$$

Evaluating the relations of $G(t)$ in $G(s)$ translates to the following (matrix) equations.

$$\begin{aligned}
&(1) && t_{123}m_{33} = s_{123}\det(M_1) \\
&(2) && \mathrm{diag}(t_{134}, t_{235})M_3 = m_{33}M_1\,\mathrm{diag}(s_{134}, s_{235}) \\
&(3) && (t_{124}, t_{125})M_3 - (s_{124}, s_{125})\det(M_1) = (m_{23}, -m_{13})M_1\,\mathrm{diag}(s_{134}, s_{235}) + s_{123}(x, y)
\end{aligned}$$

In Equation (3) we used the following definitions.

$$\begin{aligned}
x &= s_{134}(m_{11}^2m_{22} - m_{12}m_{21}^2 - \det(M_1))/2 - \det(M_1)m_{33}^{-1}m_{34} \\
y &= s_{235}(m_{11}m_{22}^2 - m_{12}^2m_{21} - \det(M_1) + 2m_{12}m_{22}(m_{11} - m_{21}))/2 - \det(M_1)m_{33}^{-1}m_{35}
\end{aligned}$$

Equation (1) implies that $s_{123} = t_{123}$ and $m_{33} = \det(M_1)$. If Equation (2) has a solution over $\mathbb{Z}_p$ for a prime p, then the diagonal matrices $\mathrm{diag}(s_{134}, s_{235})$ and $\mathrm{diag}(t_{134}, t_{235})$ have the same elementary divisors over $\mathbb{Z}_p$. Thus $\nu_p(s_{134}) = \nu_p(t_{134})$ and $\nu_p(s_{235}) = \nu_p(t_{235})$; see also [11, Sec. IV, §7]. As a consequence, Equation (2) has a solution for all primes p if and only if $s_{134} = t_{134}$ and $s_{235} = t_{235}$. In this case, Equation (2) implies $m_{45} = km_{12}m_{33}$ and thus $M_3 \in D_k(\mathbb{Z}_p)$. We now fix a prime p for the remainder of the proof.

If all m_{ij} range independently over $\mathbb{Z}_p$, then so do x and y. Further $(m_{23}, -m_{13})M_1$ ranges over all of $\mathbb{Z}_p^2$, since we assume that M_1 is invertible. It follows that the right hand side of Equation (3) takes arbitrary values of the form $(a\gcd(s_{123}, s_{134}), b\gcd(s_{123}, s_{235}))$

with $a, b \in \mathbb{Z}_p$, and thus (s_{124}, s_{125}) and (t_{124}, t_{125}) are in the same orbit under the action of $D_k(\mathbb{Z}_p)$ on $L(s_{134}, s_{235}, s_{123})$.

Vice versa, assume that $(t_{124}, t_{125})A \equiv (s_{124}, s_{125}) \bmod (\gcd(s_{123}, s_{134}), \gcd(s_{123}, s_{235}))$ for some $A \in D_k(\mathbb{Z}_p)$. It suffices to determine $M_1, M_3 \in \mathrm{GL}(2, \mathbb{Z}_p)$ with $\det(M_1)^{-1}M_3 = A$ that solve Equation (2). Note that if M_1 and M_3 solve Equation (2), then $\det(M_3) = \det(M_1)^3$ holds, and thus $\det(M_1) = \det(A)$ and $M_3 = \det(A)A$. This yields a solution to Equations (1–3) over $\mathbb{Z}_p$. •

14 *Remark:* Fix integers $a, b, c > 0$ with $a \mid b$ and a prime p, and let $k = b/a$. Let us give an explicit description of the orbits of $D_k(\mathbb{Z}_p)$ on $L(a, b, c)$. Let $(d, e), (f, g) \in L(a, b, c)$ and write $\ell = \nu_p(\gcd(a, c))$ and $m = \nu_p(\gcd(b, c))$.

If there is a matrix $A \in D_k(\mathbb{Z}_p)$ with $(d, e)A \equiv (f, g) \mod (\gcd(a, c), \gcd(b, c))$, then there are $\alpha, \beta, \gamma, \delta \in \mathbb{Z}$ with

$$p^\ell \mid (d\alpha + e\gamma - f), \quad p^m \mid (dk\beta + e\delta - g) \quad \text{and} \quad p \nmid (\alpha\delta - k\beta\gamma).$$

The other way round, such integers $\alpha, \beta, \gamma, \delta$ immediately yield a matrix

$$A = \begin{pmatrix} \alpha & k\beta \\ \gamma & \delta \end{pmatrix} \in D_k(\mathbb{Z}_p)$$

with $(d, e)A \equiv (f, g) \mod (\gcd(a, c), \gcd(b, c))$.

References

[1] B. Assmann and S. Linton. Using the Malcev correspondence for collection in polycyclic groups. *J. Algebra*, 316(2):828–848, 2007.

[2] A. Cant and B. Eick. Polynomials describing the multiplication in finitely generated torsion-free nilpotent groups. *J. Symbolic Comput.*, 92:203–210, 2019.

[3] A. Cant and B. Eick. GAP code for the computation of equations related to isomorphisms between p-adic completions of finitely generated torsion-free nilpotent groups. `https://www.tu-braunschweig.de/en/iaa/personal/cant`, 2023.

[4] B. Eick and A.-K. Engel. The isomorphism problem for torsion free nilpotent groups of Hirsch length at most 5. *Groups Complex. Cryptol.*, 9(1):55–75, 2017.

[5] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.12.2*, 2022.

[6] F. Grunewald and P. Zalesskii. Genus for groups. *J. Algebra*, 326(1):130–168, 2011.

[7] F. J. Grunewald, P. F. Pickel, and D. Segal. Polycyclic groups with isomorphic finite quotients. *Ann. Math. (2)*, 111:155–195, 1980.

[8] F. J. Grunewald and R. Scharlau. A note on finitely generated torsion-free nilpotent groups of class 2. *J. Algebra*, 58(1):162–175, 1979.

[9] P. Hall. *The Edmonton notes on nilpotent groups.* Queen Mary College Mathematics Notes. Queen Mary College, Mathematics Department, London, 1969.

[10] R. Hirshon. Some cancellation theorems with applications to nilpotent groups. *J. Austral. Math. Soc. Ser. A*, 23(2):147–165, 1977.

[11] H. Lombardi and C. Quitté. *Commutative Algebra: Constructive Methods.* Springer Netherlands, 2015.

[12] A. I. Malcev. On a class of homogeneous spaces. *Amer. Math. Soc. Translation*, 1951(39):33, 1951.

[13] V. Nekrashevych. An uncountable family of 3-generated groups with isomorphic profinite completions. *Internat. J. Algebra Comput.*, 24(1):33–46, 2014.

[14] N. Nikolov and D. Segal. Constructing uncountably many groups with the same profinite completion. *New Zealand J. Math.*, 52:765–771, 2021 [2021–2022].

[15] P. F. Pickel. Finitely generated nilpotent groups with isomorphic finite quotients. *Trans. Amer. Math. Soc.*, 160:327–341, 1971.

[16] P. F. Pickel. Finitely generated nilpotent groups with isomorphic finite quotients. *Bull. Amer. Math. Soc.*, 77:216–219, 1971.

[17] P. F. Pickel. Nilpotent-by-finite groups with isomorphic finite quotients. *Trans. Amer. Math. Soc.*, 183:313–325, 1973.