Average edge order of normal 3-pseudomanifolds

Biplab Basak and Raju Kumar Gupta

Department of Mathematics, Indian Institute of Technology Delhi, New Delhi 110016, India[1].

June 20, 2024

Abstract

In their work [8], Feng Luo and Richard Stong introduced the concept of the average edge order, denoted as $\mu_0(K)$. They demonstrated that if $\mu_0(K) \leq \frac{9}{2}$ for a closed 3-manifold K, then K must be a sphere. Building upon this foundation, Makoto Tamura extended similar results to 3-manifolds with non-empty boundaries in [10, 11]. In our present study, we extend these findings to normal 3-pseudomanifolds. Specifically, we establish that for a normal 3-pseudomanifold K with singularities, $\mu_0(K) \geq \frac{30}{7}$. Moreover, equality holds if and only if K is a one-vertex suspension of $\mathbb{RP}^2$ with seven vertices. Furthermore, we establish that when $\frac{30}{7} \leq \mu_0(K) \leq \frac{9}{2}$, the 3-pseudomanifold K can be derived from some boundary complexes of 4-simplices by a sequence of possible operations, including connected sums, bistellar 1-moves, edge contractions, edge expansions, vertex folding, and edge folding. In the end, we discuss some normal 3-pseudomanifolds exhibiting higher average edge orders.

MSC 2020 : Primary 05E45; Secondary 05C07, 05A20.
Keywords: Normal pseudomanifolds; Average edge order; Suspension.

1 Introduction

Consider a normal 3-pseudomanifold denoted as K. Its *average edge order*, denoted by $\mu_0(K)$, is defined as $\mu_0(K) = \frac{3F(K)}{E(K)}$, where $F(K)$ and $E(K)$ represent the number of faces and edges in K respectively. Essentially, $\mu_0(K)$ signifies the average number of faces incident on edges within K. In their research [8], Feng Luo and Richard Stong demonstrated that a small average edge order in a closed connected 3-manifold suggests relatively simple topology and imposes constraints on its triangulation. The implication of this finding can be summarized as:

---

[1] *E-mail addresses:* `biplab@iitd.ac.in` (B. Basak), `Raju.Kumar.Gupta@maths.iitd.ac.in` (R. K. Gupta).

Proposition 1 ([8])**.** *Let K be any triangulation of a closed connected 3-manifold M. Then*

- *(i)* $3 \leq \mu_0(K) < 6$, *equality holds if and only if K is the triangulation of the boundary of a 4-simplex.*
- *(ii) For any $e > 0$, there are triangulations K_1 and K_2 of M such that $\mu_0(K_1) < 4.5 + e$ and $\mu_0(K_2) > 6 - e$.*
- *(iii) If $\mu_0(K) < 4.5$, then K is a triangulation of $\mathbb{S}^3$.*
- *(iv) If $\mu_0(K) = 4.5$, then K is a triangulation of $\mathbb{S}^3$, $\mathbb{S}^2 \times \mathbb{S}^1$, or $\mathbb{S}^2 \tilde{\times} \mathbb{S}^1$. Furthermore, in the last two cases the triangulations can be described.*

Building upon the previous discussion, Makoto Tamura contributed further insights in [10] by extending similar results to 3-manifolds with non-empty boundaries. In [11], Tamura revised the definition of the average edge order as follows: $\mu_0(K) = 3\frac{F_0(K)}{E_0(K)}$, where $E_0(K) = E_i(K) + \frac{E_\partial(K)}{2}$ and $F_0(K) = F_i(K) + \frac{F_\partial(K)}{2}$. Here, $E_i(K)$ (respectively, $F_i(K)$) represents the number of edges (respectively, faces) within int $K = K \setminus \partial(K)$, while $E_\partial(K)$ (respectively, $F_\partial(K)$) denotes the number of edges (respectively, faces) within $\partial(K)$. With this refined definition of the average edge order, Tamura established the following theorem:

Proposition 2 ([11], Theorem 1.2)**.** *Let K be any triangulation of a compact connected 3-manifold M with non-empty boundary. Then*

- *(i)* $2 \leq \mu_0(K) < 6$, *and equality holds if and only if K is the triangulation of one 3-simplex.*
- *(ii) For any rational number r with $4 < r < 6$, there is a triangulation K' of M such that $\mu_0(K') = r$.*
- *(iii) If $\mu_0(K) < 4$, then K is a triangulation of $\mathbb{B}^3$. There are an infinite number of distinct such triangulations, but for any constant $c < 4$ there are only finitely many triangulations K with $\mu_0(K) \leq c$.*
- *(iv) If $\mu_0(K) = 4$, then K is a triangulation of $\mathbb{B}^3, \mathbb{D}^2 \times \mathbb{S}^1$ or $\mathbb{D}^2 \tilde{\times} \mathbb{S}^1$. Furthermore, in the last two cases the triangulations can be described.*

In [4], similar work is done for a 3-manifold colored triangulation. In this article, we establish comparable conclusions for normal 3- pseudomanifolds without boundaries. In fact, we prove the following result:

Theorem 3. *Let K be a normal 3-pseudomanifold with singularities, and let $\mu_0(K)$ represent the average edge order of K. Then*

- *(i)* $\mu_0(K) \geq \frac{30}{7}$, *and equality holds if and only if K is a one-vertex suspension of $\mathbb{RP}^2$ with seven vertices.*

(ii) *If* $\mu_0(K) \leq \frac{9}{2}$, *then* K *is obtained from some boundary complexes of* 4*-simplices by a sequence of possible operations, including connected sums, bistellar* 1*-moves, edge contractions, edge expansions, vertex folding, and edge folding. In particular, either* $|K|$ *is a handlebody with its boundary coned off or a suspension of* $\mathbb{RP}^2$.

(iii) *If* K *contains exactly* n *singular vertices, then* $\mu_0(K) < 6 + n$. *Additionally, a sequence of normal* 3*-pseudomanifolds, denoted as* $\{K_m\}_{m\geq 1}$, *with two singularities exists, such that* $\mu_0(K_m) \to 8$ *as* $m \to \infty$.

2 Preliminaries

A *d-simplex* is the convex hull of $d+1$ affinely independent points. A 0-simplex is a point, a 1-simplex is an edge, a 2-simplex is a triangle, and so on. Let σ be a simplex. Then, *faces* of σ are defined as the convex hulls of non-empty subsets of the vertex set of σ. We denote a face τ of σ as $\tau \leq \sigma$. A *simplicial complex* K is a finite collection of simplices in $\mathbb{R}^m$ for some $m \in \mathbb{N}$ that satisfies the following conditions: if $\sigma \in K$, then $\tau \leq \sigma$ implies $\tau \in K$; for any two faces σ and τ in K, either $\sigma \cap \tau = \emptyset$ or $\sigma \cap \tau \leq \sigma$ and $\sigma \cap \tau \leq \tau$. We assume that the *empty simplex* $\emptyset$ (considered as a simplex of dimension -1) is a member of every simplicial complex. If K is a simplicial complex, then its *geometric carrier* $|K| = \cup_{\sigma \in K}\sigma$ is the union of all the simplices in K together with the subspace topology induced from $\mathbb{R}^n$ for some $n \in \mathbb{N}$. The *dimension* of K is defined as the maximum of the dimension of simplices in K. If σ and τ are two skew simplices in $\mathbb{R}^n$ for some $n \in \mathbb{N}$, then their *join* $\sigma * \tau$ is defined as $\{\lambda p + \mu q \,|\, p \in \sigma, q \in \tau;\ \lambda, \mu \in [0,1] \text{ and } \lambda + \mu = 1\}$. Similarly, we define the *join* of two skew simplicial complexes K_1 and K_2 as $\{\sigma * \tau | \sigma \in K_1, \tau \in K_2\}$. The *link* of a face σ in a simplicial complex K is defined as $\{\tau \in K \,|\, \sigma \cap \tau = \emptyset, \tau * \sigma \in K\}$. The link of σ is denoted by $\mathrm{lk}\,(\sigma, K)$. The *star* of a face σ in a simplicial complex K is defined as $\{\beta | \beta \leq \sigma * \alpha \text{ and } \alpha \in \mathrm{lk}\,(\sigma, K)\}$, and is denoted by $\mathrm{st}\,(\sigma, K)$.

A simplicial complex K is called *pure* if all its facets possess the same dimension. A simplicial complex K is called a *normal d-pseudomanifold without a boundary* if it is pure, every face of dimension $d-1$ is contained in precisely two facets, and the link of every face of dimension $\leq d-2$ is connected. If some faces of dimension $d-1$ are contained in only one facet, then K is called a *normal d-pseudomanifold with boundary.* For a normal 3-pseudomanifold K, the link of a vertex v is a triangulated closed connected surface. If $|\mathrm{lk}\,(v)| \cong \mathbb{S}^2$, then we call v a non-singular vertex; otherwise, v is called a singular vertex. Let K be a simplicial complex. Let $V(K)$, $E(K)$, $F(K)$, and $T(K)$ represent the counts of vertices, edges, faces, and tetrahedra in K, respectively. In this article, we will use the abbreviations V, E, F, and T to refer to $V(K)$, $E(K)$, $F(K)$, and $T(K)$ when the context is clear that we are discussing simplicial complex K. The numbers g_2 and g_3 for a 3-dimensional simplicial complex K are defined as $g_2(K) = E - 4V + 10$, and $g_3(K) = F - 3E + 6V - 10$.

Remark 4. Let $\{[1,2,4]$, $[1,3,4]$, $[1,2,5]$, $[1,3,6]$, $[1,5,6]$, $[2,4,6]$, $[2,3,5]$, $[2,3,6]$, $[3,4,5]$, $[4,5,6]\}$ denote a set of facets comprising the triangulation L of $\mathbb{RP}^2$ with six vertices. Consider $K = (uv * \mathrm{lk}\,(6)) \cup (u * L \setminus \{6\}) \cup (v * L \setminus \{6\})$. We define K as the one-vertex suspension of the complex L. Topologically, $|K|$ represents the suspension of $|L|$. Consequently, $\{[1,2,4,u]$, $[1,3,4,u]$, $[1,2,5,u]$, $[1,2,4,v]$, $[1,3,4,v]$, $[1,2,5,v]$, $[1,3,u,v]$, $[1,5,u,v]$, $[2,4,u,v]$, $[2,3,5,u]$, $[2,3,5,v]$, $[2,3,u,v]$, $[3,4,5,u]$, $[3,4,5,v]$, $[4,5,u,v]\}$ forms the set of facets of K.

Proposition 5 (Lemma 2.6, [2])**.** *Let K be a normal 3-pseudomanifold, and $v \in K$ be a vertex in K. Then $g_2(K) \geq g_2(\mathrm{st}\,(v, K)) = g_2(\mathrm{lk}\,(v, K))$.*

For definitions of operations like connected sum, bistellar 1-move, edge contraction, edge expansion, vertex folding, and edge folding, please see [2]. The subsequent results can be obtained by substituting $n = 1$ into [2, Theorem 4.4] and $m = 1$ into [2, Theorem 5.5].

Proposition 6. *Let K be a normal 3-pseudomanifold with exactly one singularity at t, such that $|\mathrm{lk}\,(t, K)|$ is a torus or Klein bottle. Then, $g_2(K) \leq 15$ implies that K is obtained from some boundary complexes of 4-simplices by a sequence of operations of types connected sums, bistellar 1-moves, edge contractions, edge expansions, and a vertex folding. Furthermore, $|K|$ is a handlebody with its boundary coned off.*

Proposition 7. *Let K be a normal 3-pseudomanifold with exactly two $\mathbb{RP}^2$ singularities. Then, $g_2(K) \leq 12$ implies K is obtained from some boundary complexes of 4-simplices by a sequence of operations of types connected sums, bistellar 1-moves, edge contractions, edge expansions, and an edge folding.*

Remark 8. If K is a normal 3-pseudomanifold and K^{ψ}_{uv} is obtained from K by an edge folding at uv, then $g_2(K^{\psi}_{uv}) = g_2(K) + 3$. Furthermore, if K is a triangulation of a 3-sphere then K^{ψ}_{uv} is a triangulation of the suspension of $\mathbb{RP}^2$ because $|K^{\psi}_{uv}|$ is obtained from a 3-ball by the following procedure: (i) choose two connected regions in the boundary of the 3-ball having a point in common, (ii) identify those two regions in reverse orientation, and (iii) then take the resulting topological space with its boundary conned off.

3 Average edge order of a normal 3-pseudomanifold

Consider K as a normal 3-pseudomanifold with singularities, containing n singular vertices denoted by $t_1, t_2, \dots, t_r, p_1, p_2, \dots, p_{r'}$, where $r + r' = n$ for some $r, r' \in \mathbb{N}$. According to surface classification, each vertex $x \in K$ has a link, $\mathrm{lk}\,(x)$, which is either a triangulated sphere, a triangulation of connected sums of tori, or a triangulation of connected sums of projective planes. Specifically, if $|\mathrm{lk}\,(t_i)| \cong \#h_i\mathbb{T}^2$ and $|\mathrm{lk}\,(p_j)| \cong \#m_j\mathbb{RP}^2$ for $h_i \geq 0$, $m_j \geq 0$, $1 \leq i \leq r$, and $1 \leq j \leq r'$, where $\mathbb{T}^2$ and $\mathbb{RP}^2$ represent a torus and a projective plane, respectively, then $\chi(t_i) = 2 - 2h_i$ and $\chi(p_j) = 2 - m_j$.

Let V, E, F, and T denote the count of vertices, edges, faces, and tetrahedra in K, respectively. Considering $g_2(K) = E - 4V + 10$ and $g_3(K) = F - 3E + 6V - 10$, it follows that $g_2(K) + g_3(K) = F - 2E + 2V$.

Additionally, we have

$$g_2(K) + g_3(K) = \sum_{v \in K} (2 - \chi(\mathrm{lk}\,(v))).$$

Thus, the average edge order of a normal 3-pseudomanifold is,

$$\begin{aligned}
\mu_0(K) &= \frac{3F}{E} = 3\frac{(g_2(K) + g_3(K) + 2E - 2V)}{E} \\
&= 3\frac{(\sum_{v \in K}(2 - \chi(\mathrm{lk}\,(v))) + 2E - 2V)}{E} \\
&= 3\frac{(2V - \sum_{v \in K}\chi(\mathrm{lk}\,(v)) + 2E - 2V)}{E} \\
&= 3\frac{(-\sum_{v \in K}\chi(\mathrm{lk}\,(v)) + 2E)}{E} \\
&= 6 - 3\frac{\sum_{v \in K}\chi(\mathrm{lk}\,(v))}{E} \\
&= 6 - 3\frac{2(V-n)}{E} - 3\frac{\sum_{i=1}^{r}\chi(\mathrm{lk}\,(t_i))}{E} - 3\frac{\sum_{i=1}^{r'}\chi(\mathrm{lk}\,(p_i))}{E} \\
&= 6 - 6\frac{(V-n)}{E} - 3\frac{\sum_{i=1}^{r}(2 - 2h_i)}{E} - 3\frac{\sum_{i=1}^{r'}(2 - m_i)}{E} \\
&= 6 - 6\frac{V}{E} + 6\frac{n}{E} - 6\frac{(r + r')}{E} + 6\frac{\sum_{i=1}^{r} h_i}{E} + 3\frac{\sum_{i=1}^{r'} m_i}{E} \\
&= 6 - 6\frac{V}{E} + 6\frac{\sum_{i=1}^{r} h_i}{E} + 3\frac{\sum_{i=1}^{r'} m_i}{E}.
\end{aligned}$$

Proof of Theorem 3 (*ii*):
Suppose that $\mu_0(K) \leq \frac{9}{2}$, then we obtain

$$\begin{aligned}
& 6 - 6\frac{V}{E} + 6\frac{\sum_{i=1}^{r} h_i}{E} + 3\frac{\sum_{i=1}^{r'} m_i}{E} \leq \frac{9}{2} \\
\implies & -6\frac{V}{E} + 6\frac{\sum_{i=1}^{r} h_i}{E} + 3\frac{\sum_{i=1}^{r'} m_i}{E} \leq \frac{9}{2} - 6 \\
\implies & 4\frac{V}{E} - 4\frac{\sum_{i=1}^{r} h_i}{E} - 2\frac{\sum_{i=1}^{r'} m_i}{E} \geq 1 \\
\implies & E - 4V + 10 \quad \leq 10 - 4\sum_{i=1}^{r} h_i - 2\sum_{i=1}^{r} m_i \\
\implies & g_2(K) \quad \leq 10 - 4\sum_{i=1}^{r} h_i - 2\sum_{i=1}^{r} m_i.
\end{aligned}$$

Consider h as the maximum value in the set $\{h_1, h_2, ..., h_r\}$, and let m be the maximum value in $\{m_1, m_2, ..., m_r'\}$. Given Lemma 5, which states that $g_2(K) \geq$

$g_2(\mathrm{lk}\,(t))$ for any $t \in K$, we derive:

$$6h \leq 10 - 4\sum_{i=1}^{r} h_i - 2\sum_{i=1}^{r} m_i$$

and

$$3m \leq 10 - 4\sum_{i=1}^{r} h_i - 2\sum_{i=1}^{r} m_i.$$

It is evident that the inequalities above are satisfied if and only if one of the following situations arises:
(a) $n = 1$, $h = 1$, $m = 0$
(b) $n = 1$, $h = 0$, $m = 1$
(c) $n = 1$, $h = 0$, $m = 2$
(d) $n = 2$, $h = 0$, $m = 1$
(e) $n = 3$, $h = 0$, $m = 1$.

Due to the fact that the sum of Euler characteristics for the vertex links of a normal 3-pseudomanifold is always even, we can infer that cases (b) and (e) are not feasible.

Cases $(a), (c)$**:** In both cases, $g_2(K) \leq 6$, and K features precisely one singularity at either t_1 or p_1, where $\mathrm{lk}\,(t_1)$ forms a triangulated torus and $\mathrm{lk}\,(p_1)$ constitutes a triangulated Klein bottle. Moreover, $g_2(\mathrm{lk}\,(t_1)) = g_2(\mathrm{lk}\,(p_1)) = 6$. Additionally, according to Proposition 5, we establish $g_2(K) \geq 6$. Consequently, $g_2(K) = 6$. Henceforth, according to Proposition 6, K is obtained from some boundary complexes of 4-simplices by a sequence of operations, including connected sums, bistellar 1-moves, edge contractions, edge expansions, and vertex folding. Furthermore, $|K|$ is a handlebody with its boundary coned off.
Case (d)**:** In this instance, K encompasses two $\mathbb{RP}^2$ singularities at vertices p_1 and p_2. According to Proposition 5, we have $3 \leq g_2(K) \leq 6$. Consequently, by Proposition 7, we deduce that K is obtained from some boundary complexes of 4-simplices by a sequence of possible operations, including connected sums, bistellar 1-moves, edge contractions, edge expansions, and an edge folding. Moreover, K is the triangulation of suspension of $\mathbb{RP}^2$ by Remark 8.

Hence, by combining cases $(a), (c)$ and (d), proof of Theorem 3 (ii) follows.

Proof of Theorem 3 (i):
Consider K as a normal 3-pseudomanifold with singularities, where $\mu_0(K) \leq \frac{30}{7}$. Therefore, $\mu_0(K) \leq \frac{30}{7} < \frac{9}{2}$. Consequently, based on the preceding discussions, we have only three potential cases for K: (a), (c), and (d). In both cases (a) and (c), $\mu_0(K) = \frac{9}{2} > \frac{30}{7}$, rendering them implausible. Now, let's discuss the case (d). Assume K contains two $\mathbb{RP}^2$ singularities at vertices u and v. Hence, $3 \leq g_2(K) \leq 6$. Then, we obtain

$$\begin{aligned}
\mu_0(K) = 6 - \frac{6(V-1)}{E} &\leq \frac{30}{7} \\
\implies 6 - \frac{6(V-1)}{4V - 10 + g_2(K)} &\leq \frac{30}{7} \\
\implies \frac{18V + 6(g_2(K-9))}{4V - 10 + g_2(K)} &\leq \frac{30}{7} \\
\implies 126V - 378 + 42g_2(K) &\leq 120V + 30g_2(K) - 300 \\
\implies V \quad &\leq 13 - 2g_2(K).
\end{aligned}$$

Considering that for any normal 3-pseudomanifold Δ, $V(\Delta) \geq 7$, the aforementioned inequality holds if and only if $g_2(K) = 3$ and $V(K) = 7$. Consequently, this implies that $\mu_0(K) = \frac{30}{7}$, and K represents a triangulation of the one-vertex suspension of $\mathbb{RP}^2$. Conversely, according to Remark 4, we obtain a triangulation K of the one-vertex suspension of $\mathbb{RP}^2$ with seven vertices, satisfying $\mu_0(K) = \frac{30}{7}$ and $g_2(K) = 3$. This concludes the proof of Theorem 3 (i).

Proof of Theorem 3 (iii):
Let h and m denote the same as previously stated. Therefore, $g_2(K) \geq 6h$, and $g_2(K) \geq 3m$. Consequently, $E(K) \geq 6h$ and $E(K) \geq 3m$.
Additionally, we have

$$\mu_0(K) = 6 - 6\frac{V}{E} + 6\frac{\sum_{i=1}^{r} h_i}{E} + 3\frac{\sum_{i=1}^{r'} m_i}{E}.$$

Thus,

$$\begin{aligned}
\mu_0(K) &= 6 - 6\frac{V}{E} + 6\frac{\sum_{i=1}^{r} h_i}{E} + 3\frac{\sum_{i=1}^{r'} m_i}{E} \\
&\leq 6 + 6\frac{rh}{E} + 3\frac{r'm}{E} \\
&< 6 + 6\frac{rh}{6h} + 3\frac{r'm}{3m} \\
&= 6 + r + r' \\
&= 6 + n.
\end{aligned}$$

Hence, $\mu_0(K) < 6 + n$.

Consider V_m, E_m, and F_m as the number of vertices, edges, and faces, respectively, of the minimal triangulation $\mathbb{T}_m$ representing a torus with genus $m \geq 3$. According to [6], we find: $V_m = \lceil \frac{7+\sqrt{1+48m}}{2} \rceil$, $E_m = 3V_m + 6m - 6$, $F_m = 2V_m + 4m - 4$. Now, if we take the suspension of the surface $\mathbb{T}_m$, we obtain a normal 3-pseudomanifold denoted as K_m with two singularities, where: $V(K_m) = V_m + 2$, $E = 2V_m + E_m$, and $F = 2E_m + F_m$. Then,

$$\mu_0(K_m) = \frac{3F}{E} = \frac{6E_m + 3F_m}{2V_m + E_m} = \frac{24V_m + 48m - 48}{5V_m + 6m - 6}.$$

Thus, $\mu_0(K_m) \to 8$ as $m \to \infty$. This completes the proof of Theorem 3(iii).

Remark 9. If K_1 represents a normal 3-pseudomanifold obtained from a stacked sphere K' with $g_2(K') = 0$ via a vertex folding, then the following relations hold: $E(K') = 4V(K') - 10$, $F(K') = 6V(K') - 20$, $T(K') = 3V(K') - 10$. Moreover, $V(K_1) = V(K') - 3$, $E(K_1) = E(K') - 6 = 4V(K') - 16$, $F(K_1) = F(K') - 4 = 6V(K') - 24$, $T(K_1) = T(K') - 2 = 3V(K') - 12$. Now, we find that

$$\mu_0(K_1) = \frac{3F}{E} = 3\frac{6V-24}{4V-16} = \frac{9}{2}(\frac{V-4}{V-4}) = \frac{9}{2}.$$

Corollary 10. *If K is a normal 3-pseudomanifold and $\mu_0(K) < \frac{30}{7}$, then K is triangulation of a 3-sphere.*

Proof. If K represents a normal 3-pseudomanifold and $\mu_0(K) < \frac{30}{7}$, according to Theorem 3, K does not contain any singular vertices. This indicates that K forms a triangulation of a 3-manifold, where $\mu_0(K) < \frac{30}{7}$. Consequently, the result follows from Proposition 1(iii). $\square$

4 On the upper bound of average edge order

The upper bound provided in Theorem 3 varies according to the number of singularities. Consequently, the task of determining an upper bound for the average edge orders of all normal 3-pseudomanifolds remains unresolved. In this context, we present several examples of normal 3-pseudomanifolds with singularities that possess a higher average edge order, sourced from the simplicial complex library of GAP [5]. Denote by [SCLib, n] the simplicial complex located at the n-th position within the simplicial complex library in GAP.

Examples:

1. Let $K_1 = [\text{SCLib}, 60]$ and $K_2 = [\text{SCLib}, 61]$, which represent two normal 3-pseudomanifolds with the vector $(V, E, F, T) = (11, 55, 154, 77)$. Notably, both K_1 and K_2 exhibit $\mu_0(K_1) = \mu_0(K_2) = 8.4$. In both complexes, the link of each vertex forms a triangulation consisting of 6 copies of $\mathbb{RP}^2$. The automorphism groups for K_1 and K_2 are D_{22} (a dihedral group with 22 elements) and C_{11} (a cyclic group with 11 elements), respectively.

2. Let $K_3 = [\text{SCLib}, 540]$ be a normal 3-pseudomanifold with the vector (V, E, F, T) $= (17, 136, 544, 272)$. Consequently, $\mu_0(K_3) = 12$. The link of each vertex in K_3 forms a triangulation consisting of 18 copies of $\mathbb{RP}^2$. The automorphism group of K_3 is $C_{17} : C_{16}$, a semidirect product of cyclic groups of orders 16 and 17.

3. Let $K_5 = [\text{SCLib}, 587]$ be a normal 3-pseudomanifold with the vector (V, E, F, T) $= (19, 171, 684, 342)$. Consequently, $\mu_0(K_5) = 12$. The link of each vertex in K_5 forms a triangulation consisting of 20 copies of $\mathbb{RP}^2$. The automorphism group of K_5 is $C_{19} : C_{18}$.

4. Let $K_4 = [\text{SCLib}, 541]$ be a normal 3-pseudomanifold with the vector (V, E, F, T) $= (17, 136, 680, 340)$. Consequently, $\mu_0(K_4) = 15$. The link of each vertex in K_4

forms a triangulation consisting of 26 copies of $\mathbb{RP}^2$. The automorphism group of K_4 is the same as the automorphism group of K_3, i.e., $C_{17} : C_{16}$.

It is interesting to note that in all the above examples, each vertex in the complexes is a singular vertex. This draws attention towards the study of normal 3-pseudomanifolds with all vertices being singular. Additionally, it's worth mentioning that all of these complexes exhibit neighborly properties, with even the example of K_4 being 3-neighborly. If a normal 3-pseudomanifold K is 3-neighborly, then $\mu_0(K) = \frac{3\binom{V}{3}}{\binom{V}{2}} = V - 2$. Based on all the observations we obtained, we arrive at the following questions and conjectures:

Question 11. *How many vertices are needed at minimum to construct a neighborly normal* 3*-pseudomanifold, where the link of each vertex forms a triangulated surface of genus* g *for* $g \in \mathbb{N}$*?*

Question 12. *Is there a* 3*-neighborly normal* 3*-pseudomanifold* K *with* $V(K) > n$ *for every positive integer* $n \in \mathbb{N}$*?*

If an affirmative response exists for the preceding question, then the following conjecture is easily valid.

Conjecture 13. *For every positive integer* $n \in \mathbb{N}$*, there exists a normal* 3*-pseudomanifold* K *such that* $\mu_0(K) > n$*.*

Acknowledgement: The first author is supported by the Mathematical Research Impact Centric Support (MATRICS) Research Grant (MTR/2022/000036). The second author is supported by CSIR (India).

Data availability: The authors declare that all data supporting the findings of this study are available within the article.

Declarations

Conflict of interest: No potential conflict of interest was reported by the authors.

References

[1] B. Bagchi and B. Datta, Lower bound theorem for normal pseudomanifolds, *Expo. Math.* 26 (2008), 327–351.

[2] B. Basak, R.K. Gupta, S. Sarkar, A characterization of normal 3-pseudomanifolds with at most two singularities, *Discrete Math.* 346 (2023), no. 12, Paper No. 113588, 15 pp.

[3] B. Basak and E. Swartz, Three-dimensional normal pseudomanifolds with relatively few edges, *Advances in Mathematics* 365 (2020) 107035, 1-25.

[4] M. R. Casali, The average edge order of 3-manifold coloured triangulations, *Canad. Math. Bull.* 37(2) (1994) 154-161.

[5] F. Effenberger and J. Spreer, simpcomp-a GAP toolkit for simplicial complexes, Version 4.12.0 (2022) http://www.igt.uni-stuttgart.de/LstDiffgeo/simpcomp.

[6] M. Jungerman and G. Ringel, Minimal triangulations on orientable surfaces. *Acta Math.* 145(1980), 121-154.

[7] G. Kalai, Rigidity and the lower bound theorem I, *Invent. Math.* 88 (1987), 125–151.

[8] F. Luo and R. Stong, Combinatorics of triangulations of 3-manifolds, *Trans. Amer. Math. Soc.* 337 (1993), 891-906.

[9] E. Swartz, Thirty-five years and counting, arXiv:1411.0987.

[10] M. Tamura, The average edge order of triangulations of 3-manifolds, *Osaka J. Math.* 33 (1996), 761-773.

[11] M. Tamura, The average edge order of triangulations of 3-manifolds with boundary, *Trans. Amer. Math. Soc.* 350 (1998), 2129-2140.