Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 6 February 2018 (MN LaTeX style file v2.2)

A model of rotating hotspots for 3:2 frequency ratio of HFQPOs in black hole X-ray binaries

Ding-Xiong Wang[1,3], Yong-Chun Ye[1], Guo-Zheng Yao[1,2] and Ren-Yi Ma[1]
1 *Department of Physics, Huazhong University of Science and Technology, Wuhan,430074,China*
2 *Department of Physics, Beijing Normal University, Beijing 100875, China*
3 *Send offprint requests to: D.-X. Wang (dxwang@hust.edu.cn)*

6 February 2018

ABSTRACT

We propose a model to explain a puzzling 3:2 frequency ratio of high frequency quasi-periodic oscillations (HFQPOs) in black hole (BH) X-ray binaries, GRO J1655-40, GRS 1915+105 and XTE J1550-564. In our model a non-axisymmetric magnetic coupling (MC) of a rotating black hole (BH) with its surrounding accretion disc coexists with the Blandford-Znajek (BZ) process. The upper frequency is fitted by a rotating hotspot near the inner edge of the disc, which is produced by the energy transferred from the BH to the disc, and the lower frequency is fitted by another rotating hotspot somewhere away from the inner edge of the disc, which arises from the screw instability of the magnetic field on the disc. It turns out that the 3:2 frequency ratio of HFQPOs in these X-ray binaries could be well fitted to the observational data with a much narrower range of the BH spin. In addition, the spectral properties of HFQPOs are discussed. The correlation of HFQPOs with jets from microquasars is contained naturally in our model.

Key words:
accretion, accretion discs — black hole physics — magnetic field — instabilities — stars: individual (GRO J1655-40) — stars: individual (GRS 1915+105) — stars: individual (XTE J1550-564) — stars: oscillations — X-rays: stars

1 INTRODUCTION

Quasi-periodic oscillations in X-ray binaries have become a very active research field since the launch of the *Rossi X-Ray Timing Explorer* (*RXTE*; Bradt, Rothschild & Swank 1993). A key feature in these sources is that some of high frequency quasi-periodic oscillations (HFQPOs) appear in pairs. Five black hole (BH) X-ray binaries exhibit transient HFQPOs, of which three sources have pairs occurring in GRO J1655-40 (450, 300∆ruby∆∆rb∆Hz∆/rb∆∆rt∆

2002b; Wang et al. 2003a, hereafter W03a). In addition, we explained the HFQPOs in X-ray binaries based on a model of a rotating hotspot due to the MC of a rotating BH with the inner region of the disc (Wang et al. 2003b, hereafter W03b). However, the 3:2 ratio of HFQPOs has not been discussed by virtue of the MC process.

Very recently, we discussed the condition for the coexistence of the BZ and MC processes, and found that this coexistence always accompanies the screw instability of the magnetic field in BH magnetosphere, provided that the BH spin and the power-law index for the variation of the magnetic field on the disc are greater than some critical values (Wang et al. 2003c, 2004, hereafter W03c and W04, respectively).

In this paper we propose a model to explain the 3:2 ratio of HFQPOs in BH X-ray binaries by virtue of non-axisymmetric MC of a rotating BH with its surrounding accretion disc. The upper frequency arises from a rotating hotspot near the inner edge of the disc, and the lower frequency is produced by the screw instability somewhere away from the inner edge. It turns out that the 3:2 ratio of HFQPOs in GRO J1655-40, GRS 1915+105 and XTE J1550-564 is well fitted by energy transferred in the MC process, where six parameters are used to describe the BH mass, spin and distribution of a non-axisymmetric magnetic field. The 3:2 ratio obtained in our model provides a much narrower range of the BH spin compared with the other models. Furthermore, the jets from these BH binaries are explained naturally by the BZ process coexisting with the MC process.

This paper is organized as follows. In section 2 we give a brief description of our model. In section 3 we fit the upper and lower frequencies of HFQPOs by the inner and outer hotspots co-rotating with the disc. The discussion is focused at the issues of the outer hotspot. A scenario of how the presence of the screw instability leads to the outer hotspot is given. In addition, the time-scale of the screw instability is estimated by using an equivalent R-L circuit, and the spectral properties of HFQPOs are discussed by assuming the existence of corona above the disc. In section 4 we discuss some issues of astrophysical meanings related to the 3:2 ratio. Throughout this paper the geometric units $G = c = 1$ are used.

2 DESCRIPTION OF OUR MODEL

We intend to fit the 3:2 ratio of HFQPOs by virtue of an rotating BH surrounded by a magnetized accretion disc based on the following assumptions.

(1) The configuration of the poloidal magnetic field is shown in Fig. 1. The radius r_{ms} indicates the inner edge of the disc, being the innermost stable circular orbit (ISCO) (Novikov & Thorne 1973, hereafter NT73), and r_S is the radius where the screw instability of the magnetic field might occur, which is related to the angle θ_S by the same mapping relation given in W04. As shown in Fig. 1, θ_S is the angular boundary between the open and closed field lines on the horizon, and θ_L is the lower boundary angle for the closed field lines connecting the BH with the disc. Throughout this paper $\theta_L = 0.45\pi$ is assumed in calculations.



Figure 1. Poloidal magnetic field connecting a rotating BH with a remote astrophysical load and the surrounding disc.

(2) The poloidal magnetic field on the BH horizon is non-axisymmetric as assumed in W03b, varying with the azimuthal angle ϕ as follows (see Fig.2 in W03b),

$$B_H^p(\phi) = \sqrt{\left\langle (B_H^p)^2 \right\rangle} f(\phi), \quad f(\phi) \equiv \left\{ \begin{array}{l} 1+\delta,\ 0<\phi<\Delta\phi, \\ 1,\ \Delta\phi \leqslant \phi \leqslant 2\pi, \end{array} \right. \tag{1}$$

where $\sqrt{\left\langle (B_H^p)^2 \right\rangle}$ is root-mean-square of the poloidal magnetic field over the angular coordinate from $\theta = 0$ to θ_L. The parameter δ is used to describe the strength of the bulging magnetic field in the azimuthal region $0 < \phi < \Delta\phi$.

(3) The magnetosphere is assumed to be force-free outside the BH and the disc, and the closed magnetic field lines are frozen in the disc. The disc is thin and perfectly conducting, lies in the equatorial plane of the BH with the inner boundary being at ISCO.

(4) The poloidal magnetic field varies as a power law with the radial coordinate of the disc as follows,

$$B_D^p \propto \xi^{-n}, \tag{2}$$

where B_D^p represents the poloidal magnetic field on the disc, and the parameter n is the power-law index, and the dimensionless radial coordinate is defined as $\xi \equiv r/r_{ms}$. The radius r_{ms} is related to the BH mass M and the dimensionless spin parameter $a_* \equiv a/M$ as given in NT73.

(5) The magnetic flux connecting the BH with its surrounding disc takes precedence over that connecting the BH with the remote load. As argued in W03c and W04, assumption 5 is crucial for the coexistence of the BZ and MC processes.

The magnetosphere anchored in a Kerr BH and its surrounding disc is described in Boyer-Lindquist coordinates, in which the following parameters are involved (MacDonald and Thorne 1982, hereafter MT82).

$$\left\{ \begin{array}{l} \Sigma^2 = \left(r^2+a^2\right)^2 - a^2\Delta\sin^2\theta,\ \rho^2 = r^2 + a^2\cos^2\theta, \\ \Delta = r^2 + a^2 - 2Mr,\ \varpi = (\Sigma/\rho)\sin\theta, \\ \alpha = \rho\sqrt{\Delta}\big/\Sigma. \end{array} \right. \tag{3}$$

In W03b the BZ and MC powers in non-axisymmetic case are related to those in axisymmetic case by

$$P_{BZ}^{NA} = \lambda P_{BZ}^{A}, \tag{4}$$

$$P_{MC}^{NA} = \lambda P_{MC}^{A}, \tag{5}$$

where P_{BZ}^{NA} and P_{MC}^{NA} are the powers in non-axisymmetic case, and P_{BZ}^{A} and P_{MC}^{A} are the powers in axisymmetic case, respectively. The parameter λ is expressed in terms of δ and $\varepsilon \equiv \Delta\phi/2\pi$ by

$$\lambda = \left[(1+\delta)\,\varepsilon + (1-\varepsilon)\right]^2 = (1+\delta\varepsilon)^2 . \tag{6}$$

Similarly, the BZ and MC torques in non-axisymmetic case are related to those in axisymmetic case by

$$T_{BZ}^{NA} = \lambda T_{BZ}^{A}, \tag{7}$$

$$T_{MC}^{NA} = \lambda T_{MC}^{A}, \tag{8}$$

where T_{BZ}^{NA} and T_{MC}^{NA} are the torques in non-axisymmetic case, and T_{BZ}^{A} and T_{MC}^{A} are the torques in axisymmetic case, respectively. The powers P_{BZ}^{A} and P_{MC}^{A}, and the torques T_{BZ}^{A} and T_{MC}^{A} are given in W02 as follows.

$$P_{BZ}^{A}/P_0 = 2a_*^2 \int_0^{\theta} \frac{k\,(1-k)\sin^3\theta d\theta}{2-(1-q)\sin^2\theta}, \quad 0 < \theta < \theta_S, \tag{9}$$

$$P_{MC}^{A}/P_0 = 2a_*^2 \int_{\theta_S}^{\theta} \frac{\beta\,(1-\beta)\sin^3\theta d\theta}{2-(1-q)\sin^2\theta}, \quad \theta_S < \theta < \theta_L. \tag{10}$$

$$\begin{aligned} &T_{BZ}^{A}/T_0 = \\ &\quad 4a_*\,(1+q) \int_0^{\theta} \tfrac{(1-\beta)\sin^3\theta d\theta}{2-(1-q)\sin^2\theta}, \quad 0 < \theta < \theta_S, \end{aligned} \tag{11}$$

$$\begin{aligned} &T_{MC}^{A}/T_0 = \\ &\quad 4a_*\,(1+q) \int_{\theta_S}^{\theta} \tfrac{(1-\beta)\sin^3\theta d\theta}{2-(1-q)\sin^2\theta}, \quad \theta_S < \theta < \theta_L. \end{aligned} \tag{12}$$

In equations (9)—(12), k and β are the ratios of the angular velocities of the magnetic field lines to the angular velocity of the BH horizon in the BZ and MC processes, respectively. The quantity $q \equiv \sqrt{1-a_*^2}$ is a function of the BH spin, and P_0 and T_0 are defined as

$$\left\{ \begin{array}{l} P_0 \equiv \left\langle (B_H^p)^2 \right\rangle M^2 \approx B_4^2 m_{BH}^2 \times 6.59 \times 10^{28} erg \cdot s^{-1}, \\ T_0 \equiv \left\langle (B_H^p)^2 \right\rangle M^3 \approx B_4^2 m_{BH}^3 \times 3.26 \times 10^{23} g \cdot cm^2 \cdot s^{-2}, \end{array} \right. \tag{13}$$

where B_4 is the strength of the poloidal magnetic field on the horizon in units of $10^4 gauss$, and m_{BH} is defined as $m_{BH} \equiv M/M_{\odot}$.

So far six parameters are involved in our model, in which m_{BH} and a_* are used for describing the mass and spin of the Kerr BH, and the three parameters, B_4, δ and ε are used for describing the non-axisymmetric magnetic field on the horizon, the power-law index n is for the variation of the magnetic field with the radial coordinate of the disc.

3 INNER AND OUTER HOTSPOTS FOR 3:2 FREQUENCY RATIO

As argued in W03b, the upper frequency of HFQPOs is modulated by an inner hotspot, which corresponds to the maximum of function F_{QPO} as follows,

$$F_{QPO} \equiv r^2 F / r_{ms}^2 F_0 = \xi^2 (F_{DA} + F_{MC})/F_0, \tag{14}$$

where F_{DA} is the radiation flux due to disc accretion, and F_{MC} is radiation flux arising from the energy transferred from the rotating BH into the inner disc in the MC process. We find that the radial coordinate of the inner hotspot, $\xi_{\max}$, only depends on the parameters a_*, and n, i.e., it is independent of the parameters m_{BH}, B_4, δ and ε. Considering that the hotspot is frozen at the disc, we have the upper frequency ν_{QPO}^{upper} by substituting $\xi_{\max}$ into the Keplerian angular velocity as follows,

$$\nu_{QPO} = \nu_0 (\xi^{3/2} \chi_{ms}^3 + a_*)^{-1}, \tag{15}$$

where $\nu_0 \equiv (m_{BH})^{-1} \times 3.23 \times 10^4 Hz$. It turns out that ν_{QPO}^{upper} depends on the three parameters, m_{BH}, a_* and n.

The lower frequency of HFQPOs is fitted by an outer hotspot rotating with the Keplerian angular velocity expressed by equation (15), and a scenario for the production of the outer hotspot is given as follows.

It is well known that the magnetic field configurations with both poloidal and toroidal components can be screw instable (Kadomtsev 1966; Bateman 1978). According to the Kruskal-Shafranov criterion (Kadomtsev 1966), the screw instability will occur, if the toroidal magnetic field becomes so strong that the magnetic field line turns around itself about once. Recently, some authors discussed the screw instability in BH magnetosphere (Gruzinov 1999a; Li 2000b; Tomimatsu et el. 2001). In W04 we suggested that the criterion for the screw instability in the MC process could be expressed by

$$F_{Screw}\,(a_*; \xi, n) - L/(2\pi\varpi_D) \leqslant 0, \tag{16}$$

where L is the poloidal length of the closed field line connecting the BH and the disc, ϖ_D is the cylindrical radius on the disc, and $F_{Screw}\,(a_*; \xi, n)$ is a function of a_*, ξ and n.

Setting equality in equation (16), we obtain the critical radial coordinate $\xi_S \equiv r_S/r_{ms}$ for the screw instability, which depends on the parameters a_* and n.

It is believed that a disc is probably surrounded by a high-temperature corona analogous to the solar corona (Liang & Price 1977; Haardt 1991; Zhang et al 2000). Very recently, some authors argued that the coronal heating in some stars including the Sun is probably related to dissipation of currents, and very strong X-ray emissions arise from variation of magnetic fields (Galsgaard & Parnell 2004; Peter et al. 2004).

Analogously, if the corona exists above the disc in our model, we expect that the corona above ξ_S might be heated by the induced current due to the screw instability of the non-axisymmetric magnetic field. Therefore a very strong X-ray emission would be produced above the rotating outer hotspot, and the lower frequency ν_{QPO}^{lower} is obtained directly by substituting ξ_S into equation (15).

The time-scale of the screw instability can be estimated by using an equivalent circuit as shown in Fig. 2, in which the segments $\boldsymbol{LM}$ and $\boldsymbol{KN}$ represent the two adjacent magnetic surfaces connecting the BH horizon and the outer hotspot as shown in Fig. 2a. Considering the existence of the toroidal magnetic field threading each loop, we introduce an inductor into the circuit, which is represented by the symbol ΔL in Fig. 2b.

$$\Delta\varepsilon_H = (\Delta\Psi^p/2\pi)\,\Omega_H, \quad \Delta\varepsilon_D = -(\Delta\Psi^p/2\pi)\,\Omega_D, \tag{17}$$

where $\Delta\Psi^p$ is the flux of the poloidal magnetic field sandwiched by the magnetic surfaces $\boldsymbol{LM}$ and $\boldsymbol{KN}$. The resistance at the BH horizon is given by

$$\Delta R_H = \sigma_H \frac{\Delta l}{2\pi\varpi_H} = \frac{2\rho_H \Delta\theta_H}{\varpi_H}, \tag{18}$$

where $\sigma_H = 4\pi = 377 ohm$ is the surface resistivity of the BH horizon (MT82), and $\Delta\theta_H$ is the angle on the horizon spanned by the two surfaces. The quantities ρ_H and ϖ_H are the Kerr metric parameters at the horizon. The inductance ΔL in the R-L circuit is defined by

$$\Delta L = \Delta\Psi^T \big/ I^p, \tag{19}$$

where I^p and $\Delta\Psi^T$ are the poloidal current flowing in the circuit and the flux of the toroidal magnetic field threading the circuit, respectively. The flux $\Delta\Psi^T$ can be integrated over the region $KLMN$ as follows,

$$\Delta\Psi^T = \oint_{loop} B^T \sqrt{g_{rr} g_{\theta\theta}}\, dr d\theta, \tag{20}$$

where the toroidal magnetic field measured by "zero-angular-momentum observers" (ZAMOs) is given as

$$B^T = 2I^p/(\alpha\varpi), \tag{21}$$

where α is the lapse function defined in equation (3).

Unfortunately, the geometric shapes of the magnetic surfaces $\boldsymbol{LM}$ and $\boldsymbol{KN}$ are unknown. As an approximate estimation we assume that the surfaces $\boldsymbol{LM}$ and $\boldsymbol{KN}$ are formed by rotating the two adjacent circles around the symmetric axis, and $\Delta\Psi^T$ can be calculated by integrating over the region $\boldsymbol{KLMN}$ as shown in Fig. 3. Finally, we calculate the quantity $\tau \equiv \Delta L/\Delta R_H$ by incorporating equations (18)—(21), which is independent of I^P.

As a simple analysis we can divide one event of the screw instability into two stages. In stage 1 the instability starts, and the magnetic energy is released rapidly, the poloidal current and thus the toroidal magnetic field decrease to zero in very short time. In stage 2 the toroidal magnetic field recovers gradually with the increasing poloidal current I^P in the R-L circuit, until the next event of the instability occurs according to the criterion (16). It seems reasonable that the duration in stage 1 is much less than that in stage 2, and the time-scale of the screw instability depends mainly on the duration of stage 2, which is regarded as the relax time of the poloidal current I^P in the R-L circuit. The current I^P is governed by the following equation,

$$\Delta L \frac{dI^P}{dt} + \Delta R_H I^P = \Delta\varepsilon_H + \Delta\varepsilon_L. \tag{22}$$

Setting the initial condition, $I^P = 0$, we have the solution,

$$I^P(t) = I^p_{steady}\left(1 - e^{-t/\tau}\right), \tag{23}$$

where I^p_{steady} is the steady current in the R-L circuit, and it reads

$$I^p_{steady} = (\Delta\varepsilon_H + \Delta\varepsilon_L)/\Delta R_H. \tag{24}$$

From equation (23) we obtain that the poloidal current attains 99.3% of I^p_{steady} in the relax time $t_{relax} = 5\tau$, implying the recovery of the toroidal magnetic field. Thus the time-scale of the screw instability can be estimated as

$$t_{Screw} > t_{relax} = 5\tau. \tag{25}$$

Summarizing the above results, we have the upper and lower frequencies of HFQPOs, and the corresponding values of $\xi_{\max}$, ξ_S and t_{relax} as shown in Table 1. It is easy to check from Table 1 that the time-scale of the screw instability is generally greater than the corresponding periods of HFQPOs, i.e.,

$$t_{Screw} > t_{relax} > 1\big/\nu^{upper}_{QPO} > 1\big/\nu^{lower}_{QPO}. \tag{26}$$



Figure 2. (a) Two adjacent magnetic surfaces connecting the BH horizon and the outer hotspot, (b) one loop of equivalent circuit for screw instability.

As pointed out in MR03, the spectral properties of accreting black holes in states which show HFQPOs are usually dominated by a steep power-law (SPL) component. However, the origin of X-ray power-law remains controversial. Most of models for the SPL state invoke inverse Compton scattering as the operant radiation mechanism, and the radiation spectrum formed through Comptonization of low-frequency photons in a hot thermal plasma cloud may be described by a power law (Pozdnyakov, Sobol & Syunyaev 1983). It is noted that the origin of the Comptonizing electrons involves magnetic instabilities in the accretion disc (Poutanen & Fabian 1999), with which the screw instability in our model is consistent.

Suppose that the spectral properties of the inner and outer hotspots arise from blackbody spectra, the effective radiation temperature $(T_{HS})_{eff}$ is expressed by

$$T_{eff} = T_0\left[f_{DA} + f_{MC}\right]^{1/4}, \tag{27}$$

$$T_0 = (F_0/\sigma)^{1/4} \approx 4.8 \times 10^5 B_4^{1/2}\,(K), \tag{28}$$

Table 1. The 3:2 ratio of HFQPOs produced by the inner and outer hotspots with $\delta = 0.5$ and $\varepsilon = 0.2$. Henceforth BHB-I, II and III represent GRO J1655-40, GRS 1915+105 and XTE J1550-564, respectively.

BHB	a_*	n	m_{BH}	t_{relax} (sec)	Inner Hotspot			Outer Hotspot		
					$\xi_{\max}$	$E_{HS}^{upper} \Big/ B_4^{1/2}$ *(kev)*	ν_{QPO}^{upper} *(Hz)*	ξ_{SC}	$E_{HS}^{lower} \Big/ B_4^{1/2}$ *(kev)*	ν_{QPO}^{lower} *(Hz)*
I	0.777	6.22	6.8	7.61×10^{-3}	1.277	7.23×10^{-3}	450	2.015	4.96×10^{-3}	300
	0.730	6.27	5.8	6.57×10^{-3}	1.281	6.72×10^{-3}	450	2.097	4.46×10^{-3}	300
II	0.773	6.23	18	2.02×10^{-2}	1.277	7.19×10^{-3}	168	2.021	4.92×10^{-3}	113
	0.603	6.17	10	1.11×10^{-2}	1.297	5.55×10^{-3}	168	2.428	3.27×10^{-3}	113
III	0.788	6.19	11.5	1.28×10^{-2}	1.276	7.37×10^{-3}	276	2.000	5.09×10^{-3}	184
	0.696	6.29	8.5	9.72×10^{-3}	1.284	6.38×10^{-3}	276	2.159	4.14×10^{-3}	184

Note: The value ranges of the BH mass corresponding to GRO J1655-40, GRS 1915+105 and XTE J1550-564 are adopted from Greene et al. (2001), MR03 and Orosz et al. (2002), respectively.



Figure 3. Two adjacent magnetic surfaces produced by two rotating circles, where the arrows represent the current flowing in the R-L circuit.

where σ is the Stefan-Boltzmann constant. The energy of the inner and outer hotspots can be estimated by

$$E_{HS} \equiv k_B T_{eff} = E_0 \left[f_{DA} + f_{MC} \right]^{1/4}, \tag{29}$$

$$E_0 = k_B \left(F_0 / \sigma \right)^{1/4} \approx 4.14 \times 10^{-2} B_4^{1/2} \left(kev \right), \tag{30}$$

where k_B is the Boltzmann constant. Substituting $\xi_{\max}$ and ξ_S into equation (29), we have the energy of the inner and outer hotspots E_{HS}^{upper} and E_{HS}^{lower} as shown in Table 1.

Inspecting equation (29) and the expressions for f_{DA} and f_{MC} given in W03b, we find that both E_{HS}^{upper} and E_{HS}^{lower} depend not only on the parameters, a_* and n, but also on the parameters for the non-axisymmetric magnetic field, $B_4^{1/2}$, δ and ε. It is worthy to point out that both E_{HS}^{upper} and E_{HS}^{lower} are independent of the BH mass of the X-ray binaries.

4 DISCUSSION

Inspecting Table 1, we find that the 3:2 ratios of HFQPOs in GRO J1655-40, GRS 1915+105 and XTE J1550-564 are well fitted by adjusting the two parameters, a_* and n, for the given value ranges of m_{BH}. In this section we discuss some issues of astrophysical meanings related to the 3:2 ratio of HFQPOs in the BH binaries.

4.1 3:2 ratio and magnetic field on the BH horizon

It is pointed out that the twin HFQPOs are not always detected together in GRO J1655-40, i.e., 300 Hz QPO is observed sometimes without 450 Hz detection, while 450 Hz QPO is observed sometimes without 300 Hz detection (S01a). In our model the upper and lower frequencies of HFQPOs are produced by the inner and outer hotspots, which are located at different sites of the disc. In addition, our model is proposed based on the non-axisymmetric magnetic field on the horizon, and two different physical mechanisms are involved, i.e., the MC process for the inner hotspot and the screw instability for the outer hotspot. Since the non-axisymmetric magnetic field cannot be stationary on the horizon, we can understand that the twin HFQPOs do not have to occur simultaneously in some cases.

In Table 1, both E_{HS}^{upper} and E_{HS}^{lower} reach the energy level of emitting X-ray, provided that the root-mean square of the magnetic field on the horizon is strong enough to be $\sim 10^9 gauss$ ($B_4 \sim 10^5$). Considering that a system cannot radiate a given flux at less than the blackbody temperature (Frank et al. 1992), the strength of magnetic field of $\sim 10^9 gauss$ might be regarded as a upper limit for energy level of emitting X-ray in BH binaries.

It is pointed out in S01a that the energy spectra of the 300 Hz QPO appear to be significantly different from those of the 450 Hz QPO in GRO J1655-40: The former has a typical amplitude in the 2–12 keV band, while the latter is detected in the hard band, 13–27 keV. It is also shown in Fig. 4.16 of MR03 that the power densities corresponding to the upper frequency are greater than those corresponding to the lower frequency of HFQPOs in some BH X-ray binaries, such as GRO J1655-40 (450, 300ΔrubyΔΔrbΔHzΔ/rbΔΔrtΔ

(276, 184∆ruby∆∆rb∆Hz∆/rb∆∆rt∆

Table 2. The jet powers accompanying 3:2 ratio of HFQPOs with $\delta = 0.5$ and $\varepsilon = 0.2$.

BHB	a_*	n	m_{BH}	θ_S	$P_{BZ}^{NA}/B_4^2\,(erg/s)$
I	0.777	6.22	6.8	50.7^0	7.09×10^{28}
	0.730	6.27	5.8	48.9^0	3.98×10^{28}
II	0.773	6.23	18	50.5^0	4.87×10^{29}
	0.603	6.17	10	42.4^0	4.68×10^{28}
III	0.788	6.19	11.5	50.9^0	2.14×10^{29}
	0.696	6.29	8.5	47.7^0	7.01×10^{28}

(i) The minus sign in equation (33) shows that the current flows in the opposite direction to the disc matter; (ii) the power-law index $n+1$ in equation (33) implies that the profile of the current density in the central disc is steeper than that of the magnetic field. We give a primary explanation for these features as follows.

Probably the direction of the current arises from the effect of the closed magnetic field lines on the charged particles of the disc matter. In our model the angular velocity of the BH is required to be greater than that of the disc to produce the inner and outer hotspots on the disc. Therefore the charged particles of the disc matter will be dragged forward by the closed field lines. Considering that the mass of the positive charged particle, such as proton, is much greater than that of electron, we think that the bulk flow of electrons might be a little ahead of that of protons. Probably the minute difference in the two kinds of the bulk flows results in a current in the opposite direction to the disc.

On the other hand, the flux of angular momentum transferred from the rotating BH to the disc is given by

$$H_{MC}^{NA} = \frac{\partial T_{MC}^{NA}/\partial r}{4\pi r}. \tag{34}$$

Substituting equations (8) and (12) into equation (34), we have

$$\begin{aligned} H_{MC}^{NA} &= \tfrac{1}{4\pi\xi r_{ms}^2}\left(\partial\theta/\partial\xi\right)\left(\partial T_{MC}^{NA}/\partial\theta\right) \\ &= -\tfrac{\lambda a_*(1+q)}{\pi r_{ms}^2}\,\tfrac{(1-\beta)\xi^{-1}\mathrm{G}(a_*;\xi,n)}{\left[2\csc^2\theta-(1-q)\right]}. \end{aligned} \tag{35}$$

Incorporating equation (2) with equation (35), we have

$$H_{MC}^{NA} \propto \xi^{-n}. \tag{36}$$

Considering that the flux of angular momentum, H_{MC}^{NA}, transferred to the disc will block the accreting particles, and the blocking effect would be much more significant for electrons than protons due to the mass differences. Thus more electrons would stay in the region very close to ISCO. That is why the current density consist of the bulk flow of electrons and it varies with ξ in a power law expressed by equation (33).

As a summary we intend to give a comment for the six parameters involved in our model. The BH mass m_{BH} has its range given by observations, and $B_4 \approx 10^5$ is required by the hotspots for emitting X-ray. The parameters ε and δ are used to describe the non-axisymmetric magnetic field on the BH horizon. In fact, neither the position nor the energy of the inner and outer hotspots is sensitive to the values of ε and δ, and we take $\delta = 0.5$ and $\varepsilon = 0.2$ in calculations. Thus the rest two parameters, a_* and n are adjustable in fitting the 3:2 ratio, which play very important roles in fitting the 3:2 ratio of HFQPOs.

In this paper a non-axisymmetric magnetic field on the BH horizon is assumed in an ad hoc way. Unfortunately, we cannot give a good explanation for its existence at present. Probably the non-axisymmetric magnetic field on the BH horizon arises from non-axisymmetric accretion disc, since the magnetic field is brought and held by the surrounding magnetized disc. We hope to approach this difficult task in future.

Acknowledgments. We thank the anonymous referee for his/her many constructive suggestions. This work is supported by the National Natural Science Foundation of China under Grant Numbers 10173004, 10373006 and 10121503.

REFERENCES

Abramowicz, M. A., & Kluzniak, W., 2001, A&A, 374, L19

Bateman G., *MHD Instabilities*, 1978, (Cambridge: The MIT Press)

Blandford, R. D., & Znajek, R. L. 1977, MNRAS, 179, 433

Blandford, R. D., 1999, in ASP Conf. Ser. 160, Astrophysical Discs: An EC Summer School, ed. J. A. Sellwood & J. Goodman (San Francisco: ASP), 265 (B99)

Bradt, H. V., Rothschild, R. E., & Swank, J. H., 1993, A&AS, 97, 355

Frank J., King A. R., & Raine D. L., 1992, *Accretion Power in Astrophysics*, 2nd ed., Cambridge Univ. Press, Cambridge

Galsgaard, K., & Parnell, C., *Procedings of SOHO 15 Coronal Heating, ESA publication*, astro-ph/0409562

Greene, J., Bailyn, C. D., & Orosz, J. A., 2001, ApJ, 554, 1290

Gruzinov A., 1999a, astro-ph/9908101

Gruzinov A., 1999b, astro-ph/9910335

Haardt F., & Maraschi L., 1991, ApJ, 380, L51

Homan, J., 2004, astro-ph/0406334

Kadomtsev B. B., 1966, Rev. Plasma Phys., 2, 153

Liang E. P. T., Price R. H., 1977, ApJ, 218, 247

Li, L. -X., 2000a, ApJ, 533, L115

Li, L. -X., 2000b, ApJ, 531, L111

Li, L. -X., 2002a, ApJ, 567, 463

Li, L. -X., 2002b, A&A, 392, 469

Macdonald D., & Thorne K. S., 1982, MNRAS, 198, 345 (MT82)

McClintock J. E., & Remillard, R. A., 2003, in "Compact Stellar X-ray Sources," eds. W. H. G. Lewin & M. van der Klis, (Cambridge U. Press), in press; astro-ph/0306213 (MR03)

Mirabel & Rodrigues, 1998, Nature, 392, 673

Mirabel & Rodrigues, 1999, ARA&A, 37, 409

Miller, J. M. et al., 2001, ApJ, 563, 928

Novikov, I. D., & Thorne, K. S., 1973, in *Black Holes*, ed. Dewitt C, (Gordon and Breach, New York) p.345

Orosz et al., 2002, ApJ, 568, 845

Peter, H., Gudiksen, B., & Nordlund, A., *Procedings of SOHO 15 Coronal Heating*, ESA publication, astro-ph/0409504

Poutanen, J., & Fabian, A. C., 1999, MNRAS, 306, L31

Pozdnyakov, L. A., Sobol, I. M., & Sunyaev, R. A., 1983, in Sov. Sci. Rev./Sec. E: Astr. Space Phys. Rev., Vol. 2, ed. R. A. Sunyaev (New York: Harwood), 189

Remillard, R. A., et al., 1999, ApJ, 522, 397

Remillard, R. A., et al., 2002, ApJ, 564, 962

Remillard, R. A., et al., 2004, astro-ph/0407025

Sobczak, G. I., et al., 1999, ApJ, 520, 776

Strohmayer, T. E., 2001a, ApJ, 552, L49

Strohmayer, T. E., 2001b, ApJ, 554, L169

Tomimatse A., Matsuoka T., & Takahashi M., 2001, Phys. Rev. D64, 123003

Wagoner, R. V., Silbergleit, A. S., & Ortega-Rodriguez, M. 2001, ApJ, 559, L25

Wang, D.-X., Xiao K., & Lei W.-H. 2002, MNRAS, 335, 655 (W02)

Wang, D.-X., Lei W.-H., & Ma R.-Y., 2003, MNRAS, 342, 851 (W03a)

Wang D.-X., Ma R.-Y., Lei W.-H., & Yao G.-Z., 2003a, MNRAS, 344, 473 (W03b)

Wang D.-X., Ma R.-Y., Lei W.-H., & Yao G.-Z., 2003b, ApJ, 595, 109 (W03c)

Wang D.-X., Ma R.-Y., Lei W.-H., & Yao G.-Z., 2004, ApJ, 601, 1031 (W04)

Wilms J. et al., 2001, MNRAS, 328, L27

Zhang, S. N., Cui, W., & Chen W., 1997, ApJ, 482, L155

Zhang S. N. et al., 2000, Science, 287, 1239