



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2023 – 03:07 PM EDT

PDB ID : 1PL4  
Title : Crystal Structure of human MnSOD Y166F mutant  
Authors : Fan, L.; Tainer, J.A.  
Deposited on : 2003-06-06  
Resolution : 1.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4690 (1.50-1.46)
Clashscore	141614	4955 (1.50-1.46)
Ramachandran outliers	138981	4846 (1.50-1.46)
Sidechain outliers	138945	4844 (1.50-1.46)
RSRZ outliers	127900	4614 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	198	 90% 9% .
1	B	198	 85% 13% ..
1	C	198	 85% 14% ..
1	D	198	 89% 9% ...

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6933 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Superoxide dismutase [Mn], mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	198	1572	1008	275	285	4	0	0	0
1	B	196	1553	996	271	282	4	0	0	0
1	C	197	1562	1002	273	283	4	0	0	0
1	D	197	1562	1002	273	283	4	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	166	PHE	TYR	engineered mutation	UNP P04179
B	166	PHE	TYR	engineered mutation	UNP P04179
C	166	PHE	TYR	engineered mutation	UNP P04179
D	166	PHE	TYR	engineered mutation	UNP P04179

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mn	0	0
			1	1		
2	B	1	Total	Mn	0	0
			1	1		
2	C	1	Total	Mn	0	0
			1	1		
2	D	1	Total	Mn	0	0
			1	1		


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	188	Total 188	O 188	0	0
3	B	171	Total 171	O 171	0	0
3	C	141	Total 141	O 141	0	0
3	D	180	Total 180	O 180	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Superoxide dismutase [Mn], mitochondrial

Chain A:  90% 9%





- Molecule 1: Superoxide dismutase [Mn], mitochondrial

Chain B:  85% 13%




- Molecule 1: Superoxide dismutase [Mn], mitochondrial

Chain C:  85% 14%



- Molecule 1: Superoxide dismutase [Mn], mitochondrial

Chain D:  89% 9%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.69Å 77.71Å 135.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.47 19.85 – 1.47	Depositor EDS
% Data completeness (in resolution range)	84.0 (30.00-1.47) 90.3 (19.85-1.47)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.91 (at 1.47Å)	Xtrriage
Refinement program	SHELXL	Depositor
R, $R_{free}$	0.185 , 0.237 0.192 , 0.200	Depositor DCC
$R_{free}$ test set	6188 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.1	Xtrriage
Anisotropy	0.208	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 57.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6933	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 57.30 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4150e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/1618	1.19	10/2195 (0.5%)
1	B	0.45	0/1599	1.11	7/2173 (0.3%)
1	C	0.44	0/1608	1.24	16/2184 (0.7%)
1	D	0.51	0/1608	1.28	11/2184 (0.5%)
All	All	0.47	0/6433	1.21	44/8736 (0.5%)

There are no bond length outliers.

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	192	ARG	CD-NE-CZ	19.48	150.88	123.60
1	D	192	ARG	NE-CZ-NH1	-16.12	112.24	120.30
1	A	78	TRP	CD1-NE1-CE2	10.12	118.11	109.00
1	A	45	TYR	CB-CG-CD2	9.50	126.70	121.00
1	D	31	HIS	CG-ND1-CE1	9.24	121.13	108.20
1	C	34	TYR	CG-CD1-CE1	8.25	127.90	121.30
1	D	192	ARG	NE-CZ-NH2	8.23	124.42	120.30
1	D	176	TYR	CB-CG-CD1	8.09	125.85	121.00
1	C	78	TRP	CD1-NE1-CE2	8.02	116.21	109.00
1	A	192	ARG	NE-CZ-NH1	7.69	124.14	120.30
1	B	78	TRP	CD1-NE1-CE2	7.35	115.62	109.00
1	B	99	ARG	NE-CZ-NH1	7.35	123.97	120.30
1	C	78	TRP	CB-CG-CD1	-7.19	117.65	127.00
1	B	144	ASP	CB-CG-OD1	6.92	124.53	118.30
1	C	34	TYR	CD1-CE1-CZ	-6.48	113.97	119.80
1	A	78	TRP	CG-CD1-NE1	-6.42	103.68	110.10
1	D	31	HIS	ND1-CG-CD2	-6.37	97.08	106.00
1	C	34	TYR	CZ-CE2-CD2	6.32	125.49	119.80
1	A	31	HIS	CG-ND1-CE1	6.24	116.94	108.20
1	C	34	TYR	CG-CD2-CE2	-6.21	116.33	121.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	45	TYR	CB-CG-CD2	6.13	124.68	121.00
1	A	45	TYR	CB-CG-CD1	-6.00	117.40	121.00
1	A	78	TRP	CB-CG-CD1	-5.98	119.22	127.00
1	C	176	TYR	CB-CG-CD1	5.87	124.52	121.00
1	B	78	TRP	CG-CD1-NE1	-5.78	104.32	110.10
1	C	151	GLY	C-N-CA	5.75	136.08	121.70
1	D	144	ASP	CB-CG-OD2	-5.68	113.19	118.30
1	C	99	ARG	NE-CZ-NH1	5.59	123.09	120.30
1	D	176	TYR	CB-CG-CD2	-5.53	117.68	121.00
1	B	78	TRP	NE1-CE2-CZ2	5.39	136.33	130.40
1	C	78	TRP	CG-CD1-NE1	-5.37	104.73	110.10
1	B	100	ASP	CB-CG-OD2	5.34	123.11	118.30
1	A	78	TRP	NE1-CE2-CD2	-5.34	101.96	107.30
1	D	31	HIS	CB-CG-CD2	5.33	147.31	130.80
1	D	91	GLY	C-N-CA	5.32	135.00	121.70
1	C	176	TYR	CG-CD1-CE1	5.32	125.55	121.30
1	A	99	ARG	NE-CZ-NH1	5.27	122.94	120.30
1	C	132	ARG	NE-CZ-NH1	5.27	122.93	120.30
1	D	99	ARG	CD-NE-CZ	5.25	130.95	123.60
1	C	90	LYS	CA-CB-CG	5.24	124.93	113.40
1	C	31	HIS	CG-ND1-CE1	5.21	115.50	108.20
1	B	78	TRP	CB-CG-CD1	-5.20	120.24	127.00
1	C	99	ARG	CD-NE-CZ	5.18	130.85	123.60
1	A	170	LYS	CA-C-N	5.05	128.30	117.20

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1572	0	1528	9	0
1	B	1553	0	1502	19	0
1	C	1562	0	1515	11	0
1	D	1562	0	1515	10	0
2	A	1	0	0	0	0

Continued on next page...



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	188	0	0	3	0
3	B	171	0	0	6	0
3	C	141	0	0	2	0
3	D	180	0	0	3	0
All	All	6933	0	6060	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:109:GLU:HG2	3:B:616:HOH:O	1.93	0.67
1:C:1:LYS:HB3	3:C:730:HOH:O	1.96	0.65
1:A:60:LEU:HG	3:A:639:HOH:O	1.96	0.64
1:C:47:GLU:O	1:C:51:LYS:HG3	1.99	0.63
1:C:104:PHE:CE2	1:C:108:LYS:HD2	2.34	0.62
1:A:171:ASN:HB3	3:A:867:HOH:O	1.99	0.61
1:B:178:LYS:O	1:B:178:LYS:HE2	2.01	0.60
1:B:5:PRO:O	1:B:27:HIS:HE1	1.86	0.58
1:B:11:TYR:OH	1:B:27:HIS:HD2	1.88	0.56
1:C:8:PRO:HD2	1:C:9:TYR:CE2	2.40	0.56
1:B:31:HIS:NE2	1:B:71:HIS:HD2	2.05	0.54
1:B:49:LEU:HD23	1:C:46:GLN:OE1	2.08	0.53
1:A:193:TYR:CZ	1:A:197:LYS:HD2	2.44	0.53
1:D:31:HIS:CE1	1:D:71:HIS:HA	2.44	0.53
1:C:193:TYR:CZ	1:C:197:LYS:HD2	2.44	0.52
1:B:65:LYS:HE3	1:B:144:ASP:OD1	2.10	0.52
1:A:171:ASN:HB3	3:B:539:HOH:O	2.11	0.51
1:B:24:GLN:HG2	3:B:967:HOH:O	2.10	0.50
1:D:8:PRO:HD2	1:D:9:TYR:CE2	2.46	0.50
1:A:104:PHE:CE2	1:A:108:LYS:HD2	2.48	0.49
1:B:43:GLU:O	1:B:47:GLU:HG3	2.13	0.49
1:B:8:PRO:HD2	1:B:9:TYR:CE2	2.47	0.49
1:D:92:GLU:OE1	1:D:197:LYS:HE3	2.12	0.49
1:D:121:SER:HB3	1:D:161:TRP:CE2	2.49	0.48
1:B:108:LYS:HG2	1:B:181:TRP:CH2	2.49	0.48
1:A:119:GLN:HB3	1:B:66:PHE:CE1	2.50	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:131:GLU:HG3	3:D:849:HOH:O	2.16	0.45
1:C:51:LYS:HE3	3:C:831:HOH:O	2.17	0.45
1:C:103:SER:OG	1:C:106:LYS:HE3	2.16	0.45
1:D:106:LYS:O	1:D:110:LYS:HG2	2.18	0.44
1:B:109:GLU:HG2	3:B:536:HOH:O	2.18	0.44
1:D:61:GLN:N	3:D:889:HOH:O	2.51	0.43
1:B:121:SER:HB3	1:B:161:TRP:CE2	2.53	0.43
1:B:46:GLN:NE2	1:C:49:LEU:HD23	2.35	0.42
1:A:121:SER:HB3	1:A:161:TRP:CE2	2.55	0.42
3:A:655:HOH:O	1:B:170:LYS:HE3	2.18	0.42
1:B:71:HIS:HE1	3:B:835:HOH:O	2.01	0.41
1:C:120:GLY:HA2	1:D:161:TRP:CH2	2.55	0.41
1:A:129:ASN:C	1:A:129:ASN:HD22	2.23	0.41
1:B:178:LYS:HD3	3:B:788:HOH:O	2.20	0.41
1:C:121:SER:HB3	1:C:161:TRP:CE2	2.56	0.41
1:D:9:TYR:HB2	1:D:13:ALA:CB	2.49	0.41
1:D:170:LYS:N	1:D:170:LYS:HD2	2.35	0.41
1:D:171:ASN:HB2	3:D:971:HOH:O	2.20	0.40
1:A:47:GLU:O	1:A:51:LYS:HG3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/198 (99%)	190 (97%)	6 (3%)	0	100	100
1	B	194/198 (98%)	189 (97%)	5 (3%)	0	100	100
1	C	195/198 (98%)	189 (97%)	6 (3%)	0	100	100
1	D	195/198 (98%)	190 (97%)	5 (3%)	0	100	100
All	All	780/792 (98%)	758 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/162 (100%)	160 (99%)	2 (1%)	71	46
1	B	160/162 (99%)	157 (98%)	3 (2%)	57	26
1	C	161/162 (99%)	156 (97%)	5 (3%)	40	10
1	D	161/162 (99%)	157 (98%)	4 (2%)	47	16
All	All	644/648 (99%)	630 (98%)	14 (2%)	52	20

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	LYS
1	A	129	ASN
1	B	90	LYS
1	B	129	ASN
1	B	178	LYS
1	C	16	PRO
1	C	84	ASN
1	C	90	LYS
1	C	177	LEU
1	C	185	ASN
1	D	1	LYS
1	D	31	HIS
1	D	106	LYS
1	D	170	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	A	119	GLN
1	A	129	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	147	GLN
1	B	27	HIS
1	B	71	HIS
1	B	129	ASN
1	B	147	GLN
1	C	21	GLN
1	C	67	ASN
1	C	147	GLN
1	C	185	ASN
1	D	24	GLN
1	D	142	ASN
1	D	147	GLN
1	D	188	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	198/198 (100%)	0.07	0 <a href="#">100</a>   <a href="#">100</a>	10, 14, 27, 36	0
1	B	196/198 (98%)	0.14	2 (1%) <a href="#">82</a>   <a href="#">84</a>	10, 16, 28, 53	0
1	C	197/198 (99%)	0.20	3 (1%) <a href="#">73</a>   <a href="#">77</a>	10, 16, 32, 50	0
1	D	197/198 (99%)	-0.01	0 <a href="#">100</a>   <a href="#">100</a>	10, 14, 26, 46	0
All	All	788/792 (99%)	0.10	5 (0%) <a href="#">89</a>   <a href="#">91</a>	10, 15, 28, 53	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	90	LYS	3.3
1	B	8	PRO	2.6
1	C	197	LYS	2.5
1	C	93	LEU	2.1
1	B	1	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MN	B	200	1/1	0.99	0.04	10,10,10,10	0
2	MN	D	200	1/1	0.99	0.03	9,9,9,9	0
2	MN	C	200	1/1	1.00	0.03	10,10,10,10	0
2	MN	A	200	1/1	1.00	0.03	11,11,11,11	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.