



HAL
open science

Evolution: Back to heavy bones in salty seas

Alexandra Houssaye

► **To cite this version:**

Alexandra Houssaye. Evolution: Back to heavy bones in salty seas. *Current Biology - CB*, 2022, 32 (1), pp.R42-R44. 10.1016/j.cub.2021.11.049 . hal-03820094

HAL Id: hal-03820094

<https://hal.science/hal-03820094>

Submitted on 18 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evolution: Back to heavy bones in salty seas

Alexandra Houssaye

Departement Adaptations du vivant, UMR 7179 CNRS/Museum National d'Histoire Naturelle, 57 rue Cuvier CP-55, 75005 Paris, France

Correspondence: houssaye@mnhn.fr

<https://doi.org/10.1016/j.cub.2021.11.049>

[In amniote vertebrates that transitioned from land to sea, bone mass typically increases and later decreases](#)

[as active swimming evolves. A new study now has found that heavy bones re-evolved in some fossil marine](#)

[mammals, suggesting that this trajectory can be reversed.](#)

Amniotes — the vertebrate clade comprising reptiles (including birds) and mammals — evolved on land but various groups became aquatic from the Early Permian (about 300 million years ago) onward and especially after the EndPermian mass extinction (about 250 million years ago). These numerous land-to-sea transitions occurred independently in various mammal and reptile lineages, such as cetaceans, sirenians, ichthyosaurs or plesiosaurs. Such a change of milieu, and thus of environmental constraints, is naturally associated with various morphological, locomotor, physiological and reproductive modifications¹, including nearly always changes in the internal structure of bones². Two main specializations stand out: first, an increase in bone mass, generally in less active swimmers hovering at shallow depths. Such heavy bones serve as ballast for hydrostatic control of buoyancy when diving and for maintaining a horizontal orientation of the body while swimming. Second, in active swimmers with great maneuvering and acceleration capabilities, which rely on hydrodynamic control of buoyancy and body trim, bones become spongy, showing a cancellous organization^{3,4}. Increased bone mass characterizes especially the early stages of a lineage's transition to an open ocean lifestyle. The affected animals inhabit shallow waters, show a more 'terrestrial-like' morphology and rely on drag-based propulsion with paddling appendages. Conversely, later evolutionary stages are characterized by a cancellous organization in more active swimmers with streamlined bodies, paddles and reliance on an oscillating hydrofoil, as in whales and mosasaurs (a clade of Mesozoic marine reptiles)^{5,6}. This apparently stepwise trajectory has led to the suggestion that a compact structure was a necessary transitional step towards a cancellous organization⁷. This evolutionary shift in bone microanatomy was thus considered unidirectional.

However, in a new study in this issue of *Current Biology*, Leonard Dewaele, Vivian de Buffrenil and colleagues [8](#) highlight the first known reversal of this evolutionary trajectory, with the re-emergence of increased bone mass in pinnipeds and cetaceans from the Paratethys sea during the Miocene (23–5 million years ago).

Increased bone mass consists of an increase in bone compactness through unbalanced remodeling (a process in which old bone is resorbed and new bone is deposited, and which maintains or adapts bone structure and repairs the skeleton) with inhibited resorption or excessive secondary bone deposition. It can also be caused by additional peripheral deposits⁴. In the fossil record, localized bone mass increase is observed in many early whales from the Eocene (56–34 million years ago), with varying degrees of intensity and distribution, but vanishes later in their evolutionary history^{9,10}. The fossil record thus appears to be consistent with the unidirectional trajectory of bone mass evolution.

In their new study, however, Dewaele and colleagues⁸, based on an analysis of the inner structure of ribs and long bones, show that bone mass increased in pinnipeds and cetaceans from the Miocene of the Paratethys sea. The occurrence of this specialization in both clades of whale — odontocetes and mysticetes — clearly indicates a reversal from a cancellous to a compact condition in the cetaceans analyzed ([Figure 1](#)). For pinnipeds (seals, sea lions, walruses), data are scarcer: modern pinnipeds and the few fossil phocids for which data are available⁸ show a cancellous organization. However, Dewaele and colleagues⁸ observed a very high bone mass in the Miocene Paratethys

R42 *Current Biology* 32, R32–R54, January 10, 2022 ^a 2021 Published by Elsevier Inc.



Dispatches

phocids. In addition, a Miocene phocine from Belgium (*Nanophoca vitulinoides*) shows a surprising case of generalized bone mass increase in its postcranial skeleton¹¹. Unfortunately, the phylogenetic relationships between all of these phocid species remain unknown (especially due to the lack of cranial material) and conditions in other early pinnipeds are not documented so the shift from a cancellous to a compact structure, although highly likely to have occurred

several times, cannot be ascertained in this lineage.

The independent, yet concomitant reappearance of this specialization in these three distinct lineages — phocids, baleen whales and toothed whales — with different functional requirements, such as swimming, diving and acceleration, clearly highlights its convergent nature and seems to reflect strong selective pressure. All the specimens studied by Dewaele and colleagues⁸ come from the Miocene Paratethys, suggesting local, convergent evolution. The Paratethys was a shallow sea that extended from the European Alps through Central Europe to the Aral Sea in Central Asia. Tectonics and sea-level variation resulted in various degrees of extension of the Paratethys — it reached its largest size during the late Miocene — and connections with other seas. Today, the remnants of the Paratethys are the Black Sea, the Caspian Sea and the almost vanished Aral Sea, as well as a few lakes¹². During the middle Miocene, between 13.8 and 13.4 million years ago, connection between the Mediterranean and the Paratethys was lost leading to the latter becoming hypersaline, a period known as the Badenian Salinity Crisis¹³. However, increased bone mass is known not to be directly influenced by local environmental conditions, as taxa with varied bone structures are observed together in similar outcrops or present-day habitats². Instead, different bone structures reflect species-specific adaptations. Dewaele and colleagues⁸ do not call this statement into question. Rather, the authors conclude that while local environmental conditions appear to be the cause of this increase in bone mass, it is likely through the response of the skeleton to a change in functional constraints. Indeed, hypersalinity increases the buoyancy that aquatic mammals have to overcome when they dive. Greater bone mass would have been favored, resulting in this evolutionary shift.

The shift in bone mass observed by Dewaele and colleagues⁸ concerns marine mammals that lived approximately one to six million years after the Badenian Salinity Crisis, suggesting it was a relatively long-term adaptive process. Moreover, it is also found in forms that lived further East, where waters were brackish and not hypersaline. Although the increase in bone mass evolved during the crisis, this feature remained characteristic of the Miocene aquatic

mammal fauna endemic to the Paratethys, particularly through the dispersal of the lineages involved. The reason for a potentially similar independent change in the Phocid *Nanophoca* from the Miocene of Belgium¹¹, and thus not from the Paratethys, should be investigated to better understand the selective pressures driving this cancellous-compact shift and how it took place.

The increase in bone mass evolved comparatively fast, over less than a million years, which is perhaps not surprising. The internal structure of bone is known to be plastic and appears to evolve more rapidly than bone shape, as evidenced by significant changes in bone inner structure in taxa with otherwise limited adaptations to an aquatic life-style^{2,14}. Dewaele and colleagues⁸ highlight that heavy bones persisted after the crisis was over and salinity returned to normal. The reasons for this persistence remain unclear; perhaps they were an exaptation to benthic foraging, as mentioned by the authors⁸, or a more general lack of strong selective pressure against high bone mass.

The study by Dewaele and colleagues⁸ on the bone structure of fossil seals, dolphins and whales from the Miocene Paratethys add a twist to our understanding of adaptation to the aquatic realm in these mammal lineages. But more generally, this first documented

Humeri

Ribs

Pakicetidae

Protocetidae

Pachyacanthus

Platanista

Delphinus

Phocoena

Delphinida indet.

Cethotheriidae indet.

Cethotheriidae indet.

Basilosauridae

* *

*

Current Biology

Figure 1. Shifts in bone microanatomy during whale evolution.

Black and white representation of humerus and rib sections from various cetaceans for which data are available plotted on a simplified phylogeny (modified from Gatesy et al.¹⁵) showing the independent shifts from a cancellous to a compact structure (/). *Specimens from the Miocene Paratethys.

Current Biology 32, R32–R54, January 10, 2022 R43



Dispatches

shift from a cancellous to a compact bone structure extends our understanding of the development of these specializations in amniotes that adapted to an aquatic

lifestyle.

REFERENCES

1. Kelley, N.P., and Pyenson, N.D. (2015). Evolutionary innovation and ecology in marine tetrapods from the Triassic to the Anthropocene. *Science* 348, 6232.
2. Houssaye, A., and de Buffrenil, V. (2021). Bone histology and the adaptation to aquatic life in tetrapods. In *Vertebrate Skeletal Histology and Paleohistology*, V. de Buffrenil, A.J. de Ricqlès, L. Zylberberg, and K. Padian, eds. (CRC Press), pp. 744–756.
3. Fish, F.E. (1996). Transitions from drag-based to lift-based propulsion in mammalian swimming. *Am. Zool.* 36, 628–641.
4. de Ricqlès, A., and de Buffrenil, V. (2001). Bone histology, heterochronies and the return of tetrapods to life in water: where are we? In *Secondary Adaptation of Tetrapods to Life in Water*, J.M. Mazin and V. de Buffrenil, eds. (München), pp. 289–310.
5. Gray, N., Kainec, K., Madar, S., Tomko, L., and Wolfe, S. (2007). Sink or swim? Bone density as a mechanism for buoyancy control in early cetaceans. *Anat. Rec.* 290, 638–653.
6. Houssaye, A., Lindgren, J., Pellegrini, R., Lee, A.H., Germain, D., and Polcyn, M.J. (2013). Microanatomical and histological features in the long bones of mosasaurine mosasaurs (Reptilia, Squamata) — implications for aquatic adaptation and growth rates. *PLoS One* 8, e76741.
7. Zangerl, R., and Peyer, B. (1935). Die Triasfauna der Tessiner Kalkalpen IX. *Pachypleurosaurus edwardsi*, *Cornalia* sp. *Osteologie-Variationsbreite-Biologie. Abh. Schweiz. Palaeontol. Ges.* 56, 1–80.
8. Dewaele, L., Gol'din, P., Marx, F.G., Lambert, O., Laurin, M., Obada, T., and de Buffrenil, V. (2022). Hypersalinity drives convergent bone mass increases in Miocene marine mammals from the Paratethys. *Curr. Biol.* 32, 248–255.
9. Houssaye, A., Tafforeau, P., De Muizon, C., and Gingerich, P.D. (2015). Transition of Eocene whales from land to sea: evidence from bone microstructure. *PLoS One* 10, e0118409.
10. Canoville, A., and Laurin, M. (2010). Evolution of humeral microanatomy and lifestyle in amniotes, and some comments on palaeobiological inferences. *Biol. J. Linn. Soc.* 100, 384–406.
11. Dewaele, L., Lambert, O., Laurin, M., De Kock, T., Louwye, S., and de Buffrenil, V. (2019). Generalized osteosclerotic condition in the skeleton of *Nanophoca vitulinoides*, a dwarf seal from the Miocene of Belgium. *J. Mammal. Evol.* 26, 517–543.
12. Rogli, F. (1999). Palaeogeographic considerations for Mediterranean and Paratethys seaways. *Ann. Nat. Mus. Wien.* 99, 279–310.
13. Palcu, D.V., Patina, I.S., Şandric, I., Lazarev, S., Vasiliev, I., Stoica, M., and Krijgsman, W. (2021). Late Miocene megalake regressions in Eurasia. *Sci. Rep.* 11, 11471.
14. Amson, E., de Muizon, C., Laurin, M., Argot, C., and de Buffrenil, V. (2014). Gradual adaptation of bone structure to aquatic lifestyle in extinct sloths from Peru. *Proc. Biol. Sci.* 281, 20140192.
15. Gatesy, J., Geisler, J.H., Chang, J., Buell, C., Berta, A., Meredith, R.W., Springer, M.S., and McGowen, M.R. (2013). A phylogenetic blueprint for a modern whale. *Mol. Phylogenet. Evol.* 66, 479–506.