

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/127249/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Graham, Meghan R, Tierney, Stephanie, Chisholm, Amy and Fox, John R E 2020. The lived experience of working with people with eating disorders: A meta-ethnography. *International Journal of Eating Disorders* 53 (3) , pp. 422-441. 10.1002/eat.23215

Publishers page: <http://dx.doi.org/10.1002/eat.23215>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1

The lived experience of working with people with eating disorders:

2

A meta-ethnography

3

Meghan R. Graham^a, Stephanie Tierney^b, Amy Chisholm^c, John R.E. Fox^{d*}

4

5

^aDepartment of Psychology, Royal Holloway, University of London, Egham, England

6

^bNuffield Department of Primary Care Health Sciences, University of Oxford, Oxford,

7

England

8

^cVincent Square Eating Disorders Service, London, England

9

^dSchool of Psychology, Cardiff University, Cardiff, Wales

10

11

* Corresponding author: John R.E. Fox, South Wales Clinical Psychology Doctoral

12

Programme, School of Psychology, Cardiff University, Cardiff, CF10 3AT, UK

13

E-mail address: foxj10@cardiff.ac.uk

14

Declaration of interests: None

15

Word count of Abstract: 209

16

Word count of Manuscript: 7,466

17

18

19

20

Abstract

1

2

3 Objective: Working with people with eating disorders (EDs) is known to elicit strong
4 emotional reactions, and the therapeutic alliance has been shown to affect outcomes with this
5 clinical population. As a consequence, it is important to understand healthcare professionals’
6 (HCPs’) experiences of working with this client group.

7 Method: A meta-synthesis was conducted of qualitative research on HCPs’ lived experiences
8 of working with people with EDs. The results from the identified studies were analyzed using
9 Noblit and Hare’s meta-ethnographic method. Data were synthesized using reciprocal
10 translation, and a line of argument was developed.

11 Results: Thirty-seven studies met the inclusion criteria. Reciprocal translation resulted in a
12 key concept: “Coping with caring without curing.” This was underpinned by the following
13 third-order concepts: (a) “The dissonance and discomfort of being a helper struggling to
14 help,” (b) “Defending against the dissonance,” and (c) “Accepting the dissonance to provide
15 safe and compassionate care.” These concepts were used to develop a line-of-argument
16 synthesis, which was expressed as a new model for understanding HCPs’ experiences of
17 working with people who have an ED.

18 Discussion: While the conflict associated with being a helper struggling to help led some
19 HCPs to avoid and blame people with EDs, others adopted a compassionate stance
20 characterized by humanity, humility, balance, and awareness.

21

22

23 Key words:

24 Systematic review

WORKING WITH EATING DISORDERS

1 Meta-ethnography

2 Eating disorders

3 Healthcare professionals

4 Qualitative research

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

1

2

Introduction

3

Eating disorders (EDs) have a reputation for being challenging to treat (Startup, Mountford,

4

Lavender, & Schmidt, 2015) in the context of high levels of risk, relapse and resistance

5

(Jansen, 2016; Reilly, Anderson, Gorrell, Schaumberg, & Anderson, 2017; Zaitsoff, Pullmer,

6

Cyr, & Aime, 2015). Anorexia nervosa (AN) and bulimia nervosa (BN) are associated with

7

many medical complications, some of which can be fatal (Westmoreland, Krantz, & Mehler,

8

2016), which means that healthcare professionals (HCPs) working with this client group are

9

required to manage high levels of physical and psychiatric risk concurrently (Le, Barendregt,

10

Hay, & Mihalopoulos, 2017). Managing significant risk is linked to emotional exhaustion,

11

particularly where HCPs have unrealistic self-expectations (Kleespies & Dettmer, 2000).

12

Recovery is often elusive, with only 31% of people with AN and 68% of those with BN

13

recovering within nine years (Eddy et al., 2017). A further challenge of the work is many

14

clients value aspects of their ED (Fassino & Abbate-Dage, 2013) and therefore regard HCPs

15

as a threat, which fosters inaccurate self-report and disengagement (Gregertsen, Mandy, &

16

Serpell, 2017). Moreover, there are few evidence-based treatments for EDs, particularly AN

17

(Lock, 2015). Given that thinness can elicit envy in societies in which it is culturally

18

endorsed (Beggan, & DeAngelis, 2015), HCPs may have mixed feelings towards

19

underweight clients, which can further complicate the work. Concerns have been raised that

20

the training of HCPs working with ED clients is inadequate (e.g., Hay, Darby, & Mond,

21

2007; Jones, Saeidi, & Morgan, 2013).

22

HCPs caring for people with EDs frequently report feelings of anxiety, distress, anger,

23

and exasperation (Golan, Yaroslavski, & Stein, 2009; Land, 2004), and often have

24

stigmatizing views of these clients (Raveneau, Feinstein, Rosen, & Fisher, 2014). Lower

25

functioning and higher levels of dysregulation among service users (Satir et al., 2009),

WORKING WITH EATING DISORDERS

1 clinician inexperience, larger caseloads and a diagnosis of AN are independently associated
2 with more negative reactions among HCPs (Franko & Rolfe, 1996). Managing negative
3 reactions via externalization of EDs, self-monitoring and discussion in supervision can reduce
4 staff burnout (Golan et al., 2009). There is evidence that acceptance-based interventions are
5 helpful for staff working with other potentially challenging populations, such as people with
6 intellectual disabilities (Noone & Hastings, 2009).

7 Understanding the lived experiences of staff in ED services is important given that the
8 clinician-client alliance has been shown to affect treatment outcomes (Graves et al., 2017).
9 Several existing reviews have examined this topic. In a mixed-methods review of
10 knowledge, attitudes and perceived challenges among HCPs working with people with EDs,
11 Seah et al. (2017) identified issues of limited knowledge and confidence, negative attitudes,
12 service-user non-adherence, high workloads, miscommunication among teams, and being
13 personally affected by the work. Thompson-Brenner, Satir, Franko, and Herzog (2012)
14 conducted a mixed-methods review on HCPs' reactions to people with EDs and highlighted
15 feelings of worry, frustration, hopelessness and incompetence. Employing thematic
16 synthesis, Sibeoni, Orri, Lachal, Moro and Revah-Levy (2017) explored views of the
17 treatment of adolescent AN and found that HCPs reported focusing on weight gain and
18 behaviors, removing control to maintain physical safety, and struggling to develop
19 therapeutic relationships due to mistrust, perceived manipulation, and a battle for control.

20 Given that these reviews lacked detail, all but one were aggregative, and the
21 interpretative synthesis by Sibeoni et al. (2017) focused on one ED and age group, a meta-
22 ethnography was planned on the lived experiences of diverse HCPs caring for service users of
23 all ages with different EDs. Whereas aggregative reviews summarize existing literature in an
24 additive manner, meta-ethnography goes beyond the original data to develop new
25 understandings (Barnett-Page & Thomas, 2009) by building an explanatory theory or model

WORKING WITH EATING DISORDERS

1 (Bondas & Hall, 2007^a; Walsh & Downe, 2005). All EDs were incorporated in the search
2 terms because they are often grouped in the literature; this grouping accords with the
3 transdiagnostic model of AN, BN and most Other Specified Feeding and Eating Disorders
4 and Unspecified Feeding and Eating Disorders as sharing a core psychopathology of over-
5 evaluation of the importance of eating, shape and weight, and their control (Fairburn, 2008).
6 The definition of lived experience that guided study selection was: “The detailed, nuanced,
7 and subjective experience, including individual perceptions, meanings, understandings,
8 descriptions, and felt somatic sense of an experience from the first-hand point of view of a
9 particular person” (Mertens, 2005, as cited in Palmer, 2015, p.123).

10 The meta-ethnography aimed to generate a model of the processes involved in
11 working with people with EDs, drawn from the lived experiences of HCPs, to improve
12 clinical practice and inform service development.

13 **Method**

14 **Systematic literature search**

15 The following databases were systematically searched in March 2018: PsycINFO,
16 PubMed, and Web of Science. Three categories of search terms – eating disorders,
17 experiences and professionals – were combined using Boolean operators. No date restriction
18 was employed. Reference lists of relevant studies were manually searched for additional
19 papers.

20 **Selecting studies**

21 Inclusion and exclusion criteria were applied by the lead author (MG) to identify
22 relevant studies. The inclusion criteria were: (a) peer-reviewed empirical studies; (b) studies
23 employing a qualitative or mixed-methods design (provided the qualitative results were
24 reported separately); (c) studies focusing on EDs; (d) studies whose participants included
25 HCPs (i.e., any worker in a healthcare setting involved in the direct care and/or treatment of

WORKING WITH EATING DISORDERS

1 service users), and (e) studies focusing on the lived experience of working with people with
2 EDs. The exclusion criteria were: (a) studies in languages other than English, and (b) book
3 chapters, books, book reviews, dissertations, opinion pieces, conference presentations, and
4 meeting abstracts.

5 Figure 1 illustrates the search strategy and process for selecting studies. The lead
6 author looked at all references (during the screening of titles and abstracts, and by reading
7 full texts where applicable), and thirty per cent of articles selected to be read in full were
8 independently screened by other members of the research team. Two discrepancies in
9 decisions were resolved via discussion and re-reading articles, leading to the exclusion of an
10 additional article.

11

12 Figure 1 to go about here

13

14 **Quality appraisal**

15 To provide a guide regarding their relative rigor, the methodological quality of the
16 included studies was assessed by MG using the Critical Appraisal Skills Programme (CASP;
17 2010) checklist for qualitative research, a tool employed in previous syntheses of qualitative
18 ED studies (e.g., Espindola & Blay, 2009; Fox, Dean, & Whittlesea, 2015). The checklist has
19 10 criteria, so each study was awarded a score out of 10, with half-points granted if a
20 criterion were partially fulfilled. In the absence of standard protocols for mixed-methods
21 studies, the qualitative components of studies of this type were evaluated using the CASP
22 checklist for qualitative studies. This review followed Fox et al. (2015) in classifying studies
23 from A to C, with A denoting studies scoring 8.5 or above and carrying a low likelihood of
24 methodological flaws; B signifying studies scoring five to eight with a moderate likelihood of

WORKING WITH EATING DISORDERS

1 methodological flaws, and C indicating a score of less than five and a high likelihood of
2 methodological flaws. Twenty per cent of included studies were independently rated by other
3 members of the research team using the CASP checklist. There was unanimous agreement
4 regarding five studies and there were minor discrepancies with two; discrepancies were
5 resolved via discussion. Table 1 shows the CASP ratings.

6

7

Table 1 to go about here

8

9 **Data synthesis**

10 Noblit and Hare's (1988) meta-ethnographic method was employed because we
11 sought to develop a new conceptual understanding of the topic by configuring existing data.
12 Meta-ethnography is well-established (Bondas & Hall, 2007^b) and entails the following
13 phases:

- 14 1. Getting started: An area of intellectual interest was identified (i.e., the lived
15 experiences of HCPs working with people with EDs).
- 16 2. Deciding which studies to include: Inclusion and exclusion criteria were applied, and
17 relevant search terms employed in a range of databases (Figure 1).
- 18 3. Reading the studies: Included articles were read multiple times to identify and extract
19 methodological features, demographic information, and key concepts.
- 20 4. Determining how the studies are related: The key concepts in each study were closely
21 compared to decide relationships between them. At this stage, accounts are either
22 deemed directly comparable and capable of being "reciprocally translated" into one
23 another; in opposition to each other and suited to "refutational translation", and/or
24 cumulatively representative of a "line of argument" that "puts any similarities and
25 dissimilarities into a new interpretive context" (Noblit & Hare, 1988, p.64).

WORKING WITH EATING DISORDERS

- 1 5. Translating the studies into one another: Reciprocal translation was used to identify
2 third-order concepts (our interpretations of the original authors' interpretations) that
3 captured similarities across studies. As there were no contradictions between
4 findings, refutational translation was not necessary.
- 5 6. Synthesizing translations: Following reciprocal translation, the concepts were
6 organized into a conceptual framework, which represented a new interpretation and
7 line of argument.
- 8 7. Expressing the synthesis: The synthesis was elaborated via narrative and diagram
9 (Figure 2).

10 Throughout, the meta-ethnography was regularly discussed among the research team. An
11 inductive approach was employed, allowing findings to emerge from the data rather than be
12 determined by an a priori theoretical framework (Dillaway, Lysack, & Luborsky, 2017).

13 **Results**

14 **Quality assessment**

15 All studies were classified as A or B, with an average rating of 7.74. The primary
16 reasons for losing points were not stating/justifying the research design; not providing a
17 rationale for the method of data collection; lack of reflexivity, and presenting insufficient data
18 to support findings; this may reflect limited word-counts rather than deficiencies in execution
19 (Walsh & Downe, 2006). The overall score and classification for each study are included in
20 Table 2. Given that all studies were rated B or above, and in line with other meta-syntheses
21 in the field (Espindola & Blay, 2009; Sibeoni et al., 2017), no studies were excluded on the
22 basis of the quality ratings.

23 **Characteristics of the included studies**

24 Across the 37 studies, data were collected from 769 HCPs. They included nurses,
25 psychologists, psychiatrists, occupational therapists, dietitians, general practitioners, and

WORKING WITH EATING DISORDERS

1 medical providers such as gynecologists. Nurses were well represented in studies with mixed
2 samples and there were nine studies with solely nurse participants. Two studies exclusively
3 involved HCPs with a history of an ED. Across the studies that did not specify personal
4 experience as an inclusion criterion and reported the number of participants who disclosed
5 such experience, 43.6% of HCPs were stated to have a personal history of an ED. Across the
6 studies in which the gender composition of participants was discernible, 91.4% were female.
7 Twenty-six studies centered on experiences of working with people with EDs; the remaining
8 11 focused on experiences of working with AN. Eight studies related specifically to work
9 with adolescents. Sixteen studies recruited exclusively from an inpatient setting. Included
10 studies were undertaken in Australia, Canada, New Zealand, Norway, Singapore, Tasmania,
11 the UK, and the USA. Most data collection proceeded via interview (generally semi-
12 structured). Authors commonly employed thematic analysis. Key characteristics of included
13 studies are shown in Table 2. Despite variety in the methodologies of included studies, and
14 the range of disciplines, settings and locations represented, there was considerable
15 concordance across accounts. However, as will be outlined below, some differences were
16 apparent according to experience level, setting, and professional background.

17

18 [Table 2 to go about here](#)

19

20 **Meta-ethnography findings**

21 The stages of analysis outlined above led to the development of a key concept to
22 depict the lived experience of working with people with EDs: “Coping with caring without
23 curing.” This described how a combination of treatment refusal, chronicity, systemic
24 challenges and/or personal factors confronted HCPs with the fact that the help they offer

WORKING WITH EATING DISORDERS

1 people with EDs may be unwanted and/or insufficient. It resulted in a sense of dissonance –
2 a mismatch between their aspiration to help, and the reality. HCPs could either defend against
3 this dissonance, or face it with courage and humanity. This key concept was underpinned by
4 the three concepts that will now be presented. The number of studies that endorsed each
5 concept and sub-theme is represented in Table 3.

6 ***Concept 1: The dissonance and discomfort of being a helper struggling to help***

7 This concept, identified in 33 studies, describes the frustration, helplessness and
8 distress that often resulted from efforts to assist people with EDs, alongside a sense of
9 scrutiny in the helping role linked to the watchfulness of service users. Together, these
10 experiences represented a painful dissonance between the expectation and reality of
11 caregiving.

12 **Emotionally draining work.**

13 In 29 studies, participants reported that their work with individuals with EDs was
14 “emotionally draining” (treatment provider; Warren et al., 2012, p.183) in the context of high
15 levels of resistance and relapse, the shock of emaciation, feeling manipulated by service
16 users, and difficulties with other professionals and families. Many clinicians described
17 experiencing frustration, and some reported anger, in response to service users’ rejection of
18 treatment, denial, and relapse: “I looked after a girl who used to rip out the naso gastric
19 feeding] tube....After some time, you would get feelings of immense anger and frustration”
20 (health professional; Walker & Lloyd, 2011, p.142). Cameron et al. (1997) noted that “the
21 frustration appeared to be linked to the inability of HCPs to control the progress of the illness,
22 about ‘not being able to do enough’” (p.28). The emaciation of service users elicited fear: “I
23 can be scared and sad for not only their quality of life but the risk of a patient dying”
24 (treatment provider; Warren et al., 2008, p.39). Some participants felt manipulated and/or

WORKING WITH EATING DISORDERS

1 attacked by service users, creating a difficult dynamic: “It’s the manipulation. You think it’s
2 a personal attack against you” (nurse; Carter et al., 2012, p.551).

3 Interactions with service users’ families could be stressful if, for example, they were
4 in denial about the ED (Harken et al., 2017). Difficulties with colleagues were also reported
5 to add to the emotional demands: “We are forcing kids to do things that we do not
6 understand. The doctors are making these decisions” (nurse; Micevski & McCann, 2005,
7 p.110). Perceived lack of understanding among peers caused frustration: “The most
8 uncomfortable thing for me is the lack of education of other people around me....in particular
9 the physicians” (dietitian nutritionist; Trammell et al., 2016, p.79). One participant described
10 the emotional toll of “caring without curing” (King & Turner, 2000, p.145) as follows:
11 “Medicine is easy when you can just prescribe something and then they are better...but
12 difficult sometimes when you have to put a lot of emotion into looking after them and
13 sometimes you don’t seem to get anywhere” (general practitioner; Reid, Williams, &
14 Hammersley, 2010, p.6).

15 **Feeling helpless and deskilled.**

16 Linked to high levels of relapse and resistance, HCPs in 21 studies described feeling
17 deskilled and ill-equipped to help service users: “When I got to my internship and I did see
18 EDs, it sort of scared me. I didn’t know what to do” (dietitian nutritionist; Trammell et al.,
19 2016, p.78). A sense of helplessness was more apparent in studies involving HCPs without
20 specialist ED experience. One participant described the dissonance of feeling helpless within
21 a helping role as follows: “We are fixers and doers by our nature, that’s why we get into the
22 profession that we do, so to not have an immediate answer is difficult” (medical provider;
23 Linville et al., 2010, p.119). For a minority of HCPs, feeling helpless led to self-judgment:
24 “Most of the time you feel you are going round and round in circles so it makes you feel like
25 a useless therapist” (health professional; Walker & Lloyd, 2011, p.386). The data suggested

WORKING WITH EATING DISORDERS

1 that organizational and sociocultural factors contributed to participants' sense of limited
2 power to help. Lack of time, the most common systemic problem, particularly affected those
3 in primary care: "You can't even scratch the surface" (general practitioner; Reid, Williams, &
4 Hammersley, 2010, p.7). At a macro-level, sociocultural pressure compounded HCPs' sense
5 of helplessness to effect change: "We often feel like we are fighting a losing battle when they
6 are exposed much more consistently to messages that contradict what we promote" (treatment
7 provider; Warren et al., 2012, p.188).

8 **Watching and being watched.**

9 In 13 studies, the necessity of close observation of service users – particularly in
10 inpatient environments (Akgül et al., 2016) – and a sense of being observed in turn, resulted
11 in interpersonal mistrust amid a culture of surveillance. Participants described feeling
12 awkward about intently monitoring service users to prevent ED behaviors: "They have to
13 have somebody sitting in their room all day long with them...I feel uncomfortable for them"
14 (nurse; Harken et al., 2017, p.e38). Some noted that this entailed a lot of work: "General
15 psychiatric is easier...there's [fewer things] to look out for...Compared to the eating disorder
16 is how you eat, what you eat, what you drink, how much you drink, where are you, what you
17 doing...everything" (nurse; Seah et al., 2018, p.141). Participants articulated a feeling that
18 their behaviors and/or appearance were scrutinized by service users, resulting in fear about
19 saying or doing the wrong thing, particularly when inexperienced. For a minority of HCPs,
20 the sense of scrutiny led to self-consciousness: "I feel like they are really watching me; I was
21 trying to drink really normal" (staff member of ED unit; Long et al., 2012, p.244). Those
22 with a personal history of an ED reported an additional layer of scrutiny, from other
23 professionals; some of Williams and Haverkamp's (2015) participants described unhelpful
24 interactions with colleagues, such as being questioned about weight loss.

25 **Moral distress.**

1 **Feeling unable to help and in conflict with service users generated “dissonance”**
2 **(ED therapist; Williams & Haverkamp, 2015, p.405) for HCPs accustomed to providing**
3 **valuable and valued care**, across 10 studies. This could lead to self-judgment for the
4 perceived violation of their core values, as illustrated in the following quotation: “My heart
5 just doesn’t warm to them any more...Disgusting – sad you know, that’s not a nurse” (nurse;
6 King & Turner, 2000, p.142). In some cases, disempowering service users – to preserve their
7 physical safety – led to self-questioning: “I didn’t want this role, I’ve always fancied myself
8 as being quite therapeutic and collaborative” (nurse; Snell et al., 2010, p.354). Moral distress
9 was most prevalent among, but not unique to, nurses.

10 ***Concept 2: Defending against the dissonance***

11 This concept, evident in 23 studies, conveyed the experience of some HCPs of
12 avoiding, blaming, and battling service users. These coping strategies could defend against
13 the dissonance of caring without curing by blocking it out or attributing “the problem” to
14 service users, but simultaneously could exacerbate moral distress due to the loss of
15 unconditional positive regard.

16 **Avoiding and distancing.**

17 Across 17 studies, there was a tendency among some HCPs, particularly nurses, to
18 unconsciously or consciously avoid negative emotions, conflict, and/or association with
19 service users. For example, Ramjan (2004) reported that some nurses requested not to work
20 with adolescents with AN to distance themselves from difficult interactions. “Shutting off”
21 and being task-oriented was another strategy employed by HCPs: “You just close off
22 everything else and just take that fixed view that yes, you’re going to eat” (clinical nurse
23 specialist; Jarman et al., 1997, p.148). This could reduce frustration and distress temporarily,
24 but sometimes led to dehumanization of service users: “They become a set of symptoms and
25 things to monitor and measure” (nurse; Davey et al., 2014, p.63). Some clinicians reported

WORKING WITH EATING DISORDERS

1 binge eating to evade burnout (Warren et al., 2012), which can be understood as avoidance of
2 emotional pain. Others used their diet to distance themselves from service users: “I find
3 myself going to an extreme, at times, of eating whatever I want and thinking it is bad to ‘eat
4 healthy.’ I feel like I try to compensate for my clients’ eating-disordered views” (treatment
5 provider; Warren et al., 2008, p.37). Cameron et al. (1997) noted that participants construed
6 people with EDs as “other” to “exclude them from the domain of socially acceptable eating
7 patterns, in which most participants located themselves” (p.27). Similarly, some HCPs with a
8 personal history of ED emphasized the “normality” of their relationship with food and their
9 bodies (Rance et al., 2010; Trammell et al., 2017), in what was interpreted by Rance et al.
10 (2010) as a binary view of recovery that entailed avoidance of shared humanity with service
11 users. One therapist described how non-disclosure at work of her personal history of ED
12 created “dissonance” due to the mismatch between her secrecy and her anti-stigma message
13 to service users (Williams & Haverkamp, 2015, p.405).

14 **Battle and blame.**

15 In 11 studies, there was evidence of HCPs blaming clients and/or regarding
16 themselves as waging battle with “rebellious and dominating” service users (Long et al.,
17 2012, p.244). Comparisons of clinicians to figures of authority jarred with the notion of
18 collaborative care: “We were just like sergeant majors, we thought we were. Standing over
19 them telling them what to do” (nurse; King & Turner, 2000, p.142). King and Turner (2000)
20 remarked that HCPs pathologized typical adolescent behaviors, and service users were
21 dismissed as “vain” by a participant in Walker and Lloyd (2011). Ramjan (2004) noted a
22 tendency to judge service users as responsible for their distress, reflected in the prison
23 metaphor the participants employed, which cast clients as criminals. Cameron et al. (1997)
24 highlighted that the “moralizing discourse” employed by participants constructed individuals
25 with EDs as “difficult people, rather than as people with a difficult illness” (p.29). While a

WORKING WITH EATING DISORDERS

1 minority of clinicians were unperturbed by this dynamic, for many it was inimical to their
2 view of themselves as caring professionals.

3 ***Concept 3: Accepting the dissonance to provide safe and compassionate care***

4 This concept, depicted in 34 studies, described some HCPs' efforts to face the reality
5 of caring without curing, without blaming or avoiding service users. This entailed adopting a
6 balanced and flexible approach that was responsive to service users' needs, and their own. It
7 also required tolerating negative emotions, uncertainty and fallibility to provide
8 compassionate care in the knowledge that it may not be perceived by service users as helpful,
9 and may be rejected altogether.

10 **Balance and flexibility.**

11 Participants in 17 studies stressed the importance of balance and flexibility.

12 Brinchmann et al. (2017) described therapists in their study as navigating a "golden middle
13 way" between extremes to create a safe environment for service users that promoted
14 development. Balance was also evident in the depiction of effective HCPs as "having one
15 arm around the client while kicking them in the butt" (therapist; Oyer et al., 2016, p.128-9).
16 This approach involved taking control where necessary and then "gradually giving service
17 users bits of control back, as and when we think [they] can cope with that" (social worker;
18 Jarman et al., 1997, p.145). While participants saw rules as important to preserve safety, they
19 emphasized the therapeutic value of flexibility, for example in adapting ethical codes to suit
20 the cultural context (Tan et al., 2013). However, flexibility was anxiety-provoking because it
21 meant clinicians could not "hide behind a method which can provide them with the safety and
22 security of being in control" (Jarman et al., 1997, p.147).

23 Balance was evident in HCPs remaining motivated to support change without
24 succumbing to rescue fantasies, and in their appreciation of the multifactorial influences on
25 their experiences (Reid, Williams, & Burr, 2010). Some HCPs recognized they did not bear

WORKING WITH EATING DISORDERS

1 sole responsibility for outcomes while continuing to work hard for service users, but this was
2 highlighted as difficult: “It’s a constant challenge to find the balance between appropriate
3 attention to my clients/doing everything I should and can for them, and letting go/bearing in
4 mind that I can’t control what they do/what happens to them” (treatment provider; Warren et
5 al., 2012, p.184).

6 **Mindful awareness, acceptance and self-observation.**

7 A balanced approach was facilitated by a mindful stance, which was articulated in 10
8 studies. This manifested in HCPs taking an objective and non-judgmental perspective; facing
9 and responding to reality, and trying to monitor, tolerate, reflect on and regulate their own
10 behaviors, reactions and relationships: “At the end of the session, I reflected and realized that
11 70% of the session was me talking AT them [the client]. I am a model of relationships, and if
12 I am doing that....that is not therapeutic” (therapist; Oyer et al., 2016, p.132). Some HCPs
13 mindfully observed and positively framed the watchfulness of service users as an opportunity
14 to model a healthy relationship with one’s body (Palmer, 2015; Seah et al., 2018).
15 Meanwhile, certain HCPs with a personal history of ED emphasized the need for self-
16 awareness, including knowing one’s residual symptoms (Williams & Haverkamp, 2015).
17 Supervision, team meetings and personal therapy were presented as fora that could promote a
18 mindful stance.

19 **Connecting with common humanity.**

20 Many HCPs, across 24 studies, described connecting with their own humanity and
21 that of service users in their work, as opposed to an us/them mentality. Connecting with
22 common humanity comprised a recognition of the individuality of clients and HCPs; attuned
23 responses; self-care; considered self-disclosure; a maternal stance, and finding reward in
24 service users’ steps towards recovery. Humanity was evident in attuned responses to service
25 users’ body language, facial expressions, and words: “I’ll just carry on walking beside

WORKING WITH EATING DISORDERS

1 them...that's helped me slow down and have that sort of metaphor of sort of walking
2 alongside at their pace and using their language and just try to get into their life world"
3 (nurse; Snell et al., 2010, p.354). Discussing topics other than their ED afforded a means of
4 acknowledging the individuality of service users, and developing relationships founded on
5 respect.

6 This respect was echoed in clinician accounts of battling alongside, rather than
7 against, service users via externalizing the ED and viewing it as separate from the client.
8 Meanwhile, self-care encompassed respect for one's own humanity, and awareness of the
9 need to preserve wellbeing in order to show humanity to service users. HCPs saw limited
10 self-disclosure as normalizing and facilitating relational safety: "I have talked about things
11 going on in my life....it....puts them on your level" (nurse; Micevski & McCann, 2005,
12 p.107). For some HCPs, predominantly nurses, maternalism – a protective and nurturing
13 stance underpinned by genuine care for service users (Wright, 2015) – facilitated
14 unconditional positive regard and reconciled the need for both empathetic support and
15 boundaries. However, these aspects of maternalism were acknowledged to be "potentially
16 conflictual," and it was suggested that nurses be supported to navigate this through
17 supervision and training (Ryan et al., 2006, p.132).

18 **Humbly open to learning and support.**

19 Awareness of their own fallibility promoted efforts by HCPs in 22 studies to self-
20 improve and engage with service users as equals. Humility was demonstrated by HCPs
21 acknowledging their limitations and valuing support: "I couldn't imagine just doing it on my
22 own...You need support" (healthcare professional; Macdonald et al., 2018, p.229). Humility
23 was also apparent in HCPs demonstrating a desire for more training; providing a rationale for
24 decisions; collaborating with service users and their families, and being open to learning from
25 conflict and mistakes: "[Conflict] gives me the opportunity for self-reflection. I always

WORKING WITH EATING DISORDERS

1 appreciate it when the dietitian that I work with, challenges my thinking” (mental health
2 professional; Dejesse & Zelman, 2013, p.198). Having personal history of ED was portrayed
3 as facilitating humility and guarding against “an us/them kind of perspective” (therapist;
4 Williams & Haverkamp, 2015, p.404).

5 **Line of argument**

6 Synthesizing the studies and reflecting on the resulting concepts generated a line of
7 argument, which is illustrated in Figure 2. The central premise is that the lived experience of
8 working with people with EDs is characterized by feeling drained, demoralized and on edge
9 due to the subversion of HCPs’ usual role as helpers against a backdrop of complexity,
10 chronicity, and treatment refusal. This foregrounds the limits of clinicians’ power to help,
11 given that what they offer may not match the needs and/or wishes of service users. Problems
12 may be beyond their power to resolve due to intrapersonal factors such as their level of
13 experience; interpersonal factors such as relationships within the team; systemic factors such
14 as the time available and cultural context, and/or the client’s readiness to change. For “fixers
15 and doers,” a painful dissonance can result from the sense of powerlessness and subversion of
16 role. One response is to defend against this by avoiding, battling, and blaming service users.
17 While this may provide temporary relief, it can exacerbate the dissonance by violating the
18 principle of unconditional positive regard, and is unlikely to be conducive to productive
19 alliances with service users. Another response is to mindfully face the dissonance with
20 humanity and humility, and seek balance. This involves remaining emotionally present and
21 compassionate while acknowledging that ideal care is illusory, and at times actions that may
22 make HCPs seem, and feel, like prison guards are necessary to preserve service users’ safety.
23 This approach entails tolerance of uncertainty, and of not always being perceived as helpful
24 by service users. However, it also facilitates the connection with clients that was the source

1 of many HCPs' enjoyment of the work, and overcomes the inhumane rigidity of avoidance
2 and blame.

3

4

Figure 2 to go about here

5

6

Discussion

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

This is the first meta-ethnography to focus exclusively on HCPs' lived experiences of working with people with EDs. As well as supporting the findings of previous reviews (Salzmann-Erikson & Dahlén, 2017; Seah et al., 2017; Sibeoni et al., 2017; Thompson-Brenner et al., 2012), it yields new insights regarding the experiences of this work, and approaches to managing challenges in clinical practice. Findings from 37 studies, which spanned 21 years and were of moderate to high quality, were synthesized to produce a key concept of "Coping with caring without curing." The meta-ethnography describes how those working in ED services are limited in their capacity to help, which creates a painful dissonance that can impact on enjoyment of the job, perceptions of service users, and working alliances. This key and novel finding captures the tension inherent in being a professional helper and not knowing how to help, or having your offer of help rejected. Inherent in the key concept is a sense that, compared with many service users, people with EDs are harder to assist because they present with both physical and psychological risks (Seah et al., 2017; Walker & Lloyd, 2011), as well as high levels of complexity and treatment refusal (Franko & Rolfe, 1996; Golan et al., 2009; Kaplan & Garfinkel, 1999).

HCPs reported experiencing work with this client group as emotionally draining, which could lead to negative judgments of service users and themselves. This finding echoes Thompson-Brenner et al.'s (2012) review, with feelings of frustration and helplessness reported as common even among highly experienced therapists in this field (Franko & Rolfe,

WORKING WITH EATING DISORDERS

1 1996). A contributor to the emotional demands was the “culture of surveillance” that some
2 HCPs, particularly those based in inpatient environments, encountered in their work. This
3 sense of being scrutinized by clients, and in some cases colleagues, was a novel finding of
4 this synthesis; it could lead to hostility and anxiety, thereby fueling negative attributions
5 (Cromby & Harper, 2009).

6 The pull towards avoidance, battle and blame identified in this meta-ethnography
7 echoes results from quantitative research showing that many HCPs have stigmatized views of
8 people with EDs (Raveneau et al., 2014). According to the model of helping behavior
9 proposed by Weiner (1986, as cited in Fox, Woodrow, & Leonard, 2012), negative
10 attributions of service users' actions render helpful interventions less likely. Our meta-
11 ethnography showed how such attributions can also lead to moral distress due to the conflict
12 with HCPs' belief in the importance of unconditional positive regard. This is consistent with
13 previous research showing that moral distress is correlated with lack of respect for patients
14 (Lamiani, Borghi, & Argentero, 2017). Individuals with EDs value empathy from and
15 collaboration with HCPs (Bezance & Holliday, 2013); however, as in Sibeoni et al. (2017),
16 the data suggested that some clinicians struggle to maintain empathy and respect in their
17 interactions with service users, and/or seek to avoid them. As well as compromising
18 relationships with clients, engaging in avoidant behaviors is a risk factor for burnout (Fearon
19 & Nicol, 2011).

20 Another contributor to the dissonance and discomfort of the work was feeling unable
21 to help, which is also a risk factor for burnout (Fearon & Nicol, 2011) and is particularly
22 challenging for HCPs with ‘rescue fantasies’ who may identify with the role of ideal carer
23 (Golan et al., 2009). Such identification is arguably understandable in the context of a
24 popular press that tends to depict HCPs as either selfless heroes or heartless villains (Barker,
25 Cornwell, & Gishen, 2016). Moreover, people with EDs commonly seek ideal care (Bell,

WORKING WITH EATING DISORDERS

1 1999; DeLucia-Waack, 1999) and family members are frequently desperate for a “cure” (Fox
2 et al., 2012). The perfectionism inherent in striving to be an ideal carer, and HCPs’
3 avoidance of painful emotions, mirror the experiences of many people with EDs (DeLucia-
4 Waack, 1999). It is possible that this parallel is linked to the high prevalence of personal
5 history of ED among HCPs in the studies.

6 Our meta-ethnography highlighted coping strategies which appear conducive to HCP
7 and service-user wellbeing. The emphasis on connecting with common humanity, and being
8 humbly open to learning and support, are similar to the findings of Salzman-Erikson and
9 Dahlén (2017), and Seah et al. (2017). What is novel is the notion of mindfully drawing on
10 this humanity and humility to adopt a compassionate approach that eschews the rigidity of
11 avoidance or blame. The importance of connecting with common humanity is underlined by
12 research showing that service users appreciate being treated as individuals (Gulliksen et al.,
13 2012). The attunement that characterizes common humanity is valuable given that people
14 with EDs have high levels of insecure attachment (Zachrisson & Skårderud, 2010) and the
15 negative effects of this can be mitigated by healthy, attuned therapeutic relationships
16 (Ardovini, 2002). Mindful awareness, acceptance and self-observation involved HCPs
17 accepting clients and themselves as human, and as and where they are. This aligns with
18 evidence that acceptance-based interventions are helpful when working with challenging
19 populations (Noone & Hastings, 2009).

20 A related healthy practice was taking a balanced and flexible approach, constituting
21 imperfect but responsive care that allows for fallibility and humanity in oneself and others.
22 This stance evokes the position of safe-uncertainty outlined by Mason (1993). According to
23 his model, teams are motivated to remain in the safe-certain position. Correspondingly, the
24 pressure to provide “ideal care” among HCPs working with people with EDs could push the
25 team towards approaches seen as safe and certain (e.g., standardized protocols). However,

WORKING WITH EATING DISORDERS

1 Mason (1993) advocated a position of safe-uncertainty, “which is always in a state of flow”
2 (p.35) and facilitates new ideas. It has been suggested that this position can free HCPs from
3 the bind of needing to know the “final answer,” while supervision and consultation keep them
4 safe in their practice (Fox et al., 2012). Tolerating imperfection is important given that self-
5 expectations of staff have been identified as a critical factor in burnout (Scully, 1983;
6 Freudenberger, 1980, as cited in Kleespies & Dettmer, 2000).

7 The findings of this meta-ethnography are consistent with a multifactorial view of
8 HCPs’ experiences as reflecting an interaction between behaviors of individual clients and
9 colleagues, systemic factors (Franko & Rolfe, 1996), and clinicians’ own practices and
10 internal conflicts (Walker & Lloyd, 2011). Organizational pressures, such as inadequate
11 staffing, contributed to the discrepancy between HCPs’ desire to provide valued care, and
12 feeling they were unable to do so; lack of time is correlated with moral distress among HCPs
13 (Lamiani et al., 2017). It is likely that staff with insufficient resources will seek a safe-certain
14 position, rather than tolerate safe-uncertainty. Furthermore, it has been established that
15 higher caseloads are associated with more negative staff reactions towards people with EDs
16 (Franko & Rolfe, 1996). On a personal level, the capacity of HCPs to be attuned and provide
17 relational safety depends, in part, on their own attachment security (Goodwin, 2003).

18 There were some differences in the data according to experience level, professional
19 background, and setting. In the included studies, less experienced staff were portrayed as
20 more fearful of saying the wrong thing, and helplessness was more prevalent among
21 generalist as opposed to specialist practitioners. Inexperience has been linked to more
22 negative reactions to service users with EDs in previous research (Franko & Rolfe, 1996). It
23 has been suggested that the difficulties generalist staff experience in identifying and
24 responding to EDs reflect the reluctance of clients to report them, and lack of knowledge
25 about treatment options (Waller, Micali, & James, 2014). The systemic pressure of lack of

WORKING WITH EATING DISORDERS

1 time seemed to be felt most keenly in primary care, while the surveillance culture and
2 frustrations around communication were predominantly reported in inpatient settings. The
3 greater likelihood that nurses would engage in avoidance and blame, and adopt a maternal
4 stance, may reflect their high level of contact with people with EDs and their involvement in
5 implementing treatment protocols, positioning them both for intimacy with service users and
6 becoming the object of their anger (Ryan et al., 2006; Zugai et al., 2018^a; Zugai et al., 2018^b).

7 Fox et al. (2012) highlighted the irony that it is generally the least experienced HCPs
8 who spend most time with service users and receive the least containment in the form of
9 space to formulate, process, and reflect. The need for reflective spaces in which to process
10 moral distress in healthcare settings has been highlighted (Kälvemark, Höglund, Hansson,
11 Westerholm, & Arnetz, 2004). It has been suggested that psychologists have an important
12 role in helping ED teams to manage distress and frustration via facilitating reflective spaces;
13 foregrounding the individuality and humanity of clients, and providing training to enhance
14 clinical practice (Fox et al., 2012).

15 Given that the clinician-client relationship is known to affect treatment outcome
16 (Graves et al., 2017) in this field, and low levels of wellbeing among HCPs are associated
17 with poor safety outcomes (Hall, Johnson, Watt, Tsipa, & O'Connor, 2016), staff experiences
18 of the work are a vital consideration for ED services. The findings of this meta-ethnography
19 suggest that HCPs can make a valuable contribution and find reward if they pursue a
20 balanced and humane approach, rather than unrealistically aspiring to be an ideal carer or
21 embracing the punitive role of a prison guard. However, they are likely to need support and
22 reflective space to tolerate the resulting uncertainty and ongoing internal conflict (Halton,
23 2003).

24 **Limitations**

WORKING WITH EATING DISORDERS

1 As almost one-third of the included studies focused exclusively on AN, findings may
2 be most pertinent to those working with this condition (Hage et al., 2017^a; Ryan et al., 2006).
3 Exclusion of unpublished studies may have led to the loss of information (Petticrew et al.,
4 2008).

5 The fact that 11 of the included studies were identified from manual searching of
6 reference lists, or by chance via background reading, raised concerns regarding the breadth of
7 our search. It was identified that nine of these 11 studies would have been located by the
8 original database searches if the ‘professionals’ category of search terms had been applied to
9 study abstracts as well as titles. As a confirmatory check, the search was rerun in the
10 databases with this update. Seven articles that were not identified by the previous search
11 were deemed suitable for full screening; none of these met the inclusion criteria. Difficulty
12 with locating qualitative papers through traditional databases (Booth, 2016), due to issues
13 with MESH terms and filters, has led to a suggestion that other means of accessing them
14 (e.g., hand searching and citation tracking) should be employed (Harris et al., 2018).
15 Reviewers have been warned to extend searches beyond well-known databases to avoid
16 missing journals with relevant information that are not referenced on them (Booth, 2016).

17 While critical appraisal has come to be an expected element of qualitative-evidence
18 synthesis, it is contentious because of the epistemological variety of qualitative research, the
19 diversity of appraisal tools, the variability in ratings within as well as between tools (Carroll
20 & Booth, 2015; Dixon-Woods et al., 2007), and the fact that such tools do not measure
21 conceptual quality (Toye et al., 2014). Applying the CASP checklist for qualitative studies to
22 mixed-methods studies is problematic because the latter should be evaluated as a whole,
23 given that the strengths of one strand can compensate for deficiencies of the other (Heyvaert,
24 Hannes, Maes, & Onghena, 2013).

WORKING WITH EATING DISORDERS

1 The prevalence of convenience and purposive sampling among the included studies
2 may have led to bias within individual papers. With convenience sampling, the researcher
3 may not know how the sample characteristics compare with those of the population of
4 interest, and thus there is a risk of inattention to sampling bias and its potential impact on
5 findings (Etikan, Musa, & Alkassim, 2016). Unlike convenience sampling, purposive
6 sampling entails deliberate selection by the researcher and, while it is regarded as more
7 credible than convenience sampling (Marshall, 1996), it is prone to the bias associated with
8 researcher subjectivity. However, both purposive and convenience sampling are consistent
9 with a qualitative approach, in which the aim is not to generalize findings by ensuring the
10 sample is representative of the population but to produce a new and nuanced understanding of
11 the topic; the objective is depth, rather than breadth (Etikan, Musa, & Alkassim, 2016).
12 Moreover, combining data from a large number of studies ensured that a range of
13 perspectives were incorporated in the final synthesis, lessening the potential impact of
14 sampling bias within individual studies (Bearman & Dawson, 2013).

15 Given that the research team includes HCPs, some of whom have worked extensively
16 with people with EDs, there was a risk we might view the data through the prism of our own
17 experiences. We endeavored to be mindful of this potential bias in team meetings, and alert to
18 differences in the data according to national context and service type. The risk of bias was
19 alleviated by the fact that collectively the research team held experience of both psychology
20 and nursing, working in the public and private sectors, and practicing in the UK and New
21 Zealand.

22 **Clinical implications**

23 Findings from this meta-ethnography can be used to inform clinical practice by
24 supporting HCPs to consider the personal impact of their work with EDs, and by raising
25 awareness of and reflection on their coping strategies. HCPs could draw on the findings to

WORKING WITH EATING DISORDERS

1 help them provide safe and compassionate care, rather than avoid and blame people with
2 EDs. This could lead to improvements in HCP wellbeing, job satisfaction, and treatment
3 outcomes.

4 The findings suggest that providing time and space for HCPs working in the ED field
5 to process the dissonance they experience would promote the wellbeing of clinicians and
6 service users (Fox et al., 2012). This provision could take the form of supervision and/or
7 reflective groups. As well as facilitating safe practice and providing containment,
8 supervision can make HCPs feel valued and listened to (Fearon & Nicol, 2011); thus, it can
9 alleviate damage to their self-concept associated with caring without curing. Supervision and
10 reflective spaces may also support HCPs to adopt positions of safe-uncertainty, and select
11 formulation- and evidence-based interventions rather than act in automatic pilot.

12 In primary care, Balint groups could enable HCPs to discuss stressors in a supportive
13 environment (Rabinowitz, Kushnir, & Ribak, 1996), while in multidisciplinary teams
14 Schwartz rounds could help clinicians to make sense of the emotional challenges of their
15 work (Barker et al., 2016). Given that the surveillance culture and frustrations around
16 communication were predominantly reported in studies with inpatient settings, reflective
17 spaces and case conferences appear to be particularly valuable in these environments (Fox et
18 al., 2012). In light of the fact that many HCPs with personal history of an ED experienced a
19 sense of dissonance in relation to non-disclosure, it would be helpful for managers of ED
20 services to promote a safe environment for sharing, perhaps by role modelling and asking in
21 supervision about personal impacts of the work.

22 Introducing mindfulness practice for HCPs could promote the aware and accepting
23 stance underpinning a balanced approach (Raab, 2014), and team formulation meetings could
24 be used to foreground the individuality of service users and collectively support safe and
25 compassionate care. The findings suggest that reducing systemic pressures (e.g., by

WORKING WITH EATING DISORDERS

1 increasing staffing levels) could decrease demands on HCPs, and mitigate against them
2 feeling emotionally drained and helpless. In primary care, closer links with specialist ED
3 services could make clinicians feel more empowered to help.

4 **Future research**

5 A future review could triangulate the findings of this synthesis with quantitative
6 studies. It would be profitable to examine the impact of HCP experiences of dissonance on
7 client outcomes. Future research could evaluate support mechanisms for HCPs working with
8 people with EDs, for example a pre-post evaluation of introducing a structured reflective
9 space such as a Balint group (Rabinowitz et al., 1996) or a Schwartz round (Barker et al.,
10 2016). Further qualitative research on clinicians' experiences of working with people with
11 binge eating disorder and BN would help to clarify whether the challenges reported in this
12 review extend to all EDs.

13 **Conclusion**

14 This meta-ethnography highlights how HCPs experience working with people with
15 EDs, namely the painful dissonance this can entail and how it can be managed in practice.
16 The synthesis provides an interpretation that can help frontline staff and managers to
17 understand and tackle the barriers to thriving in this work, so they can make a positive
18 difference to service users. A new concept was developed from the reciprocal translation:
19 "Coping with caring without curing." Furthermore, we produced a line-of-argument
20 synthesis, expressed as a new model for understanding HCPs' experiences of working with
21 people who have an ED.

22

23

24

25

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

References

Akgül, S., Pehlivan-türk-Kızıllkan, M., Ores, S., Dermal, O., Düzçeker, Y., & Kanbur, N. (2016). Type of setting for the inpatient adolescent with an eating disorder: Are specialized inpatient clinics a must or will the paediatric ward do?. *Turkish Journal of Pediatrics*, 58(6), 641-649. <https://doi.org/10.24953/turkjped.2016.06.010>

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC.

Ardovini, C. (2002). Attachment theory, metacognitive functions and the therapeutic relationship in eating disorders. *Eating and Weight Disorders – Studies on Anorexia, Bulimia and Obesity*, 7(4), 328-331. <https://doi.org/10.1007/BF03324981>

Barker, R., Cornwell, J., & Gishen, F. (2016). Introducing compassion into the education of health care professionals; Can Schwartz Rounds help?. *Journal of Compassionate Health Care*, 3(1), 3. <https://doi.org/10.1186/s40639-016-0020-0>

Barnett-Page, E., & Thomas, J. (2009). Methods for the synthesis of qualitative research: A critical review. *BMC Medical Research Methodology*, 9(1), 59. <https://doi.org/10.1186/1471-2288-9-59>

Bearman, M., & Dawson, P. (2013). Qualitative synthesis and systematic review in health professions education. *Medical Education*, 47(3), 252-260. <https://doi.org/10.1111/medu.12092>

Beggan, J. K., & DeAngelis, M. (2015). “Oh, My God, I Hate You:” The Felt Experience of Being Othered for Being Thin. *Symbolic Interaction*, 38(3), 371-392. <http://doi.org/10.1002/symb.162>

Bell, L. (1999) The spectrum of psychological problems in people with eating disorders: An analysis of 30 eating-disordered patients treated with cognitive analytic therapy. *Clinical Psychology and Psychotherapy*, 6(1), 29–38.

WORKING WITH EATING DISORDERS

- 1 [http://doi.org/10.1002/\(SICI\)1099-0879\(199902\)6:1%3c29::AID-](http://doi.org/10.1002/(SICI)1099-0879(199902)6:1%3c29::AID-)
- 2 [CPP179%3e3.0.CO;2-U](http://doi.org/10.1002/(SICI)1099-0879(199902)6:1%3c29::AID-CPP179%3e3.0.CO;2-U)
- 3 Bezance, J., & Holliday, J. (2013). Adolescents with anorexia nervosa have their say: a
- 4 review of qualitative studies on treatment and recovery from anorexia nervosa.
- 5 *European Eating Disorders Review*, 21(5), 352-360. <http://doi.org/10.1002/erv.2239>
- 6 Bondas, T., & Hall, E. O. (2007^a). A decade of metasynthesis research in health sciences: A
- 7 meta-method study. *International Journal of Qualitative Studies on Health and Well-*
- 8 *being*, 2(2), 101-113. doi: 10.1080/17482620701251684
- 9 Bondas, T., & Hall, E. O. (2007^b). Challenges in approaching metasynthesis research.
- 10 *Qualitative Health Research*, 17(1), 113-121.
- 11 <http://doi.org/10.1177/1049732306295879>
- 12 Booth, A. (2016). Searching for qualitative research for inclusion in systematic reviews: A
- 13 structured methodological review. *Systematic reviews*, 5(1), 74.
- 14 <http://doi.org/10.1186/s13643-016-0249-x>
- 15 Brinchmann, B. S., Moe, C., Valvik, M. E., Balmbra, S., Lyngmo, S., & Skarbø, T. (2017).
- 16 An Aristotelian view of therapists' practice in multifamily therapy for young adults
- 17 with severe eating disorders. *Nursing Ethics*, XX(X), 1-11.
- 18 <http://doi.org/10.1177/0969733017739780>
- 19 Cameron, P., Willis, K., & Richter, D. (1997). Working with people with eating disorders:
- 20 How do professional attitudes affect approaches to early intervention?. *Australian*
- 21 *Journal of Primary Health*, 3(3), 23-31. <http://doi.org/10.1071/PY97018>
- 22 Carroll, C. & Booth, A. (2015). Quality assessment of qualitative evidence for systematic
- 23 review and synthesis: Is it meaningful, and if so, how should it be performed?
- 24 *Research Synthesis Methods*, 6, 149-154. <http://doi.org/10.1002/jrsm.1128>

WORKING WITH EATING DISORDERS

- 1 Carter, N., Webb, C., Findlay, S., Grant, C., & Van Blyderveen, S. (2012). The integration of
2 a specialized eating disorders nurse on a general inpatient pediatric unit. *Journal of*
3 *Pediatric Nursing: Nursing Care of Children and Families*, 27(5), 549-556.
4 <http://doi.org/10.1016/j.pedn.2012.07.007>
- 5 Critical Appraisal Skills Programme (CASP) (2010). *10 questions to help you make sense of*
6 *qualitative research*. Oxford: CASP.
- 7 Cromby, J., & Harper, D. J. (2009). Paranoia: a social account. *Theory & Psychology*, 19(3),
8 335-361. <http://doi.org/10.1177/0959354309104158>
- 9 Davey, A., Arcelus, J., & Munir, F. (2014). Work demands, social support, and job
10 satisfaction in eating disorder inpatient settings: A qualitative study. *International*
11 *Journal of Mental Health Nursing*, 23(1), 60-68. <http://doi.org/10.1111/inm.12014>
- 12 DeJesse, L. D., & Zelman, D. C. (2013). Promoting optimal collaboration between mental
13 health providers and nutritionists in the treatment of eating disorders. *Eating*
14 *Disorders*, 21(3), 185-205. <http://doi.org/10.1080/10640266.2013.779173>
- 15 DeLucia & Waack, J. L. (1999). Supervision for counselors working with eating disorders
16 groups: Countertransference issues related to body image, food, and weight. *Journal*
17 *of Counseling & Development*, 77(4), 379-388. <http://doi.org/10.1002/j.1556->
18 6676.1999.tb02463.x
- 19 Dillaway, H., Lysack, C., & Luborsky, M. R. (2017). Qualitative approaches to interpreting
20 and reporting data. In R.R. Taylor (Ed.) *Kielhofner's Research in Occupational*
21 *Therapy: Methods of Inquiry for Enhancing Practice* (2nd ed., pp.228-43).
22 Philadelphia, Pennsylvania: F.A. Davis.
- 23 Dixon-Woods, M., Sutton, A., Shaw, R., Miller, T., Smith, J., Young, B., ... & Jones, D.
24 (2007). Appraising qualitative research for inclusion in systematic reviews: A

WORKING WITH EATING DISORDERS

- 1 quantitative and qualitative comparison of three methods. *Journal of Health Services*
2 *Research & Policy*, 12(1), 42-47. <http://doi.org/10.1258/135581907779497486>
- 3 Eddy, K.T., Tabri, N., Thomas, J.J., Murray, H.B., Keshaviah, A. Hastings, E., Edkins, K.,
4 Krishna, M., Herzog, D.B., Keel, P.K., Franko, D.L. (2017). Recovery from anorexia
5 nervosa and bulimia nervosa at 22-year follow-up. *The Journal of Clinical*
6 *Psychiatry*, 78(2), 184-189. <http://doi.org/10.4088/JCP.15m10393>
- 7 Espindola, C.R. & Blay, S.L. (2009). Anorexia nervosa's meaning to patients: A qualitative
8 synthesis. *Psychopathology*, 42, 69-80. <http://doi.org/10.1159/000203339>
- 9 Etikan, I., Musa, S. A., & Alkassim, R. S. (2016). Comparison of convenience sampling and
10 purposive sampling. *American journal of theoretical and applied statistics*, 5(1), 1-4.
11 <http://doi.org/10.11648/j.ajtas.20160501.11>
- 12 Fairburn, C.G. (2008). *Cognitive behavior therapy and eating disorders*. New York: Guilford
13 Press.
- 14 Fassino, S., & Abbate-Daga, G. (2013). Resistance to treatment in eating disorders: a critical
15 challenge. <http://doi.org/10.1186/1471-244X-13-282>
- 16 Fearon, C., & Nicol, M. (2011). Strategies to assist prevention of burnout in nursing staff.
17 *Nursing Standard*, 26(14), 35-39. <http://doi.org/10.7748/ns2011.12.26.14.35.c8859>
- 18 Fox, J. R., Dean, M., & Whittlesea, A. (2015). The Experience of Caring For or Living with
19 an Individual with an Eating Disorder: A Meta Synthesis of Qualitative Studies.
20 *Clinical Psychology & Psychotherapy*, 24(1), 103-125.
21 <http://doi.org/10.1002/cpp.1984>
- 22 Fox, J.R.E., Woodrow, C., & Leonard, K. (2012). Working with anorexia nervosa on an
23 eating disorders inpatient unit: Consideration of the issues. In J.R.E. Fox & K.P. Goss
24 (Eds.), *Eating and its disorders* (pp.344-359). Chichester, West Sussex: John Wiley &
25 Sons Ltd.

WORKING WITH EATING DISORDERS

- 1 Franko, D. L., & Rolfe, S. (1996). Countertransference in the treatment of patients with
2 eating disorders. *Psychiatry*, *59*(1), 108-116.
3 <http://doi.org/10.1080/00332747.1996.11024753>
- 4 Golan, M., Yaroslavski, A., & Stein, D. (2009). Managing eating disorders:
5 Countertransference processes in the therapeutic milieu. *International Journal of*
6 *Child Health and Adolescent Health*, *2*(2), 213-227.
- 7 Goodwin, I. (2003). The relevance of attachment theory to the philosophy, organization, and
8 practice of adult mental health care. *Clinical Psychology Review*, *23*(1), 35-56.
9 [http://doi.org/10.1016/S0272-7358\(02\)00145-9](http://doi.org/10.1016/S0272-7358(02)00145-9)
- 10 Graves, T. A., Tabri, N., Thompson-Brenner, H., Franko, D. L., Eddy, K. T., Bourion-Bedes,
11 S., & Isserlin, L. (2017). A meta-analysis of the relation between therapeutic alliance
12 and treatment outcome in eating disorders. *International Journal of Eating Disorders*,
13 *50*(4), 323-340. <http://doi.org/10.1002/eat.22672>
- 14 Gregertsen, E. C., Mandy, W., & Serpell, L. (2017). The egosyntonic nature of anorexia: An
15 impediment to recovery in anorexia nervosa treatment. *Frontiers in Psychology*, *8*,
16 2273. <http://doi.org/10.3389/fpsyg.2017.02273>
- 17 Gulliksen, K. S., Espeset, E., Nordbø, R. H., Skårderud, F., Geller, J., & Holte, A. (2012).
18 Preferred therapist characteristics in treatment of anorexia nervosa: The patient's
19 perspective. *International Journal of Eating Disorders*, *45*(8), 932-941.
20 <http://doi.org/10.1002/eat.22033>
- 21 Hage, T. W., Rø, Ø., & Moen, A. (2017^a). “Do you see what I mean?” staff collaboration in
22 eating disorder units during mealtimes. *BMC nursing*, *16*(1), 40.
23 <http://doi.org/10.1186/s12912-017-0233-3>

WORKING WITH EATING DISORDERS

- 1 Hage, T. W., Rø, Ø., & Moen, A. (2017^b). To bend or not to bend? Rule adherence among
2 staff at an eating disorder unit. *Eating disorders*, 25(2), 134-150.
3 <http://doi.org/10.1080/10640266.2016.1260361>
- 4 Hall, L. H., Johnson, J., Watt, I., Tsipa, A., & O'Connor, D. B. (2016). Healthcare staff
5 wellbeing, burnout, and patient safety: A systematic review. *PLoS One*, 11(7),
6 e0159015. <http://doi.org/10.1371/journal.pone.0159015>
- 7 Halton, W. (2003). Some unconscious aspects of organizational life: Contributions from
8 psychoanalysis. In A. Obholzer & V.Z. Roberts (Eds.) *The unconscious at work*
9 (pp.31-38). London: Routledge.
- 10 Harken, W., Maxwell, J., Hainline, M., Pollack, L., & Roberts, C. (2017). Perceptions of
11 caring for adolescents with eating disorders hospitalized on a general pediatric unit.
12 *Journal of Pediatric Nursing: Nursing Care of Children and Families*, 34, e34-e41.
13 <http://doi.org/10.1016/j.pedn.2017.02.008>
- 14 Harris, J.L., Booth, A., Cargo, M., Hannes, K., Harden, A., Flemming, K.... & Noyes, J.
15 (2018). Cochrane qualitative and implementation methods group guidance series –
16 Paper 2: Methods for question formulation, searching, and protocol development for
17 qualitative evidence synthesis. *Journal of Clinical Epidemiology*, 97(39-48).
18 <http://doi.org/10.1016/j.jclinepi.2017.10.023>
- 19 Hay, P., Darby, A., & Mond, J. (2007). Knowledge and beliefs about bulimia nervosa and its
20 treatment: A comparative study of three disciplines. *Journal of Clinical Psychology in*
21 *Medical Settings*, 14(1), 59-68. <http://doi.org/10.1007/s10880-007-9057-8>
- 22 Heyvaert, M., Hannes, K., Maes, B., & Onghena, P. (2013). Critical appraisal of mixed
23 methods studies. *Journal of Mixed Methods Research*, 7(4), 302-327.
24 <http://doi.org/10.1177/1558689813479449>

WORKING WITH EATING DISORDERS

- 1 Hunt, D., & Churchill, R. (2013). Diagnosing and managing anorexia nervosa in UK primary
2 care: a focus group study. *Family Practice*, 30(4), 459-465.
3 <http://doi.org/10.1093/fampra/cmt013>
- 4 Jansen, A. (2016). Eating disorders need more experimental psychopathology. *Behaviour*
5 *Research and Therapy*, 86, 2-10. <http://doi.org/10.1016/j.brat.2016.08.004>
- 6 Jarman, M., Smith, J. A., & Walsh, S. (1997). The psychological battle for control: A qualitative
7 study of health-care professionals' understandings of the treatment of anorexia nervosa.
8 *Journal of Community & Applied Social Psychology*, 7(2), 137-152.
9 [http://doi.org/10.1002/\(SICI\)1099-1298\(199704\)7:2<137::AID-CASP404>3.0.CO;2-G](http://doi.org/10.1002/(SICI)1099-1298(199704)7:2<137::AID-CASP404>3.0.CO;2-G)
- 10 Jones, W. R., Saeidi, S., & Morgan, J. F. (2013). Knowledge and attitudes of psychiatrists
11 towards eating disorders. *European Eating Disorders Review*, 21(1), 84-88.
12 <http://doi.org/10.1002/erv.2155>
- 13 Källemark, S., Höglund, A. T., Hansson, M. G., Westerholm, P., & Arnetz, B. (2004). Living
14 with conflicts-ethical dilemmas and moral distress in the health care system. *Social*
15 *Science & Medicine*, 58(6), 1075-1084. [http://doi.org/10.1016/S0277-9536\(03\)00279-](http://doi.org/10.1016/S0277-9536(03)00279-)
16 X
- 17 Kaplan, A. S., & Garfinkel, P. E. (1999). Difficulties in treating patients with eating
18 disorders: A review of patient and clinician variables. *The Canadian Journal of*
19 *Psychiatry*, 44(7), 665-670. <http://doi.org/10.1177/070674379904400703>
- 20 King, S. J., & Turner, D. S. (2000). Caring for adolescent females with anorexia nervosa:
21 Registered nurses' perspective. *Journal of Advanced Nursing*, 32(1), 139-147.
22 <http://doi.org/10.1046/j.1365-2648.2000.01451.x>
- 23 Kleespies, P. M., & Dettmer, E. L. (2000). The stress of patient emergencies for the clinician:
24 Incidence, impact, and means of coping. *Journal of Clinical Psychology*, 56(10),

WORKING WITH EATING DISORDERS

- 1 1353-1369. [http://doi.org/10.1002/1097-4679\(200010\)56:10%3c1353::AID-](http://doi.org/10.1002/1097-4679(200010)56:10%3c1353::AID-)
- 2 [JCLP7%3e3.0.CO;2-3](http://doi.org/10.1002/1097-4679(200010)56:10%3c1353::AID-JCLP7%3e3.0.CO;2-3)
- 3 Lamiani, G., Borghi, L., & Argentero, P. (2017). When healthcare professionals cannot do
- 4 the right thing. A systematic review of moral distress and its correlates. *Journal of*
- 5 *Health Psychology, 22*(1), 51-67. <http://doi.org/10.1177/1359105315595120>
- 6 Land, P. (2004). Thinking about feelings: working with the staff of an eating disorders unit.
- 7 *Psychoanalytic Psychotherapy, 18*(4), 390-403.
- 8 <http://doi.org/10.1080/1474973042000297344>
- 9 Le, L. K. D., Barendregt, J. J., Hay, P., & Mihalopoulos, C. (2017). Prevention of eating
- 10 disorders: A systematic review and meta-analysis. *Clinical Psychology Review, 53*,
- 11 46-58. <http://doi.org/10.1016/j.cpr.2017.02.001>
- 12 Linville, D., Aoyama, T., Knoble, N. B., & Gau, J. (2013). The effectiveness of a brief eating
- 13 disorder training programme in medical settings. *Journal of Research in Nursing,*
- 14 *18*(6), 544-558. <http://doi.org/10.1177/1744987112452182>
- 15 Linville, D., Benton, A., O'Neil, M., & Sturm, K. (2010). Medical providers' screening,
- 16 training and intervention practices for eating disorders. *Eating disorders, 18*(2), 110-
- 17 131. <http://doi.org/10.1080/10640260903585532>
- 18 Long, S., Wallis, D. J., Leung, N., Arcelus, J., & Meyer, C. (2012). Mealtimes on eating
- 19 disorder wards: A two-study investigation. *International Journal of Eating Disorders,*
- 20 *45*(2), 241-246. <http://doi.org/10.1002/eat.20916>
- 21 Lock, J. (2015). An update on evidence-based psychosocial treatments for eating disorders in
- 22 children and adolescents. *Journal of Clinical Child & Adolescent Psychology, 44*(5),
- 23 707-721. <http://doi.org/10.1080/15374416.2014.971458>
- 24 Macdonald, P., Kan, C., Stadler, M., De Bernier, G. L., Hadjimichalis, A., Le Coguic, A. S.
- 25 & Treasure, J. (2018). Eating disorders in people with Type 1 diabetes: experiential

WORKING WITH EATING DISORDERS

- 1 perspectives of both clients and healthcare professionals. *Diabetic Medicine*, 35(2),
2 223-231. <http://doi.org/10.1111/dme.13555>
- 3 Marshall, M. N. (1996). Sampling for qualitative research. *Family practice*, 13(6), 522-526.
4 <https://doi.org/10.1093/fampra/13.6.522>
- 5 Masson, P. C., & Sheeshka, J. D. (2009). Clinicians' perspectives on the premature
6 termination of treatment in patients with eating disorders. *Eating Disorders*, 17(2),
7 109-125. <http://doi.org/10.1080/10640260802714548>
- 8 Micevski, V., & McCann, T. V. (2005). Developing interpersonal relationships with
9 adolescents with anorexia nervosa. *Contemporary Nurse*, 20(1), 102-116.
10 <http://doi.org/10.5172/conu.20.1.102>
- 11 Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G., The PRISMA Group (2009). Preferred
12 reporting items for systematic reviews and meta-analyses: The PRISMA statement.
13 *PLoS Med*, 6(7). <http://doi.org/10.1371/journal.pmed.1000097>
- 14 Noblit, G.W., & Hare, H.D. (1988). *Meta-ethnography: Synthesising qualitative studies*.
15 Thousand Oaks, California: Sage.
- 16 Noone, S. J., & Hastings, R. P. (2009). Building psychological resilience in support staff
17 caring for people with intellectual disabilities: Pilot evaluation of an acceptance-based
18 intervention. *Journal of Intellectual Disabilities*, 13(1), 43-53.
19 <http://doi.org/10.1177/1744629509103519>
- 20 Oyer, L., O'Halloran, M. S., & Christoe-Frazier, L. (2016). Understanding the working
21 alliance with clients diagnosed with anorexia nervosa. *Eating Disorders*, 24(2), 121-
22 137. <http://doi.org/10.1080/10640266.2015.1034050%20%20>
- 23 Palmer, A. E. (2015). The Lived Experience of Dance/Movement Therapists Working with
24 Patients with Eating Disorders. *American Journal of Dance Therapy*, 37(2), 113-134.
25 <http://doi.org/10.1007/s10465-015-9198-5>

WORKING WITH EATING DISORDERS

- 1 Petticrew, M., Egan, M., Thomson, H., Hamilton, V., Kunkler, R., & Roberts, H. (2008).
2 Publication bias in qualitative research: What becomes of qualitative research
3 presented at conferences? *Journal of Epidemiology & Community Health*, 62(6), 552-
4 554. <http://doi.org/10.1136/jech.2006.059394>
- 5 Raab, K. (2014). Mindfulness, self-compassion, and empathy among health care
6 professionals: A review of the literature. *Journal of Health Care Chaplaincy*, 20(3),
7 95-108. <http://doi.org/10.1080/08854726.2014.913876>
- 8 Rabinowitz, S., Kushnir, T., & Ribak, J. (1996). Preventing burnout: increasing professional
9 self-efficacy in primary care nurses in a Balint Group. *Aaohn Journal*, 44(1), 28-32.
10 <http://doi.org/10.1177/216507999604400108>
- 11 Ramjan, L. M. (2004). Nurses and the ‘therapeutic relationship’: Caring for adolescents with
12 anorexia nervosa. *Journal of Advanced Nursing*, 45(5), 495-503.
13 <http://doi.org/10.1046/j.1365-2648.2003.02932.x>
- 14 Ramjan, L. M., & Gill, B. I. (2012). An inpatient program for adolescents with anorexia
15 experienced as a metaphoric prison. *AJN The American Journal of Nursing*, 112(8),
16 24-33. <http://doi.org/10.1097/01.NAJ.0000418094.31317.52>
- 17 Rance, N. M., Moller, N. P., & Douglas, B. A. (2010). Eating disorder counsellors with
18 eating disorder histories: A story of being “normal”. *Eating Disorders*, 18(5), 377-
19 392. <http://doi.org/10.1080/10640266.2010.511901>
- 20 Raveneau, G., Feinstein, R., Rosen, L. M., & Fisher, M. (2014). Attitudes and knowledge
21 levels of nurses and residents caring for adolescents with an eating disorder.
22 *International Journal of Adolescent Medicine and Health*, 26(1), 131-136.
23 <http://doi.org/10.1515/ijamh-2013-0015>

WORKING WITH EATING DISORDERS

- 1 Reid, M., Williams, S., & Burr, J. (2010). Perspectives on eating disorders and service
2 provision: A qualitative study of healthcare professionals. *European Eating Disorders*
3 *Review, 18*(5), 390-398. <http://doi.org/10.1002/erv.976>
- 4 Reid, M., Williams, S., & Hammersley, R. (2009). Managing eating disorder patients in
5 primary care in the UK: A qualitative study. *Eating Disorders, 18*(1), 1-9.
6 <http://doi.org/10.1080/10640260903439441>
- 7 Reilly, E. E., Anderson, L. M., Gorrell, S., Schaumberg, K., & Anderson, D. A. (2017).
8 Expanding exposure-based interventions for eating disorders. *International Journal of*
9 *Eating Disorders, 50*(10), 1137-1141. <http://doi.org/10.1002/eat.22761>
- 10 Ryan, V., Malson, H., Clarke, S., Anderson, G., & Kohn, M. (2006). Discursive constructions
11 of 'eating disorders nursing': An analysis of nurses' accounts of nursing eating
12 disorder patients. *European Eating Disorders Review, 14*(2), 125-135.
13 <http://doi.org/10.1002/erv.666>
- 14 Salzmann-Erikson, M., & Dahlén, J. (2017). Nurses' establishment of health promoting
15 relationships: A descriptive synthesis of anorexia nervosa research. *Journal of Child*
16 *and Family Studies, 26*(1), 1-13. <http://doi.org/10.1007/s10826-016-0534-2>
- 17 Seah, X. Y., Tham, X. C., Kamaruzaman, N. R., & Yobas, P. K. (2017). Knowledge, attitudes
18 and challenges of healthcare professionals managing people with eating disorders: A
19 literature review. *Archives of Psychiatric Nursing, 31*(1), 125-136.
20 <http://doi.org/10.1016/j.apnu.2016.09.002>
- 21 Seah, X. Y., Tham, X. C., Kamaruzaman, N. R., & Yobas, P. K. (2018). Nurses' perception
22 of knowledge, attitudes and reported practice towards patients with eating disorders:
23 A concurrent mixed-methods study. *Archives of Psychiatric Nursing, 32*(1), 133-151.
24 <http://doi.org/10.1016/j.apnu.2017.11.011>

WORKING WITH EATING DISORDERS

- 1 Sibeoni, J., Orri, M., Valentin, M., Podlipski, M. A., Colin, S., Pradere, J., & Revah-Levy, A.
2 (2017). Metasynthesis of the views about treatment of anorexia nervosa in
3 adolescents: Perspectives of adolescents, parents, and professionals. *PloS one*, *12*(1),
4 e0169493. <http://doi.org/10.1371/journal.pone.0169493>
- 5 Snell, L., Crowe, M., & Jordan, J. (2010). Maintaining a therapeutic connection: Nursing in
6 an inpatient eating disorder unit. *Journal of Clinical Nursing*, *19*(3-4), 351-358.
7 <http://doi.org/10.1111/j.1365-2702.2009.03000.x>
- 8 Startup, H., Mountford, V. A., Lavender, A., & Schmidt, U. (2015). Cognitive behavioural
9 case formulation in complex eating disorders. In N. Tarrrier and J. Johnson (Eds.)
10 *Case Formulation in Cognitive Behaviour Therapy: The Treatment of Challenging*
11 *and Complex Cases* (2nd ed., pp.239-264). London: Routledge.
- 12 Tan, J. O., Karim, S. A., Lee, H. Y., Goh, Y. L., & Lee, E. L. (2013). Cultural and ethical
13 issues in the treatment of eating disorders in Singapore. *Asian Bioethics Review*, *5*(1),
14 40-55. <http://doi.org/10.1353/asb.2013.0003>
- 15 Thompson-Brenner, H., Satir, D. A., Franko, D. L., & Herzog, D. B. (2012). Clinician
16 reactions to patients with eating disorders: A review of the literature. *Psychiatric*
17 *Services*, *63*(1), 73-78. <http://doi.org/10.1176/appi.ps.201100050>
- 18 Toye, F., Seers, K., & Barker, K. L. (2017). Meta-ethnography to understand healthcare
19 professionals' experience of treating adults with chronic non-malignant pain. *BMJ*
20 *Open*, *7*(12), e018411. <http://doi.org/10.1136/bmjopen-2017-018411>
- 21 Trammell, E. L., Reed, D., & Boylan, M. (2016). Education and Practice Gaps of Registered
22 Dietitian Nutritionists Working With Clients With Eating Disorders. *Topics in*
23 *Clinical Nutrition*, *31*(1), 73-85. <http://doi.org/10.1097/TIN.0000000000000059>

WORKING WITH EATING DISORDERS

- 1 Walker, S. & Lloyd, C. (2011). Barriers and attitudes health professionals working in eating
2 disorders experience. *International Journal of Therapy & Rehabilitation*, 18(7), 383-
3 391. <http://doi.org/10.12968/ijtr.2011.18.7.383>
- 4 Waller, G., Micali, N., & James, A. (2014). General Practitioners are poor at identifying the
5 eating disorders. *Advances in Eating Disorders: Theory, Research and Practice*, 2(2),
6 146-157. <http://doi.org/10.1080/21662630.2013.859437>
- 7 Walsh, D., & Downe, S. (2005). Meta-synthesis method for qualitative research: A literature
8 review. *Journal of Advanced Nursing*, 50(2), 204-211. [http://doi.org/10.1111/j.1365-
9 2648.2005.03380.x](http://doi.org/10.1111/j.1365-2648.2005.03380.x)
- 10 Walsh, D., & Downe, S. (2006). Appraising the quality of qualitative research. *Midwifery*,
11 22(2), 108-119. <http://doi.org/10.1016/j.midw.2005.05.004>
- 12 Warren, C. S., Crowley, M. E., Olivardia, R., & Schoen, A. (2008). Treating patients with
13 eating disorders: An examination of treatment providers' experiences. *Eating
14 Disorders*, 17(1), 27-45. <http://doi.org/10.1080/10640260802570098>
- 15 Warren, C. S., Schafer, K. J., Crowley, M. E., & Olivardia, R. (2012). A qualitative analysis
16 of job burnout in eating disorder treatment providers. *Eating Disorders*, 20(3), 175-
17 195. <http://doi.org/10.1080/10640266.2012.668476>
- 18 Westmoreland, P., Krantz, M. J., & Mehler, P. S. (2016). Medical complications of anorexia
19 nervosa and bulimia. *The American Journal of Medicine*, 129(1), 30-37.
20 <https://doi.org/10.1016/j.amjmed.2015.06.031>
- 21 Williams, M., & Haverkamp, B. E. (2015). Eating disorder therapists' personal eating
22 disorder history and professional ethics: An interpretive description. *Eating
23 Disorders*, 23(5), 393-410. <http://doi.org/10.1080/10640266.2015.1013393>

WORKING WITH EATING DISORDERS

- 1 Wright, K. M. (2015). Maternalism: a healthy alliance for recovery and transition in eating
2 disorder services. *Journal of Psychiatric and Mental Health Nursing*, 22(6), 431-439.
3 <http://doi.org/10.1111/jpm.12198>
- 4 Wright, K. M., & Hacking, S. (2012). An angel on my shoulder: A study of relationships
5 between women with anorexia and healthcare professionals. *Journal of Psychiatric
6 and Mental Health Nursing*, 19(2), 107-115. [http://doi.org/10.1111/j.1365-
8 2850.2011.01760.x](http://doi.org/10.1111/j.1365-
<u>7</u> 2850.2011.01760.x)
- 8 Zachrisson, H. D., & Skårderud, F. (2010). Feelings of insecurity: Review of attachment and
9 eating disorders. *European Eating Disorders Review*, 18(2), 97-106.
10 <http://doi.org/10.1002/erv.999>
- 11 Zaitsoff, S., Pullmer, R., Cyr, M., & Aime, H. (2015). The role of the therapeutic alliance in
12 eating disorder treatment outcomes: a systematic review. *Eating disorders*, 23(2), 99-
13 114. <http://doi.org/10.1080/10640266.2014.964623>
- 14 Zugai, J. S., Stein-Parbury, J., & Roche, M. (2018^a). The nature of the therapeutic alliance
15 between nurses and consumers with anorexia nervosa in the inpatient setting: A
16 mixed-methods study. *Journal of Clinical Nursing*, 27(1-2), 416-426.
- 17 Zugai, J. S., Stein-Parbury, J., & Roche, M. (2018^b). Therapeutic alliance, anorexia nervosa
18 and the inpatient setting: A mixed methods study. *Journal of Advanced Nursing*,
19 74(2), 443-453.
20
21