

## Direct Imaging of Multiple Planets Orbiting the Star HR 8799

Christian Marois,<sup>1,2,3\*</sup> Bruce Macintosh,<sup>2</sup> Travis Barman,<sup>4</sup> B. Zuckerman,<sup>5</sup> Inseok Song,<sup>6</sup> Jennifer Patience,<sup>7</sup> David Lafrenière,<sup>8</sup> René Doyon<sup>9</sup>

<sup>1</sup>National Research Council Canada Herzberg Institute of Astrophysics, 5071 West Saanich Road, Victoria, BC, V9E 2E7, Canada. <sup>2</sup>Lawrence Livermore National Laboratory, 7000 East Avenue, Livermore, CA 94550, USA. <sup>3</sup>Astronomy Department, University of California, Berkeley, CA 94720, USA. <sup>4</sup>Lowell Observatory, 1400 West Mars Hill Road, Flagstaff, AZ 86001, USA. <sup>5</sup>Physics & Astronomy Department and Center for Astrobiology, University of California, Los Angeles, CA 90095, USA. <sup>6</sup>University of Georgia, Physics and Astronomy, 240 Physics, Athens, GA 30602, USA. <sup>7</sup>University of Exeter, School of Physics, Stocker Road, Exeter, EX4 4QL, UK. <sup>8</sup>Department of Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON, M5S 3H4, Canada. <sup>9</sup>Département de Physique and Observatoire du Mont Mégantic, Université de Montréal, C.P. 6128, Succ. Centre-Ville, Montréal, QC, H3C 3J7, Canada.

\*To whom correspondence should be addressed. E-mail: christian.marois@nrc-cnrc.gc.ca

**Direct imaging of exoplanetary systems is a powerful technique that can reveal Jupiter-like planets in wide orbits, can enable detailed characterization of planetary atmospheres, and is a key step toward imaging Earth-like planets. Imaging detections are challenging due to the combined effect of small angular separation and large luminosity contrast between a planet and its host star. High-contrast observations with the Keck and Gemini telescopes have revealed three planets orbiting the star HR 8799, with projected separations of 24, 38, and 68 astronomical units. Multi-epoch data show counter-clockwise orbital motion for all three imaged planets. The low luminosity of the companions and the estimated age of the system imply planetary masses between 5 and 13 times that of Jupiter. This system resembles a scaled-up version of the outer portion of our solar system.**

During the past decade various planet detection techniques—precision radial velocities, transits, and microlensing—have been used to detect a diverse population of exoplanets. However, these methods have two limitations. First, the existence of a planet is inferred through its influence on the star about which it orbits; the planet is not directly discerned [photometric signals from some of the closest-in giant planets have been detected by careful analysis of the variations in the integrated brightness of the system as the planet orbits its star (*I*)]. Second, these techniques are limited to small (transits) to moderate (precision radial velocity and microlensing) planet-star separation. The effective sensitivities of the latter two techniques diminish rapidly at semi-major axes beyond about 5 AU. Direct observations allow discovery of planets in wider orbits and allow the spectroscopic and photometric characterization of their complex atmospheres to derive their physical characteristics.

There is indirect evidence for planets in orbits beyond 5 AU from their star. Some images of dusty debris disks orbiting main-sequence stars (the Vega phenomenon) show spatial structure on a scale of tens to hundreds of astronomical units (2). The most likely explanation of such structure is gravitational perturbations by planets with semi-major axes comparable to the radius of the dusty disks and rings [see references in (3)].

The only technique currently available to detect planets with semi-major axes greater than about 5 AU in a reasonable amount of time is infrared imaging of young, nearby stars. The detected near-infrared radiation is escaped internal heat energy from the recently formed planets. During the past decade, hundreds of young stars with ages  $\leq 100$  Ma have been identified within  $\sim 100$  pc of Earth (4, 5), and many of these have been imaged in the near-IR with ground-based adaptive optics (AO) systems and with the Hubble Space Telescope. Direct imaging searches for companions of these stars have detected some objects that are generally considered to be near or above the mass threshold  $13.6 M_{\text{Jup}}$  dividing planets from brown dwarfs [see (6) for an example and (7) for a list of known substellar objects orbiting stars], and one planetary mass companion that is orbiting a brown dwarf, not a star (8). Recently, Lafrenière *et al.* (9) have detected a candidate planet near a young (5 Ma) star of the Upper Scorpius association, but a proper motion analysis is required to confirm that it is bound to the host star and not an unrelated low-mass member of the young association. In this issue, Kalas *et al.* report the detection, in visible light, of a candidate planetary mass companion near the inner edge of the Fomalhaut debris disk (10). Non-detections of the candidate companion at near-IR wavelengths suggest that the detected visible flux may be primarily host-star light scattering off circumplanetary dust rather than photons from

the underlying object. A statistical Bayesian analysis of a dedicated AO survey of nearby young F-, G- and K-type stars shows that exoplanets are relatively rare at separations  $>20$  AU around stars with masses similar to the Sun (11).

Bright A-type stars have been mostly neglected in imaging surveys since the higher stellar luminosity offers a less favorable planet-to-star contrast. However, main sequence A-type stars do have some advantages. The higher-mass A stars can retain heavier and more extended disks and thus might form massive planets at wide separations, making their planets easier to detect. Millimeter interferometric continuum observations of the nearest Herbig Ae stars, the precursors to A-type stars, indicate that these are encircled by disks with masses up to several times the Minimum Mass Solar Nebula (12), the minimum amount of solar abundance material ( $0.01M_{\text{Sun}}$ ) required to form all planets in the solar system (13). Associated millimeter line observations resolve these gas disks and indicate that their outer radii are 85–450 AU (12). The most exceptional example of a young A-star disk is the one orbiting IRAS 18059-3211, which is estimated to have a mass of 90 times the Minimum Mass Solar Nebula and an outer radius extending to  $\sim 3,000$  AU (14). Radial velocity surveys of evolved A stars do seem to confirm these hypotheses by showing a trend of a higher frequency of planets at wider separations (15). In this article, we describe the detection of three faint objects at  $0.63''$ ,  $0.95''$ , and  $1.73''$  (24, 38, and 68 AU projected separation) (see Fig. 1) from the dusty and young A-type main sequence star HR 8799, show that all objects are co-moving with HR 8799, and describe their orbital motion and physical characteristics.

**HR 8799 stellar properties.** HR 8799 [also V342 Peg & HIP114189, located 39.4 pc (16) from Earth] is the only star known that has simultaneously been classified as  $\gamma$  Doradus (variable),  $\lambda$  Bootis (metal-poor Population I A-type star) and Vega-like (far-IR excess emission from circumstellar dust) (17, 18). A fit to the Infrared Astronomical Satellite (IRAS) and Infrared Space Observatory (ISO) photometry indicates that it has a dominant dust disk with temperature of 50 K (3, 19). Such blackbody grains, in an optically thin disk, would reside  $\sim 75$  AU ( $\sim 2''$ ) from HR 8799. This would place the dust just outside the orbit of the most distant companion seen in our images (see Fig. 1), similar to the way the Kuiper Belt is confined by Neptune in our solar system.

The fractional IR luminosity ( $L_{\text{IR}}/L_{\text{bol}} = 2.3 \times 10^{-4}$ ) (19, 20) is too bright to come from a geometrically thin, flat disk orbiting at such large distances from HR 8799. Such an optically thin disk would need to be warped or “puffed up” in the vertical direction, plausibly by the gravitational influence of nearby planets. Submillimeter photometry indicates a dust mass of 0.1 Earth masses (21), making it one of the most massive debris disks detected by IRAS (19).

When planets form, gravitational potential energy is released and turned into heat in their interior. As planets do not possess any internal nuclear energy source to maintain their temperature, they cool down and become less luminous with time. For massive planets, this self-luminosity can dominate over their stellar insolation for hundreds of millions or billions of years. With some assumptions on the initial conditions at the time of formation, a planet’s mass can be derived simply by estimating the planet’s luminosity and the system age. Our age estimate for HR8799 is based on four lines of evidence: the star’s galactic space motion, the star’s position in a color-magnitude diagram, the typical age of  $\lambda$  Boo and  $\gamma$  Dor class stars and the large mass of the HR 8799 debris disk.

Most young stars in the solar neighborhood have Galactic space motions (UVW) that fall in limited ranges. HR 8799’s space motion with respect to the Sun, as calculated from published distance, radial velocity and proper motion, is  $UVW = (-11.9, -21.0, -6.8 \text{ km s}^{-1})$  (16, 22). This UVW is similar to that of other stars with an age between that of the TW Hydra association [8 Ma (4, 5)] and the Pleiades [ $125 \pm 8$  Ma (23)]. The UVW of HR 8799 is similar to that of members of the 30 Ma old, southern hemisphere, Columba and Carina Associations (5). Calculations of the UVW of the young stars HD 984 and HD 221318, which lie near HR8799, show that their space motions are similar to that of HR8799. We estimate the ages of HD 984 and HD 221318 to be 30 and 100 Ma, respectively, while the FEPS team estimates the age of HD 984 to be 40 Ma (24). Overall, the UVW of HR 8799 is clearly consistent with those of young clusters and associations in the solar neighborhood. Of course, in this UVW range of young stars, there are also older stars with random motions; so other, independent, methods must also be employed to place limits/constraints on the age of HR 8799.

HR 8799 is also found below the main sequence of the Pleiades,  $\alpha$  Per (70 Ma) and IC2391 (50 Ma) on a Hertzsprung-Russell diagram. This is consistent with a younger age compared to that of the Pleiades (25). Even with the more recent Tycho measurement and correcting for the star’s low metallicity, so that B-V is increased and lies between 0.26 and 0.3, HR 8799 still lies low on the Hertzsprung-Russell diagram when plotted against known young stars (25), consistent with our young age estimate.

The  $\lambda$  Boo stars are generally thought to be young, up to a few 100 Ma (26). The  $\gamma$  Dor class stars are probably also young; they are seen in the Pleiades and in NGC 2516 (age  $\sim 100$  Ma), but not in the Hyades (age  $\sim 650$  Ma) (27). Finally, the probability that a star has a massive debris disk like HR 8799 declines with age (19). Considering all of the above, we arrive at an estimate of 60 Ma and a range between 30 and 160 Ma, consistent with an earlier independent estimate of 20–150 Ma (20). The conservative age upper limit

for HR 8799 is chosen to be the  $\sim 5\sigma$  upper limit to the Pleiades age.

**Observations.** The sensitivity of high-contrast ground-based AO imaging is limited primarily by quasi-static speckle artifacts; at large separations ( $>0.5''$ ), the main source of speckles is surface errors on the telescope primary mirror and internal optics. To remove this noise, we used angular differential imaging (ADI) in our observations (28, 29). This technique uses the intrinsic field-of-view rotation of altitude/azimuth telescopes to decouple exoplanets from optical artifacts. An ADI sequence is obtained by keeping the telescope pupil fixed on the science camera and allowing the field-of-view to slowly rotate with time around the star. Our observations were obtained in the near-infrared (1.1 to 4.2 microns), a regime where the planets are expected to be bright and where the AO system provides excellent image correction. ADI sequences at various wavelengths were acquired using the adaptive optics system at the Keck and Gemini telescopes and the corresponding facility near-infrared cameras, NIRC2 and NIRI, between 2007 and 2008 (see Fig. 1). During each observing sequence, we typically obtained a mix of unsaturated short-exposure images of the star, to determine its precise location and brightness, together with a set of 30-s exposures that overexposed the central star but had maximum sensitivity to faint companions. Some coronagraphic images were also acquired with NIRC2 to benefit from the simultaneous photometric calibration achievable with a partially transmissive focal plane mask. The b and c companions were first seen in October 2007 Gemini data; the d component was first detected in Keck data in 2008. The b and c components were also visible in a re-analysis of non-ADI Keck data obtained for a related program in 2004 [the data sets and the reduction technique are described further in the supplemental online material (SOM)].

**Astrometric analysis.** Following the initial detection of the companions, we evaluated their positions relative to the star to confirm that they are co-moving with it (possibly including orbital motion) and not unrelated back/foreground objects (see SOM; Table 1, table S2, and Fig. 2). Because HR 8799b was visible in the 2004 Keck images, we have more than 4 years of time baseline for proper motion measurements. With the large proper motion of HR 8799 ( $0.13''/\text{year}$ ), the HR 8799b object is shown to be bound at a significance of 98 times the estimated 1-sigma uncertainty. Additionally, the data show that it is orbiting counter-clockwise. It moved  $25 \pm 2$  mas/year ( $0.98$  AU/year) southeast during the 4 year period. Its detected orbital motion is near perpendicular to the line connecting the planet and primary, suggesting that the system is viewed nearly pole-on and that the orbit is not very eccentric. The near face-on perspective is further supported by the slow projected rotational velocity of HR 8799 [ $\sim 40$  km sec $^{-1}$  (17)]; this is

well below average for late-A and early-F type stars (30). If we assume that it has a semi-major axis of 68 AU, a circular orbit, a pole-on view, and a host stellar mass equal to 1.5 solar masses, then the orbital period and motion of HR 8799b are  $\sim 450$  years and  $0.93$  AU/year ( $24$  mas/year) respectively, consistent with our measurements.

HR 8799c is also detected, at lower significance, in the 2004 data set. The measurement of its 4 year proper motion confirms that it is bound to the star at the 90 sigma level. Its orbit is also counter-clockwise at  $30 \pm 2$  mas/year ( $1.18$  AU/year). For its semi-major axis of 38 AU, the orbital period is  $\sim 190$  years and the expected orbital motion is  $1.25$  AU/year ( $32$  mas/year). Again, the orbital motion is close to being perpendicular to the line connecting the planet to the primary.

HR 8799d was first detected in the July 2008 data set. The two months of available proper motion measurements are sufficient to confirm that it is bound to the star at the  $\sim 6$  sigma level. The available data is also consistent with a counter-clockwise orbital motion of  $42 \pm 27$  mas/year ( $1.65$  AU/year). For a semi-major axis of 24 AU, the orbital period is 100 years and the expected orbital motion is  $1.57$  AU/year ( $40$  mas/year).

**HR 8799bcd photometric analysis.** All three companions are intrinsically faint and have red near-IR colors that are comparable to those of substellar-mass objects with low effective temperatures (see Table 1). Compared to old field brown dwarfs (objects with masses between planets and stars), all three companions lie at the base of the L dwarf spectral sequence—objects known to be cool and have dusty clouds in their atmospheres (Fig. 3). Two candidate free-floating Pleiades brown dwarfs, with comparable colors and absolute K-band magnitudes to HR 8799c and d, are consistent with a mass of  $\sim 11 M_{\text{Jup}}$  from evolutionary models (31). If HR 8799 is (as is likely) younger than the Pleiades, the c and d companions would be even less massive. HR 8799b is fainter than all of the known Pleiades substellar members and thus is below  $11 M_{\text{Jup}}$  (Fig. 3). All three companions stand apart from the older, more massive brown dwarfs in a color-magnitude diagram. The known distance to HR 8799, and photometry for each companion that covers a substantial fraction of the spectral energy distribution (SED), allow for a robust measurement of the bolometric luminosity ( $L_{\text{bol}}$ ). We fit a variety of synthetic SEDs (generated with the PHOENIX model atmosphere code) to the observed photometry for each companion assuming their atmospheres were either cloud-free, very cloudy, partly cloudy (50% coverage), or radiated like black bodies. This fitting process is equivalent to simultaneously determining bolometric corrections for each band-pass for various model assumptions. Luminosities were also obtained using the K-band bolometric corrections for brown dwarfs (32). Although



the different models produce different estimates of effective temperature, the range of  $L_{\text{bol}}$  for each object is small (see Table 1), indicating that our estimate is robust against the uncertainty in the details of the atmosphere and clouds (see the SOM for more details).

The cooling of hydrogen-helium brown dwarfs and giant planets is generally well understood; however, the initial conditions associated with the formation of objects from collapsing molecular clouds or core accretion inside a disk are uncertain. Consequently, theoretical cooling tracks of objects at young ages may not be reliable. Recent efforts to establish initial conditions for cooling tracks based on core-accretion models have produced young Jupiter-mass planets substantially fainter ( $<10^{-5} L_{\text{Sun}}$ ) than predicted by traditional models (33). However, these hybrid models do not yet include a realistic treatment of the complex radiative transfer within the accretion-shock and thus provide only lower-limits on the luminosity at young ages. Warmer, more luminous planets originating from core accretion cannot be ruled out.

Although HR 8799 is young, its upper age limit ( $\sim 160$  Ma) is near the time when the differences among cooling tracks with various initial conditions are not so dramatic and, given the uncertainties associated with all planet evolution models, standard cooling tracks are as reliable at these ages as other hybrid models. Figure 4 compares the measured luminosities and age range for HR 8799 bcd to theoretical “hot start” cooling tracks for a variety of masses (34).

The region occupied by all three companions falls below the lowest mass brown dwarf, well inside the planet regime. The masses derived from the luminosities, cooling tracks, and best age for bcd are respectively 7, 10, and 10  $M_{\text{Jup}}$ . See table 1 for values of additional important properties derived from the cooling track comparison, with uncertainties based on our current best age range. In the very unlikely event that the star is older than our estimated upper limit, it would need to be  $>300$  Ma for all three objects to be brown dwarfs.

The large planet masses and orbital radii in the HR 8799 system are challenging to explain in the context of a core accretion scenario. A number of factors such as stellar mass (35), metallicity (36), disk surface density (37), and planet migration in the disk (38) influence the core accretion process. The stellar mass of HR 8799 is larger than the Sun. The star’s metallicity is low, especially in refractory elements, but for a  $\lambda$  Boo star this is usually attributed to the details of the star’s accretion and atmospheric physics rather than an initial low metallicity for the system (26).

The exceptionally dusty debris disk around HR 8799 may indicate that the proto-planetary disk was massive and had a high surface density, factors conducive to planet formation. Alternatively, the giant planets in the HR 8799 system may have formed rapidly from a gravitational instability in the

early disk (39, 40). Some models (40) of such instabilities do favor the creation of massive planets ( $>6 M_{\text{Jup}}$ ).

As suggested by the color-magnitude diagram (Fig. 3), each companion appears to be at the edge of (or inside) the transition region from cloudy to cloud-free atmospheres. Current planet atmosphere models have difficulties fitting the color and spectrum features of these objects. The physical mechanism responsible for the clearing of clouds in ultra-cool atmospheres is not fully understood, but recent cloud models with vertical stratification have had some success at simulating/producing photometric properties in this transition region (41). A modified PHOENIX atmospheric model was developed that incorporates cloud stratification. These updated models were found to match well-known brown dwarfs located in the cloudy/cloud-free transition region. With the cloud stratification model, PHOENIX is capable of producing spectra that are consistent with the observed photometry and the bulk properties (effective temperature, radius, and gravity) predicted by the cooling tracks (Fig. 5). Clearly these synthetic models do not reproduce all of the photometric data, but given the difficulty of cloud modeling, the agreement is sufficient to support the effective temperatures and radii determined from the cooling tracks.

**Conclusions.** The three co-moving companions of HR 8799 are significantly different from known field objects of similar effective temperature; the only similar object known is the planetary mass companion to the brown dwarf 2M1207. Low luminosities of these companions and the young age for HR 8799 indicate that they have planetary masses and are not brown dwarfs. The nature of the system provides an additional indirect line of evidence for planetary-mass companions (and hence a low age). There are no known systems where multiple brown dwarfs independently orbit an early-type star; the only systems we know of with multiple companions in independent orbits are the exoplanetary systems discovered from the precision radial velocity method. Interestingly, our observations show that the HR 8799 planets orbit in the same direction, similar to the planets in our own solar system and consistent with models of planet formation in a disk. In many ways this resembles a scaled-up version of our solar system. HR 8799 has a luminosity of  $4.9 L_{\text{Sun}}$ , so the radius corresponding to a given equilibrium temperature is 2.2 times larger than the corresponding radius in our solar system. Because formation processes will be affected by luminosity - e.g. the location of the snow line where water can condense on rocky material to potentially form giant planet cores - one can view the three planetary companions as having temperature-equivalent projected orbital separations of 11, 17 and 31 AU, to be compared with 9.5, 19, and 30 AU for Saturn, Uranus, and Neptune. The HR 8799 planets are also consistent with formation through instabilities in a massive protoplanetary disk, which may form objects with

masses above 5  $M_{\text{Jup}}$  (40), but the core accretion scenario cannot yet be ruled out.

The presence of these massive planets still leaves dynamic room for other Jovian-mass planets or even lower mass terrestrial planets in the inner part of the system. In our survey, we only observed a few early-type stars before making this detection, compared to similar imaging surveys of young G-, K- and M-type stars that have covered more than a few hundred targets. This may indicate that Jovian-mass planetary companions to early-type stars are much more common at separations beyond  $\sim 20$  AU, consistent with what was suggested by radial velocity surveys of evolved A-type stars (15).

## References and Notes

1. D. Deming, *Proceedings of IAU Symposium 253*, in press; arXiv:0808.1289 (astro-ph).
2. P. Kalas, J. R. Graham, M. Clampin, *Nature* **435**, 1067 (2005).
3. B. Zuckerman, I. Song, *Astrophys J.* **603**, 738 (2004).
4. B. Zuckerman, I. Song, *Annu. Rev. Astron. Astrophys* **42**, 685 (2004).
5. C. A. O. Torres, G. R. Quast, C. H. F. Melo, M. E. Sterzik, *Handbook of Star Forming Regions Vol. II. Astronomical Society of the Pacific*, in press; arXiv:0808.3362 (astro-ph).
6. T. Nakajima et al., *Nature* **378**, 463 (1995).
7. B. Zuckerman, I. Song, *Astron. Astrophys*, in press; arXiv:0811.0429 (astro-ph).
8. G. Chauvin et al., *Astron. Astrophys* **425**, 29 (2004).
9. D. Lafrenière et al., *Astrophys J.*, in press (arXiv:0809.1424v2 (astro-ph)).
10. P. Kalas et al., *Science* (2008); published online 13 November 2008; 10.1126/science.1166609.
11. D. Lafrenière et al., *Astrophys J.* **660**, 770 (2007).
12. V. Mannings, A. I. Sargent, *Astrophys J.* **490**, 792 (1997).
13. S. J. Weidenschilling, *Astrophys. Space Sci* **51**, 153 (1977).
14. V. Bujarrabal, K. Young, D. Fong, *Astron. Astrophys* **483**, 839 (2008).
15. J. Johnson et al., *Astrophys J.* **690**, 833 (2007).
16. E. van Leeuwen, *Astron. Astrophys* **474**, 653 (2007).
17. R. O. Gray, A. B. Kaye, *Astron. J.* **118**, 2993 (1999).
18. K. Sadakane, M. Nishida, *Publ. Astron. Soc. Pac* **98**, 685 (1986).
19. J. H. Rhee, I. Song, B. Zuckerman, M. McElwain, *Astrophys J.* **660**, 1556 (2007).
20. A. Moór et al., *Astrophys J.* **644**, 525 (2006).
21. J. P. Williams, S. M., *Andrews Astrophys J.* **653**, 1480 (2006).
22. R. E. Wilson, *General Catalogue of Stellar Radial Velocities*, Carnegie Inst. Washington D.C. Publ. 601 (1953).
23. J. R. Stauffer, G. Schultz, J. D. Kirkpatrick, *Astrophys J.* **499**, 199 (1998).
24. J. Najita, J. P. Williams, *Astrophys J.* **635**, 625 (2005).
25. B. Zuckerman, *Annu. Rev. Astron. Astrophys* **39**, 549 (2001).
26. R. O. Gray, C. J. Corbally, *Astron. J.* **124**, 989 (2002).
27. K. Krisciunas, R. A. Crowe, K. D. Luedeke, M. Roberts, *Mon. Not. R. Astron. Soc* **277**, 1404 (1995).
28. C. Marois et al., *Astrophys J.* **641**, 556 (2006).
29. D. Lafrenière et al., *Astrophys J.* **660**, 770 (2007).
30. F. Royer, J. Zorec, A. E. Gómez, *Astron. Astrophys* **463**, 671 (2007).
31. S. L. Casewell et al., *Mon. Not. R. Astron. Soc* **378**, 1131 (2007).
32. D. A. Golimowski et al., *Astrophys J.* **127**, 3516 (2004).
33. M. S. Marley et al., *Astrophys J.* **655**, 541 (2007).
34. I. Baraffe et al., *Astron. Astrophys* **402**, 701 (2003).
35. S. Ida, D. N. C. Lin, *Astrophys J.* **626**, 1045 (2005).
36. S. Ida, D. N. C. Lin, *Astrophys J.* **616**, 567 (2004).
37. J. J. Lissauer, *Icarus* **69**, 249 (1987).
38. W. K. M. Rice, P. J. Armitage, *Astrophys J.* **598**, 55 (2003).
39. A. P. Boss, *Science* **276**, 1836 (1997).
40. R. R. Rafikov, *Astrophys J.* **621**, 69 (2005).
41. D. Saumon, M. S., Marley, *Astrophys J.*, arXiv:0808.2611 (2008).
42. C., Ducourant et al., *Astron. Astrophys* **477**, 1 (2008).
43. We thank the Keck and Gemini staff, particularly T. Armandroff, B. Goodrich, and J.-R. Roy for support with the follow-up observations. We thank the UCLA galactic center team, especially J. Lu, for the NIRC2 plate scale and North orientation errors. We are indebted to E. Becklin and R. Racine for their contributions in the earliest stages of this research. C.M. and D.L. are supported in part through postdoctoral fellowships from the Fonds Québécois de la Recherche sur la Nature et les Technologies. Portions of this research were performed under the auspices of the U.S. Department of Energy by LLNL under contract DE-AC52-07NA27344, and also supported in part by the NSF Science and Technology CFAO, managed by the UC Santa Cruz under cooperative agreement AST 98-76783. We acknowledge support by NASA grants to UCLA and Lowell Observatory. R.D. is supported through a grant from the Natural Sciences and Engineering Research Council of Canada. The data were obtained at the W.M. Keck and Gemini Observatories. This publication makes use of data products from the Two Micron All Sky Survey and the SIMBAD database.

## Supporting Online Material

[www.sciencemag.org/cgi/content/full/1166585/DC1](http://www.sciencemag.org/cgi/content/full/1166585/DC1)  
Materials and Methods

30 September 2008; accepted 5 November 2008

Published online 13 November 2008;

10.1126/science.1166585

Include this information when citing this paper.

**Fig. 1.** HR 8799bcd discovery images after the light from the bright host star has been removed by ADI processing. (**Upper left**) A Keck image acquired in July 2004. (**Upper right**) Gemini discovery ADI image acquired in October 2007. Both b and c are detected at the 2 epochs. (**Bottom**) one of our most sensitive ADI image sets acquired at the Keck telescope in September 2008 with all three companions detected. The central region of each image is masked-out due to saturation or to occultation of the bright central star.

**Fig. 2.** HR 8799bcd astrometric analysis. The positions of HR 8799bcd at each epoch are shown in both the overall field of view and in the zoomed-in insets. The solid oscillating line originating from the first detected epoch of each planet is the expected motion of a unbound background objects relative to the star over a duration equal to the maximum interval over which the companions were detected (4 years for b and c, two months for d.) All three companions are confirmed as co-moving with HR 8799 to  $98\sigma$  for b,  $90\sigma$  for c and  $\sim 6\sigma$  for d. Counter-clockwise orbital motion is observed for all three companions. The dashed lines in the small insets connect the position of the planet at each epoch with the star. A schematic dust disk—at 87 AU separation to be in 3:2 resonance with d while also entirely consistent with the far-infrared dust spectrum—is also shown. The inner gray ellipses are the outer Jovian-mass planets of our solar system (Jupiter, Saturn, Uranus & Neptune) and Pluto shown to scale.

**Fig. 3.** Absolute magnitude in H-band versus H-K color. Old, field (grey dots) and young Pleiades brown dwarfs (plusses) are shown along with 2 very low-mass brown dwarfs/planetary mass companions (filled black symbols). Open symbols are HR 8799b (square), c (diamond), and d (circle).

**Fig. 4.** Luminosity versus time for a variety of masses (34). The three coeval points are HR 8799b (square), c (diamond), and d (circle); c and d data points are displaced horizontally for clarity. The locations of the low mass object AB Pic b on the planet/brown dwarf dividing line and a planetary mass companion (2M1207b) to the brown dwarf 2M1207 are also shown [note that alternative models proposed for 2M1207 lead to somewhat larger luminosity and mass ( $\sim 8 M_{\text{Jup}}$ ) for the companion (42)]. The deuterium burning mass limit, currently believed to be  $\sim 13.6 M_{\text{Jup}}$ , has been incorporated into a “working definition” of a planet by the International

Astronomical Union and is used here to separate planets (which also must orbit a star) from brown dwarfs. The boundary between stars and brown dwarfs is set by stable hydrogen burning.


**Fig. 5.** Synthetic spectra from model atmospheres containing clouds located between 10 and 0.1 bar of pressure are compared to the measured fluxes (with 3 sigma error bars) for HR 8799 b, c and d. Response curves for each filter band pass are indicated along the x axis. The predicted magnitudes from the synthetic spectra, averaged over the filter passbands, are shown by the filled symbols.

**Table 1: HR 8799 planetary system data**

| HR 8799                                     |  |                   |                   |
|---|--|-------------------|-------------------|
| Spectral type                               | A5V  |                   |                   |
| Mass  | $1.5 \pm 0.3 M_{\text{Sun}}$ (17)                                    |                   |                   |
| Luminosity                                  | $4.92 \pm 0.41 L_{\text{Sun}}$ (17)                                  |                   |                   |
| Distance                                    | $39.4 \pm 1.0 \text{ pc}$ ( $128 \pm 3 \text{ ly}$ ) (16)            |                   |                   |
| Proper motion [E,N]                         | $[107.88 \pm 0.76, -50.08 \pm 0.63] \text{ mas/year}$ (16)           |                   |                   |
| Age   | 60[30 – 160] Ma  |                   |                   |
| Metallicity                                 | $\log([M/H]/[M/H]_{\text{Sun}}) = -0.47$ (17)                        |                   |                   |
| J, H, Ks, L'                                | $5.383 \pm 0.027, 5.280 \pm 0.018, 5.240 \pm 0.018, 5.220 \pm 0.018$ |                   |                   |
| Separation w.r.t the host star in [E, N]''  |  |                   |                   |
|   | HR 8799b   | c                 | d                 |
| 2004 July 14 ( $\pm 0.005''$ )              | [1.471, 0.884]   | [-0.739, 0.612]   | –                 |
| 2007 Oct. 25 ( $\pm 0.005''$ )              | [1.512, 0.805]   | [-0.674, 0.681]   | –                 |
| 2008 July 11 ( $\pm 0.004''$ )              | [1.527, 0.799]   | [-0.658, 0.701]   | [-0.208, -0.582]  |
| 2008 Aug. 12 ( $\pm 0.002''$ )              | [1.527, 0.801]   | [-0.657, 0.706]   | [-0.216, -0.582]  |
| 2008 Sept. 18 ( $\pm 0.003''$ )             | [1.528, 0.798]   | [-0.657, 0.706]   | [-0.216, -0.582]  |
| Projected Sep. (AU)                         | 68   | 38                | 24                |
| Orbital Motion (''/year)                    | $0.025 \pm 0.002$  | $0.030 \pm 0.002$ | $0.042 \pm 0.027$ |
| Period for pole view<br>cir. orbits (years) | $\sim 460$   | $\sim 190$        | $\sim 100$        |
| $M_{\text{J}}$ ( $1.248\mu\text{m}$ )       | $16.30 \pm 0.16$   | $14.65 \pm 0.17$  | $15.26 \pm 0.43$  |
| $M_{\text{H}}$ ( $1.633\mu\text{m}$ )       | $14.87 \pm 0.17$   | $13.93 \pm 0.17$  | $13.86 \pm 0.22$  |
| $M_{\text{CH4S}}$ ( $1.592\mu\text{m}$ )    | $15.18 \pm 0.17$   | $14.25 \pm 0.19$  | $14.03 \pm 0.30$  |
| $M_{\text{CH4L}}$ ( $1.681\mu\text{m}$ )    | $14.89 \pm 0.18$   | $13.90 \pm 0.19$  | $14.57 \pm 0.23$  |
| $M_{\text{Ks}}$ ( $2.146\mu\text{m}$ )      | $14.05 \pm 0.08$   | $13.13 \pm 0.08$  | $13.11 \pm 0.12$  |
| $M_{\text{L'}}$ ( $3.776\mu\text{m}$ )      | $12.66 \pm 0.11$   | $11.74 \pm 0.09$  | $11.56 \pm 0.16$  |
| Luminosity ( $L_{\text{Sun}}$ )             | $-5.1 \pm 0.1$   | $-4.7 \pm 0.1$    | $-4.7 \pm 0.1$    |
| $T_{\text{eff}}$ (K)                        | 870[800 – 900]   | 1090[1000 – 1100] | 1090[1000 – 1100] |
| Radius ( $R_{\text{Jup}}$ )                 | 1.2[1.1 – 1.3]   | 1.2[1.2 – 1.3]    | 1.2[1.2 – 1.3]    |
| Mass ( $M_{\text{Jup}}$ )                   | 7[5 – 11]  | 10[7 – 13]        | 10[7 – 13]        |


Keck H-band July 14, 2004UT

b



Gemini CH4S Oct. 17, 2007UT

b



Keck K-band Sept. 18, 2008UT

b

