



KEY UPDATES

- Countries and territories reporting mosquito-borne Zika virus infections for the first time in the past week:
 - None
- Countries and territories reporting microcephaly and other central nervous system (CNS) malformations potentially associated with Zika virus infection for the first time in the past week:
 - None
- Countries and territories reporting Guillain-Barré syndrome (GBS) cases associated with Zika virus infection for the first time in the past week:
 - None

ANALYSIS

- Overall, the global risk assessment has not changed.
- The risk of Zika-associated complications in countries thought to be endemic for Zika virus remains largely unknown, due both to uncertainty about the risk associated with different Zika strains, and due to uncertainty about population immunity in these endemic settings. To address the latter question, seroprevalence studies are urgently needed.

SITUATION

- 73 countries and territories (Fig. 1, Table 1) have reported evidence of mosquito-borne Zika virus transmission since 2007 (67 with reports from 2015 onwards), of which:
 - 56 with a reported outbreak from 2015 onwards (Fig. 2, Table 1).
 - Seven with having possible endemic transmission or evidence of local mosquito-borne Zika infections in 2016.
 - 10 with evidence of local mosquito-borne Zika infections in or before 2015, but without documentation of cases in 2016, or with the outbreak terminated.
- Since February 2016, 12 countries have reported evidence of person-to-person transmission of Zika virus (Table 2).

- 22 countries or territories have reported microcephaly and other CNS malformations potentially associated with Zika virus infection or suggestive of congenital infection (Table 3).
- 19 countries and territories have reported an increased incidence of GBS and/or laboratory confirmation of a Zika virus infection among GBS cases (Table 4).
- In Guinea-Bissau, the investigation of five reported cases of microcephaly is ongoing.
- Operational updates:
 - In Paraguay, Mosquito Awareness Week activities are being planned by the Ministry of Health. In addition, there are ongoing studies in hospitals to assess baseline GBS and microcephaly incidence.
 - A training workshop is scheduled for October in Pakistan on prevention, surveillance and control of *Aedes* mosquito vectors in the context of emerging Zika and other *Aedes*-borne diseases in Lahore. A consultative workshop is planned for November to define an appropriate surveillance strategy for the detection of clusters of Zika virus and other arboviral diseases using both syndromic and event-based surveillance approaches.
 - In Jordan, Institute Pasteur Dakar has scheduled a Zika laboratory testing training for November.

Table 1. Countries and territories reporting mosquito-borne Zika virus transmission

Classification	WHO Regional Office	Country / territory	Total
Category 1: Countries with a reported outbreak from 2015 onwards [#]	AFRO	Cabo Verde; Guinea-Bissau	2
	AMRO/PAHO	Anguilla; Antigua and Barbuda; Argentina; Aruba; Bahamas; Barbados; Belize; Bolivia (Plurinational State of); Bonaire, Sint Eustatius and Saba – Netherlands; Brazil; British Virgin Islands; Cayman Islands; Colombia; Costa Rica; Cuba; Curaçao; Dominica; Dominican Republic; Ecuador; El Salvador; French Guiana; Grenada; Guadeloupe; Guatemala; Guyana; Haiti; Honduras; Jamaica; Martinique; Mexico; Nicaragua; Panama; Paraguay; Peru; Puerto Rico; Saint Barthélemy; Saint Kitts and Nevis; Saint Lucia; Saint Martin; Saint Vincent and the Grenadines; Sint Maarten; Suriname; Trinidad and Tobago; Turks and Caicos; United States of America; United States Virgin Islands; Venezuela (Bolivarian Republic of)	47
	WPRO	American Samoa; Fiji; Marshall Islands; Micronesia (Federated States of); Samoa; Singapore; Tonga	7
Subtotal			56
Category 2: Countries with possible endemic transmission or evidence of local mosquito-borne Zika infections in 2016	SEARO	Indonesia; Maldives; Thailand	3
	WPRO	Malaysia; New Caledonia; Philippines; Viet Nam	4
Subtotal			7
Category 3: Countries with evidence of local mosquito-borne Zika infections in or before 2015, but without documentation of cases in 2016, or outbreak terminated	AFRO	Gabon**	1
	PAHO/AMRO	ISLA DE PASCUA – Chile**	1
	SEARO	Bangladesh**	1
	WPRO	Cambodia**; Cook Islands**; French Polynesia**; Lao People's Democratic Republic; Papua New Guinea; Solomon Islands; Vanuatu	7
Subtotal			10
Total			73

[#]The wording has been revised in recognition of the fact that a country that has had a first outbreak since 2015 and in which that outbreak has since terminated, may again report a new outbreak or cases which would qualify the country to be re-included in category 1.

**These countries and territories have not reported Zika virus cases in 2015 or 2016.

Category 1: Countries with a reported outbreak from 2015 onwards[#]

- A laboratory confirmed, autochthonous, mosquito-borne case of Zika virus infection in an area where there is no evidence of circulation of the virus in the past (prior 2015), whether it is detected and reported by the country itself or by another state party diagnosing returning travellers **OR**
- A laboratory confirmed, autochthonous, mosquito-borne case of Zika virus infection in an area where transmission has been previously interrupted. The assumption is that the size of the susceptible population has built up to a sufficient level to allow transmission again; the size of the outbreak will be a function of the size of the susceptible population **OR**
- An increase of the incidence of laboratory confirmed, autochthonous, mosquito-borne Zika virus infection in areas where there is on-going transmission, above two standard deviations of the baseline rate, or doubling the number of cases over a 4-week period. Clusters of febrile illnesses, in particular when epidemiologically-linked to a confirmed case, should be microbiologically investigated.

Category 2: Countries with possible endemic transmission or evidence of local mosquito-borne Zika infections in 2016 with the reporting period beginning in 2007

- Countries or territories that have reported an outbreak with consistent presence of laboratory confirmed, autochthonous, mosquito-borne cases of Zika virus infection 12 months after the outbreak **OR**
- Countries or territories where Zika virus has been circulating for several years with consistent presence of laboratory confirmed, autochthonous, mosquito-borne cases of Zika virus infection or evidence of local mosquito-borne Zika infections in 2016. Reports can be from the country or territory where infection occurred, or from a third party where the case is first recorded according to the International Health Regulations (IHR 2005). Countries with evidence of infection prior to 2007 are listed in <http://www.who.int/bulletin/volumes/94/9/16-171082.pdf>

Category 3: Countries with evidence of local mosquito-borne Zika infections in or before 2015, but without documentation of cases in 2016, or outbreak terminated with the reporting period beginning in 2007

- Absence of confirmed cases over a 3-month period in a specific geographical area with climatic conditions suitable for year-round arbovirus transmission, or over a 12-month period in an area with seasonal vector activity.

Figure 1. Cumulative number of countries and territories by WHO region¹ reporting mosquito-borne Zika virus transmission for the first time in years (2007–2014), and monthly from 1 January 2015 to 12 October 2016




Table 2. Countries reporting non mosquito-borne Zika virus transmission since February 2016

Classification	WHO Regional Office	Country / territory	Total
Countries with evidence of person-to-person transmission of Zika virus, other than mosquito-borne transmission	AMRO/PAHO	Argentina, Canada, Chile, Peru, United States of America	5
	EURO	France, Germany, Italy, Netherlands, Portugal, Spain	6
	WPRO	New Zealand	1
Total			12

¹ <http://www.who.int/about/regions/en/>

Table 3. Countries and territories reporting microcephaly and/or CNS malformation cases potentially associated with Zika virus infection

Reporting country or territory	Number of microcephaly and/or CNS malformation cases suggestive of congenital Zika infections or potentially associated with a Zika virus infection	Probable location of infection
Brazil	2001 ²	Brazil
Cabo Verde	9	Cabo Verde
Canada	1	Undetermined
Costa Rica	1	Costa Rica
Colombia	42 ³	Colombia
Dominican Republic	10 ⁴	Dominican Republic
El Salvador	4	El Salvador
French Guiana	10 ⁵	French Guiana
French Polynesia	8	French Polynesia
Guatemala	17 ⁶	Guatemala
Haiti	1	Haiti
Honduras	1	Honduras
Marshall Islands	1	Marshall Islands
Martinique	12 ⁶	Martinique
Panama	5	Panama
Paraguay	2 ⁷	Paraguay
Puerto Rico	2 ⁸	Puerto Rico
Slovenia	1 ⁹	Brazil
Spain	2	Colombia, Venezuela (Bolivarian Republic of)
Suriname	1	Suriname
Thailand	2	Thailand
United States of America	27 ¹⁰	Undetermined*

*The probable locations of three of the infections were Brazil (1 case), Haiti (1 case) and Mexico, Belize or Guatemala (1 case).

Table 4. Countries and territories reporting Guillain-Barré syndrome (GBS) potentially associated with Zika virus infection

Classification	Country / territory
Reported increase in incidence of GBS cases, with at least one GBS case with confirmed Zika virus infection	Brazil, Colombia, Dominican Republic, El Salvador*, French Guiana, French Polynesia, Guadeloupe ¹¹ , Honduras, Jamaica, Martinique, Suriname**, Venezuela (Bolivarian Republic of)
No increase in GBS incidence reported, but at least one GBS case with confirmed Zika virus infection	Costa Rica, Grenada ¹² , Guatemala, Haiti, Mexico, Panama, Puerto Rico

*GBS cases with previous history of Zika virus infection were reported by the International Health Regulations (2005) National Focal Point in the United States of America.

**One case living in continental Netherlands was diagnosed in mid-January 2016 and reported by the Netherlands.

² Data provided by the Ministry of Health of Brazil on 7 October 2016

³ <http://www.ins.gov.co/boletin-epidemiologico/Boletn%20Epidemiolgico/2016%20Boletin%20epidemiologico%20semana%2039.pdf>

⁴ http://digeprisalud.gob.do/documentos/?drawer=Boletines%20epidemiol%C3%B3gicos*Boletines%20semanales*2016

⁵ <http://invs.santepubliquefrance.fr/fr/Publications-et-outils/Points-epidemiologiques/Tous-les-numeros/Antilles-Guyane/2016/Situation-epidemiologique-du-virus-Zika-aux-Antilles-Guyane.-Point-au-6-octobre-2016>

⁶ <http://www.mspas.gob.gt/index.php/en/mspas/noticias/1239-comunicado-ante-la-epidemia-del-virus-zika.html?tmpl=component&print=1&layout=default&page=>

⁷ <http://www.mspbs.gov.py/v3/paraguay-reporta-sus-dos-primeros-casos-de-microcefalia-asociados-al-zika/>

⁸ <http://www.salud.gov.pr/Estadisticas-Registros-y-Publicaciones/Informes%20Arbovirales/Reporte%20ArboV%20semana%2038-2016.pdf>

⁹ <http://www.neim.org/doi/pdf/10.1056/NEJMoa1600651>

¹⁰ <http://www.cdc.gov/zika/geo/pregnancy-outcomes.html>

¹¹ <http://invs.santepubliquefrance.fr/fr/Publications-et-outils/Points-epidemiologiques/Tous-les-numeros/Antilles-Guyane/2016/Situation-epidemiologique-du-virus-Zika-aux-Antilles-Guyane.-Point-au-15-septembre-2016>

¹² http://health.gov.gd/index.php?option=com_content&view=article&id=434:nine-confirmed-zika-cases-in-grenada&catid=83:latest-news&Itemid=932&lang=en