



$I(J^P) = 0(\frac{1}{2}^+)$ Status: ****

The parity of the Λ_c^+ is defined to be positive (as are the parities of the proton, neutron, and Λ). The quark content is udc . The spin J has not actually been measured. Results of an analysis of $pK^-\pi^+$ decays (JEZABEK 92) are consistent with the expected $J = 1/2$.

We have omitted some results that have been superseded by later experiments. The omitted results may be found in earlier editions.

Λ_c^+ MASS

Our value in 2004, 2284.9 ± 0.6 MeV, was the average of the measurements now filed below as "not used." The BABAR measurement is so much better that we use it alone. Note that it is about 2.6 (old) standard deviations above the 2004 value.

The fit also includes $\Sigma_c - \Lambda_c^+$ and $\Lambda_c^{*+} - \Lambda_c^+$ mass-difference measurements, but this doesn't affect the Λ_c^+ mass. The new (in 2006) Λ_c^+ mass simply pushes all those other masses higher.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2286.46 ± 0.14 OUR FIT				
2286.46 ± 0.14	4891	¹ AUBERT,B	05S BABR	$\Lambda K_S^0 K^+$ and $\Sigma^0 K_S^0 K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2284.7 ± 0.6 ± 0.7	1134	AVERY	91 CLEO	Six modes
2281.7 ± 2.7 ± 2.6	29	ALVAREZ	90B NA14	$pK^-\pi^+$
2285.8 ± 0.6 ± 1.2	101	BARLAG	89 NA32	$pK^-\pi^+$
2284.7 ± 2.3 ± 0.5	5	AGUILAR-...	88B LEBC	$pK^-\pi^+$
2283.1 ± 1.7 ± 2.0	628	ALBRECHT	88C ARG	$pK^-\pi^+, p\bar{K}^0, \Lambda 3\pi$
2286.2 ± 1.7 ± 0.7	97	ANJOS	88B E691	$pK^-\pi^+$
2281 ± 3	2	JONES	87 HBC	$pK^-\pi^+$
2283 ± 3	3	BOSETTI	82 HBC	$pK^-\pi^+$
2290 ± 3	1	CALICCHIO	80 HYBR	$pK^-\pi^+$


¹AUBERT,B 05S uses low-Q $\Lambda K_S^0 K^+$ and $\Sigma^0 K_S^0 K^+$ decays to minimize systematic errors. The error above includes systematic as well as statistical errors. Many cross checks and adjustments to properties of the BABAR detector, as well as the large number of clean events, make this by far the best measurement of the Λ_c^+ mass.

Λ_c^+ MEAN LIFE

Measurements with an error $\geq 100 \times 10^{-15}$ s or with fewer than 20 events have been omitted from the Listings.

VALUE (10^{-15} s)	EVTS	DOCUMENT ID	TECN	COMMENT
200 ± 6 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.				
204.6 $\pm 3.4 \pm 2.5$	8034	LINK	02C FOCS	$pK^-\pi^+$
198.1 $\pm 7.0 \pm 5.6$	1630	KUSHNIR...	01 SELX	$\Lambda_c^+ \rightarrow pK^-\pi^+$

$179.6 \pm 6.9 \pm 4.4$	4749	MAHMOOD	01	CLE2	$e^+ e^- \approx \gamma(4S)$
$215 \pm 16 \pm 8$	1340	FRABETTI	93D	E687	$\gamma Be, \Lambda_c^+ \rightarrow p K^- \pi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$180 \pm 30 \pm 30$	29	ALVAREZ	90	NA14	$\gamma, \Lambda_c^+ \rightarrow p K^- \pi^+$
$200 \pm 30 \pm 30$	90	FRABETTI	90	E687	$\gamma Be, \Lambda_c^+ \rightarrow p K^- \pi^+$
196^{+23}_{-20}	101	BARLAG	89	NA32	$p K^- \pi^+ + c.c.$
$220 \pm 30 \pm 20$	97	ANJOS	88B	E691	$p K^- \pi^+ + c.c.$



Λ_c^+ DECAY MODES

Nearly all branching fractions of the Λ_c^+ are measured relative to the $p K^- \pi^+$ mode, but there are no model-independent measurements of this branching fraction. We explain how we arrive at our value of $B(\Lambda_c^+ \rightarrow p K^- \pi^+)$ in a Note at the beginning of the branching-ratio measurements, below. When this branching fraction is eventually well determined, all the other branching fractions will slide up or down proportionally as the true value differs from the value we use here.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
------	--------------------------------	-----------------------------------

Hadronic modes with a p : $S = -1$ final states

Γ_1	$p\bar{K}^0$	(2.3 \pm 0.6) %
Γ_2	$pK^-\pi^+$	[a] (5.0 \pm 1.3) %
Γ_3	$p\bar{K}^*(892)^0$	[b] (1.6 \pm 0.5) %
Γ_4	$\Delta(1232)^{++}K^-$	(8.6 \pm 3.0) $\times 10^{-3}$
Γ_5	$\Lambda(1520)\pi^+$	[b] (1.8 \pm 0.6) %
Γ_6	$pK^-\pi^+$ nonresonant	(2.8 \pm 0.8) %
Γ_7	$p\bar{K}^0\pi^0$	(3.3 \pm 1.0) %
Γ_8	$p\bar{K}^0\eta$	(1.2 \pm 0.4) %
Γ_9	$p\bar{K}^0\pi^+\pi^-$	(2.6 \pm 0.7) %
Γ_{10}	$pK^-\pi^+\pi^0$	(3.4 \pm 1.0) %
Γ_{11}	$pK^*(892)^-\pi^+$	[b] (1.1 \pm 0.5) %
Γ_{12}	$p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	(3.6 \pm 1.2) %
Γ_{13}	$\Delta(1232)\bar{K}^*(892)$	seen
Γ_{14}	$pK^-\pi^+\pi^+\pi^-$	(1.1 \pm 0.8) $\times 10^{-3}$
Γ_{15}	$pK^-\pi^+\pi^0\pi^0$	(8 \pm 4) $\times 10^{-3}$
Γ_{16}	$pK^-\pi^+3\pi^0$	

Hadronic modes with a p : $S = 0$ final states

Γ_{17}	$p\pi^+\pi^-$	(3.5 \pm 2.0) $\times 10^{-3}$
Γ_{18}	$p f_0(980)^-$	[b] (2.8 \pm 1.9) $\times 10^{-3}$
Γ_{19}	$p\pi^+\pi^+\pi^-\pi^-$	(1.8 \pm 1.2) $\times 10^{-3}$
Γ_{20}	pK^+K^-	(7.7 \pm 3.5) $\times 10^{-4}$
Γ_{21}	$p\phi$	[b] (8.2 \pm 2.7) $\times 10^{-4}$
Γ_{22}	$pK^+K^- \text{ non-}\phi$	(3.5 \pm 1.7) $\times 10^{-4}$

Hadronic modes with a hyperon: $S = -1$ final states

Γ_{23}	$\Lambda\pi^+$	(1.07 \pm 0.28) %
Γ_{24}	$\Lambda\pi^+\pi^0$	(3.6 \pm 1.3) %
Γ_{25}	$\Lambda\rho^+$	< 5 % CL=95%
Γ_{26}	$\Lambda\pi^+\pi^+\pi^-$	(2.6 \pm 0.7) %
Γ_{27}	$\Sigma(1385)^+\pi^+\pi^-$, $\Sigma^{*+} \rightarrow$	(7 \pm 4) $\times 10^{-3}$
Γ_{28}	$\Lambda\pi^+$ $\Sigma(1385)^-\pi^+\pi^+$, $\Sigma^{*-} \rightarrow$	(5.5 \pm 1.7) $\times 10^{-3}$
Γ_{29}	$\Lambda\pi^-\rho^0$	(1.1 \pm 0.5) %
Γ_{30}	$\Sigma(1385)^+\rho^0$, $\Sigma^{*+} \rightarrow \Lambda\pi^+$	(3.7 \pm 3.1) $\times 10^{-3}$
Γ_{31}	$\Lambda\pi^+\pi^+\pi^-$ nonresonant	< 8 $\times 10^{-3}$ CL=90%
Γ_{32}	$\Lambda\pi^+\pi^+\pi^-\pi^0$ total	(1.8 \pm 0.8) %
Γ_{33}	$\Lambda\pi^+\eta$	[b] (1.8 \pm 0.6) %
Γ_{34}	$\Sigma(1385)^+\eta$	[b] (8.5 \pm 3.3) $\times 10^{-3}$
Γ_{35}	$\Lambda\pi^+\omega$	[b] (1.2 \pm 0.5) %
Γ_{36}	$\Lambda\pi^+\pi^+\pi^-\pi^0$, no η or ω	< 7 $\times 10^{-3}$ CL=90%
Γ_{37}	$\Lambda K^+\bar{K}^0$	(4.7 \pm 1.5) $\times 10^{-3}$ S=1.2
Γ_{38}	$\Xi(1690)^0K^+$, $\Xi^{*0} \rightarrow \Lambda\bar{K}^0$	(1.3 \pm 0.5) $\times 10^{-3}$

Γ_{39}	$\Sigma^0 \pi^+$	(1.05 \pm 0.28) %	
Γ_{40}	$\Sigma^+ \pi^0$	(1.00 \pm 0.34) %	
Γ_{41}	$\Sigma^+ \eta$	(5.5 \pm 2.3) $\times 10^{-3}$	
Γ_{42}	$\Sigma^+ \pi^+ \pi^-$	(3.6 \pm 1.0) %	
Γ_{43}	$\Sigma^+ \rho^0$	< 1.4 %	CL=95%
Γ_{44}	$\Sigma^- \pi^+ \pi^+$	(1.9 \pm 0.8) %	
Γ_{45}	$\Sigma^0 \pi^+ \pi^0$	(1.8 \pm 0.8) %	
Γ_{46}	$\Sigma^0 \pi^+ \pi^+ \pi^-$	(8.3 \pm 3.1) $\times 10^{-3}$	
Γ_{47}	$\Sigma^+ \pi^+ \pi^- \pi^0$	—	
Γ_{48}	$\Sigma^+ \omega$	[b] (2.7 \pm 1.0) %	
Γ_{49}	$\Sigma^+ K^+ K^-$	(2.8 \pm 0.8) $\times 10^{-3}$	
Γ_{50}	$\Sigma^+ \phi$	[b] (3.2 \pm 1.0) $\times 10^{-3}$	
Γ_{51}	$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow$	(8.2 \pm 3.1) $\times 10^{-4}$	
Γ_{52}	$\Sigma^+ K^+ K^-$ nonresonant	< 7 $\times 10^{-4}$	CL=90%
Γ_{53}	$\Xi^0 K^+$	(3.9 \pm 1.4) $\times 10^{-3}$	
Γ_{54}	$\Xi^- K^+ \pi^+$	(5.1 \pm 1.4) $\times 10^{-3}$	
Γ_{55}	$\Xi(1530)^0 K^+$	[b] (2.6 \pm 1.0) $\times 10^{-3}$	

Hadronic modes with a hyperon: $S = 0$ final states

Γ_{56}	ΛK^+	(5.0 \pm 1.6) $\times 10^{-4}$	
Γ_{57}	$\Lambda K^+ \pi^+ \pi^-$	< 4 $\times 10^{-4}$	CL=90%
Γ_{58}	$\Sigma^0 K^+$	(4.2 \pm 1.3) $\times 10^{-4}$	
Γ_{59}	$\Sigma^0 K^+ \pi^+ \pi^-$	< 2.1 $\times 10^{-4}$	CL=90%
Γ_{60}	$\Sigma^+ K^+ \pi^-$	(1.7 \pm 0.7) $\times 10^{-3}$	
Γ_{61}	$\Sigma^+ K^*(892)^0$	[b] (2.8 \pm 1.1) $\times 10^{-3}$	
Γ_{62}	$\Sigma^- K^+ \pi^+$	< 1.0 $\times 10^{-3}$	CL=90%

Doubly Cabibbo-suppressed modes

Γ_{63}	$p K^+ \pi^-$	< 2.3 $\times 10^{-4}$	CL=90%
---------------	---------------	------------------------	--------

Semileptonic modes

Γ_{64}	$\Lambda \ell^+ \nu_\ell$	[c] (2.0 \pm 0.6) %	
Γ_{65}	$\Lambda e^+ \nu_e$	(2.1 \pm 0.6) %	
Γ_{66}	$\Lambda \mu^+ \nu_\mu$	(2.0 \pm 0.7) %	

Inclusive modes

Γ_{67}	$e^+ \text{ anything}$	(4.5 \pm 1.7) %	
Γ_{68}	$p e^+ \text{ anything}$	(1.8 \pm 0.9) %	
Γ_{69}	$\Lambda e^+ \text{ anything}$		
Γ_{70}	$p \text{ anything}$	(50 \pm 16) %	
Γ_{71}	$p \text{ anything (no } \Lambda)$	(12 \pm 19) %	
Γ_{72}	$p \text{ hadrons}$		
Γ_{73}	$n \text{ anything}$	(50 \pm 16) %	

Γ_{74}	n anything (no Λ)	(29 ± 17) %	
Γ_{75}	Λ anything	(35 ± 11) %	S=1.4
Γ_{76}	Σ^\pm anything	[d] (10 ± 5) %	
Γ_{77}	3prongs	(24 ± 8) %	

$\Delta C = 1$ weak neutral current (C1**) modes, or
Lepton number (**L**) violating modes**

Γ_{78}	$p\mu^+\mu^-$	$C1$	< 3.4	$\times 10^{-4}$	CL=90%
Γ_{79}	$\Sigma^-\mu^+\mu^+$	L	< 7.0	$\times 10^{-4}$	CL=90%

[a] See the note on “ Λ_c^+ Branching Fractions” below.

[b] This branching fraction includes all the decay modes of the final-state resonance.

[c] An ℓ indicates an e or a μ mode, not a sum over these modes.

[d] The value is for the sum of the charge states or particle/antiparticle states indicated.

CONSTRAINED FIT INFORMATION

An overall fit to 16 branching ratios uses 30 measurements and one constraint to determine 11 parameters. The overall fit has a $\chi^2 = 15.2$ for 20 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x23	96								
x26	97	93							
x37	82	83	80						
x39	95	98	92	82					
x42	91	87	89	75	86				
x46	69	66	70	57	66	63			
x49	87	83	85	71	82	93	60		
x50	84	80	81	69	79	90	58	84	
x54	93	96	90	80	94	85	64	81	78
	x2	x23	x26	x37	x39	x42	x46	x49	x50

A REVIEW GOES HERE – Check our WWW List of Reviews

Λ_c^+ BRANCHING RATIOS

Hadronic modes with a p : $S = -1$ final states

$\Gamma(p\bar{K}^0)/\Gamma(pK^-\pi^+)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.47±0.04 OUR AVERAGE				
0.46±0.02±0.04	1025	ALAM	98	CLE2 $e^+e^- \approx \gamma(4S)$
0.44±0.07±0.05	133	AVERY	91	CLEO $e^+e^- 10.5 \text{ GeV}$
0.55±0.17±0.14	45	ANJOS	90	$\gamma\text{Be } 70\text{--}260 \text{ GeV}$
0.62±0.15±0.03	73	ALBRECHT	88C	$e^+e^- 10 \text{ GeV}$

$\Gamma(pK^-\pi^+)/\Gamma_{\text{total}}$

Γ_2/Γ

See the note on " Λ_c^+ Branching Fractions" above.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.050±0.013 OUR FIT				
0.050±0.013		PDG	02	See note at top of ratios
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.050±0.005±0.012	1205	² JAFFE	00	CLE2 $e^+e^- 10.52\text{--}10.58 \text{ GeV}$
0.041±0.010		^{3,4} ALBRECHT	920	ARG $e^+e^- \approx \gamma(4S)$
0.044±0.012		^{3,5} CRAWFORD	92	CLEO $e^+e^- 10.5 \text{ GeV}$

² JAFFE 00 assumes that a \bar{D} meson and an antiproton in opposite hemispheres tags for a Λ_c^+ in the hemisphere of the \bar{p} . The fraction of such $\bar{D}\bar{p}$ events with a $\Lambda_c^+ \rightarrow pK^-\pi^+$ decay then gives the $pK^-\pi^+$ branching fraction. See the paper for assumptions, caveats, etc.

³ To extract $\Gamma(pK^-\pi^+)/\Gamma_{\text{total}}$, we use $B(\bar{B} \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (0.28 \pm 0.06)\%$, which is the average of measurements from ARGUS (ALBRECHT 88C) and CLEO (CRAWFORD 92).

⁴ ALBRECHT 920 measures $B(\bar{B} \rightarrow \Lambda_c^+ X) = (6.8 \pm 0.5 \pm 0.3)\%$.

⁵ CRAWFORD 92 measures $B(\bar{B} \rightarrow \Lambda_c^+ X) = (6.4 \pm 0.8 \pm 0.8)\%$.

$\Gamma(p\bar{K}^*(892)^0)/\Gamma(pK^-\pi^+)$

Γ_3/Γ_2

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.31±0.04 OUR AVERAGE				
0.29±0.04±0.03		⁶ AITALA	00	E791 $\pi^- N, 500 \text{ GeV}$
$0.35^{+0.06}_{-0.07} \pm 0.03$	39	BOZEK	93	NA32 $\pi^- \text{Cu } 230 \text{ GeV}$
0.42±0.24	12	BASILE	81B	CNTR $p\bar{p} \rightarrow \Lambda_c^+ e^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.35±0.11		BARLAG	90D	NA32 See BOZEK 93

⁶ AITALA 00 makes a coherent 5-dimensional amplitude analysis of 946 ± 38 $\Lambda_c^+ \rightarrow pK^-\pi^+$ decays.

$\Gamma(\Delta(1232)^{++} K^-)/\Gamma(pK^-\pi^+)$ Γ_4/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.17±0.04 OUR AVERAGE	Error includes scale factor of 1.1.			
0.18±0.03±0.03	7	AITALA 00	E791	$\pi^- N$, 500 GeV
0.12 ^{+0.04} _{-0.05} ±0.05	14	BOZEK 93	NA32	$\pi^- Cu$ 230 GeV
0.40±0.17	17	BASILE 81B	CNTR	$p p \rightarrow \Lambda_c^+ e^- X$

⁷ AITALA 00 makes a coherent 5-dimensional amplitude analysis of $946 \pm 38 \Lambda_c^+ \rightarrow p K^- \pi^+$ decays.

 $\Gamma(\Lambda(1520)\pi^+)/\Gamma(pK^-\pi^+)$ Γ_5/Γ_2 Unseen decay modes of the $\Lambda(1520)$ are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.35±0.08 OUR AVERAGE				
0.34±0.08±0.05	8	AITALA 00	E791	$\pi^- N$, 500 GeV
0.40 ^{+0.18} _{-0.13} ±0.09	12	BOZEK 93	NA32	$\pi^- Cu$ 230 GeV

⁸ AITALA 00 makes a coherent 5-dimensional amplitude analysis of $946 \pm 38 \Lambda_c^+ \rightarrow p K^- \pi^+$ decays.

 $\Gamma(pK^-\pi^+ \text{ nonresonant})/\Gamma(pK^-\pi^+)$ Γ_6/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.55±0.06 OUR AVERAGE				
0.55±0.06±0.04	9	AITALA 00	E791	$\pi^- N$, 500 GeV
0.56 ^{+0.07} _{-0.09} ±0.05	71	BOZEK 93	NA32	$\pi^- Cu$ 230 GeV

⁹ AITALA 00 makes a coherent 5-dimensional amplitude analysis of $946 \pm 38 \Lambda_c^+ \rightarrow p K^- \pi^+$ decays.

 $\Gamma(p\bar{K}^0\pi^0)/\Gamma(pK^-\pi^+)$ Γ_7/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.66±0.05±0.07	774	ALAM 98	CLE2	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(p\bar{K}^0\eta)/\Gamma(pK^-\pi^+)$ Γ_8/Γ_2 Unseen decay modes of the η are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.25±0.04±0.04	57	AMMAR 95	CLE2	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(p\bar{K}^0\pi^+\pi^-)/\Gamma(pK^-\pi^+)$ Γ_9/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.51±0.06 OUR AVERAGE				
0.52±0.04±0.05	985	ALAM 98	CLE2	$e^+ e^- \approx \gamma(4S)$
0.43±0.12±0.04	83	AVERY 91	CLEO	$e^+ e^- 10.5$ GeV
0.98±0.36±0.08	12	BARLAG 90D	NA32	$\pi^- 230$ GeV

 $\Gamma(pK^-\pi^+\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{10}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.67±0.04±0.11	2606	ALAM 98	CLE2	$e^+ e^- \approx \gamma(4S)$

$\Gamma(pK^*(892)^-\pi^+)/\Gamma(p\bar{K}^0\pi^+\pi^-)$ Γ_{11}/Γ_9 Unseen decay modes of the $K^*(892)^-$ are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.44±0.14	17	ALEEV	94	BIS2 nN 20–70 GeV

 $\Gamma(p(K^-\pi^+)_{\text{nonresonant}}\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{12}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.73±0.12±0.05	67	BOZEK	93	NA32 π^- Cu 230 GeV

 $\Gamma(\Delta(1232)\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	35	AMENDOLIA	87	SPEC γ Ge-Si

 $\Gamma(pK^-\pi^+\pi^+\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{14}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.022±0.015		BARLAG	90D	NA32 π^- 230 GeV

 $\Gamma(pK^-\pi^+\pi^0\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{15}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.16±0.07±0.03	15	BOZEK	93	NA32 π^- Cu 230 GeV

 $\Gamma(pK^-\pi^+3\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{16}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.10±0.06±0.02	8	BOZEK	93	NA32 π^- Cu 230 GeV

Hadronic modes with a p : $S = 0$ final states

 $\Gamma(p\pi^+\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{17}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.069±0.036		BARLAG	90D	NA32 π^- 230 GeV

 $\Gamma(pf_0(980))/\Gamma(pK^-\pi^+)$ Γ_{18}/Γ_2 Unseen decay modes of the $f_0(980)$ are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.055±0.036		BARLAG	90D	NA32 π^- 230 GeV

 $\Gamma(p\pi^+\pi^+\pi^-\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{19}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.036±0.023		BARLAG	90D	NA32 π^- 230 GeV

 $\Gamma(pK^+K^-)/\Gamma(pK^-\pi^+)$ Γ_{20}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.015±0.006 OUR AVERAGE				Error includes scale factor of 2.1.

0.014±0.002±0.002	676	ABE	02C	BELL $e^+e^- \approx \gamma(4S)$
0.039±0.009±0.007	214	ALEXANDER	96C	CLE2 $e^+e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.096±0.029±0.010	30	FRABETTI	93H E687	γ Be, \bar{E}_γ 220 GeV
0.048±0.027		BARLAG	90D NA32	π^- 230 GeV

$\Gamma(p\phi)/\Gamma(pK^-\pi^+)$ Γ_{21}/Γ_2 Unseen decay modes of the ϕ are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0164±0.0032 OUR AVERAGE				Error includes scale factor of 1.2.
0.015 ± 0.002 ± 0.002	345	ABE	02C BELL	$e^+e^- \approx \gamma(4S)$
0.024 ± 0.006 ± 0.003	54	ALEXANDER	96C CLE2	$e^+e^- \approx \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.040 ± 0.027		BARLAG	90D NA32	$\pi^- 230 \text{ GeV}$

 $\Gamma(pK^+K^-\text{non-}\phi)/\Gamma(pK^-\pi^+)$ Γ_{22}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.007±0.002 ± 0.002	344	ABE	02C BELL	$e^+e^- \approx \gamma(4S)$

Hadronic modes with a hyperon: $S = -1$ final states $\Gamma(\Lambda\pi^+)/\Gamma(pK^-\pi^+)$ Γ_{23}/Γ_2

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.214±0.016 OUR FIT					Error includes scale factor of 1.1.
0.204±0.019 OUR AVERAGE					
0.217±0.013±0.020		750	LINK	05F FOCS	γ nucleus, $\bar{E}_\gamma \approx 180 \text{ GeV}$
0.18 ± 0.03 ± 0.04			ALBRECHT	92 ARG	$e^+e^- \approx 10.4 \text{ GeV}$
0.18 ± 0.03 ± 0.03		87	AVERY	91 CLEO	$e^+e^- 10.5 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.33			ANJOS	90 E691	γ Be 70–260 GeV
<0.16			ALBRECHT	88C ARG	$e^+e^- 10 \text{ GeV}$

 $\Gamma(\Lambda\pi^+\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{24}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.73±0.09±0.16	464	AVERY	94 CLE2	$e^+e^- \approx \gamma(3S), \gamma(4S)$

 $\Gamma(\Lambda\rho^+)/\Gamma(pK^-\pi^+)$ Γ_{25}/Γ_2

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.95	95	AVERY	94 CLE2	$e^+e^- \approx \gamma(3S), \gamma(4S)$

 $\Gamma(\Lambda\pi^+\pi^+\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{26}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.525±0.032 OUR FIT				
0.522±0.032 OUR AVERAGE				
0.508±0.024±0.024	1356	LINK	05F FOCS	γ nucleus, $\bar{E}_\gamma \approx 180 \text{ GeV}$
0.65 ± 0.11 ± 0.12	289	AVERY	91 CLEO	$e^+e^- 10.5 \text{ GeV}$
0.82 ± 0.29 ± 0.27	44	ANJOS	90 E691	γ Be 70–260 GeV
0.94 ± 0.41 ± 0.13	10	BARLAG	90D NA32	$\pi^- 230 \text{ GeV}$
0.61 ± 0.16 ± 0.04	105	ALBRECHT	88C ARG	$e^+e^- 10 \text{ GeV}$

 $\Gamma(\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow \Lambda\pi^+)/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_{27}/Γ_{26}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.28±0.10±0.08	LINK	05F FOCS	γ nucleus, $\bar{E}_\gamma \approx 180 \text{ GeV}$

$\Gamma(\Sigma(1385)^-\pi^+\pi^+,\Sigma^{*-}\rightarrow\Lambda\pi^-)/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_{28}/Γ_{26}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.21±0.03±0.02	LINK	05F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\Lambda\pi^+\rho^0)/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_{29}/Γ_{26}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.40±0.12±0.12	LINK	05F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\Sigma(1385)^+\rho^0,\Sigma^{*+}\rightarrow\Lambda\pi^+)/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_{30}/Γ_{26}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.14±0.09±0.07	LINK	05F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\Lambda\pi^+\pi^+\pi^-\text{nonresonant})/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_{31}/Γ_{26}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.3	90	LINK	05F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(p\bar{K}^0\pi^+\pi^-)/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_9/Γ_{26}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6±1.2	ALEEV	96	SPEC	n nucleus, 50 GeV/c
4.3±1.2	130	ALEEV	84	BIS2 nC 40–70 GeV

 $\Gamma(\Lambda\pi^+\pi^-\pi^0\text{total})/\Gamma(pK^-\pi^+)$ Γ_{32}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.36±0.09±0.09	50	10 CRONIN-HEN..03	CLE3	$e^+e^- \approx \gamma(4S)$

¹⁰ CRONIN-HENNESSY 03 finds this channel to be dominately $\Lambda\eta\pi^+$ and $\Lambda\omega\pi^+$; see below.

 $\Gamma(\Lambda\pi^+\eta)/\Gamma(pK^-\pi^+)$ Γ_{33}/Γ_2

Unseen decay modes of the η are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.36±0.07 OUR AVERAGE				
0.41±0.17±0.10	11	CRONIN-HEN..03	CLE3	$e^+e^- \approx \gamma(4S)$
0.35±0.05±0.06	116	AMMAR	95	CLE2 $e^+e^- \approx \gamma(4S)$

 $\Gamma(\Sigma(1385)^+\eta)/\Gamma(pK^-\pi^+)$ Γ_{34}/Γ_2

Unseen decay modes of the $\Sigma(1385)^+$ and η are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.17±0.04±0.03	54	AMMAR	95	CLE2 $e^+e^- \approx \gamma(4S)$

 $\Gamma(\Lambda\pi^+\omega)/\Gamma(pK^-\pi^+)$ Γ_{35}/Γ_2

Unseen decay modes of the ω are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.24±0.06±0.06	32	CRONIN-HEN..03	CLE3	$e^+e^- \approx \gamma(4S)$

 $\Gamma(\Lambda\pi^+\pi^-\pi^0,\text{no }\eta\text{ or }\omega)/\Gamma(pK^-\pi^+)$ Γ_{36}/Γ_2

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.13	90	CRONIN-HEN..03	CLE3	$e^+e^- \approx \gamma(4S)$

$\Gamma(\Lambda K^+ \bar{K}^0)/\Gamma(p K^- \pi^+)$ Γ_{37}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.093±0.018 OUR FIT	Error includes scale factor of 1.7.			
0.131±0.020 OUR AVERAGE				
0.142±0.018±0.022	251	LINK	05F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
0.12 ± 0.02 ± 0.02	59	AMMAR	95	CLE2 $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Lambda \bar{K}^0)/\Gamma(\Lambda K^+ \bar{K}^0)$ Γ_{38}/Γ_{37}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.28±0.07 OUR AVERAGE				
0.32±0.10±0.04	84±24	LINK	05F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
0.26±0.08±0.03	93	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Lambda K^+ \bar{K}^0)/\Gamma(\Lambda \pi^+)$ Γ_{37}/Γ_{23}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.43 ± 0.08 OUR FIT	Error includes scale factor of 2.0.			
0.395±0.026±0.036	460 ± 30	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^0 \pi^+)/\Gamma(p K^- \pi^+)$ Γ_{39}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.210±0.018 OUR FIT				
0.20 ± 0.04 OUR AVERAGE				
0.21 ± 0.02 ± 0.04	196	AVERY	94	CLE2 $e^+ e^- \approx \gamma(3S), \gamma(4S)$
0.17 ± 0.06 ± 0.04		ALBRECHT	92	ARG $e^+ e^- \approx 10.4$ GeV

 $\Gamma(\Sigma^0 \pi^+)/\Gamma(\Lambda \pi^+)$ Γ_{39}/Γ_{23}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.98 ± 0.05 OUR FIT				
0.98 ± 0.05 OUR AVERAGE				
0.977±0.015±0.051	33k	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$
1.09 ± 0.11 ± 0.19	750	LINK	05F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\Sigma^+ \pi^0)/\Gamma(p K^- \pi^+)$ Γ_{40}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.20±0.03±0.03	93	KUBOTA	93	CLE2 $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^+ \eta)/\Gamma(p K^- \pi^+)$ Γ_{41}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.11±0.03±0.02	26	AMMAR	95	CLE2 $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^+ \pi^+ \pi^-)/\Gamma(p K^- \pi^+)$ Γ_{42}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.73±0.08 OUR FIT				
0.68±0.09 OUR AVERAGE				
0.74±0.07±0.09	487	KUBOTA	93	CLE2 $e^+ e^- \approx \gamma(4S)$
0.54 ^{+0.18} _{-0.15}	11	BARLAG	92	NA32 π^- Cu 230 GeV

$\Gamma(\Sigma^+ \rho^0)/\Gamma(p K^- \pi^+)$					Γ_{43}/Γ_2
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.27	95	KUBOTA	93	CLE2	$e^+ e^- \approx \gamma(4S)$
$\Gamma(\Sigma^- \pi^+ \pi^+)/\Gamma(\Sigma^+ \pi^+ \pi^-)$					Γ_{44}/Γ_{42}
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.53±0.15±0.07	56	FRABETTI	94E	E687	γ Be, \bar{E}_γ 220 GeV
$\Gamma(\Sigma^0 \pi^+ \pi^0)/\Gamma(p K^- \pi^+)$					Γ_{45}/Γ_2
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.36±0.09±0.10	117	AVERY	94	CLE2	$e^+ e^- \approx \gamma(3S), \gamma(4S)$
$\Gamma(\Sigma^0 \pi^+ \pi^+ \pi^-)/\Gamma(p K^- \pi^+)$					Γ_{46}/Γ_2
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.17±0.04 OUR FIT					
0.21±0.05±0.05	90	AVERY	94	CLE2	$e^+ e^- \approx \gamma(3S), \gamma(4S)$
$\Gamma(\Sigma^0 \pi^+ \pi^+ \pi^-)/\Gamma(\Lambda \pi^+ \pi^+ \pi^-)$					Γ_{46}/Γ_{26}
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.31±0.08 OUR FIT					
0.26±0.06±0.09	480	LINK	05F	FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
$\Gamma(\Sigma^+ \omega)/\Gamma(p K^- \pi^+)$					Γ_{48}/Γ_2
Unseen decay modes of the ω are included.					
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.54±0.13±0.06	107	KUBOTA	93	CLE2	$e^+ e^- \approx \gamma(4S)$
$\Gamma(\Sigma^+ K^+ K^-)/\Gamma(p K^- \pi^+)$					Γ_{49}/Γ_2
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.057±0.008 OUR FIT					
0.070±0.011±0.011	59	AVERY	93	CLE2	$e^+ e^- \approx 10.5$ GeV
$\Gamma(\Sigma^+ K^+ K^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$					Γ_{49}/Γ_{42}
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.078±0.009 OUR FIT					
0.074±0.009 OUR AVERAGE					
0.076±0.007±0.009	246	ABE	02C	BELL	$e^+ e^- \approx \gamma(4S)$
0.071±0.011±0.011	103	LINK	02G	FOCS	γ nucleus, ≈ 180 GeV
$\Gamma(\Sigma^+ \phi)/\Gamma(p K^- \pi^+)$					Γ_{50}/Γ_2
Unseen decay modes of the ϕ are included.					
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.063±0.011 OUR FIT					
0.069±0.023±0.016	26	AVERY	93	CLE2	$e^+ e^- \approx 10.5$ GeV
$\Gamma(\Sigma^+ \phi)/\Gamma(\Sigma^+ \pi^+ \pi^-)$					Γ_{50}/Γ_{42}
Unseen decay modes of the ϕ are included.					
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.087±0.012 OUR FIT					
0.086±0.012 OUR AVERAGE					
0.085±0.012±0.012	129	ABE	02C	BELL	$e^+ e^- \approx \gamma(4S)$
0.087±0.016±0.006	57	LINK	02G	FOCS	γ nucleus, ≈ 180 GeV

$\Gamma(\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Sigma^+ K^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{51}/Γ_{42}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.023±0.005 OUR AVERAGE				
0.023±0.005±0.005	75	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$
0.022±0.006±0.006	34	LINK	02G	FOCS γ nucleus, ≈ 180 GeV

 $\Gamma(\Sigma^+ K^+ K^- \text{ nonresonant})/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{52}/Γ_{42}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.018	90	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.028	90	LINK	02G	FOCS γ nucleus, ≈ 180 GeV

 $\Gamma(\Xi^0 K^+)/\Gamma(p K^- \pi^+)$ Γ_{53}/Γ_2


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.078±0.013±0.013	56	AVERY	93	CLE2 $e^+ e^- \approx 10.5$ GeV

 $\Gamma(\Xi^- K^+ \pi^+)/\Gamma(p K^- \pi^+)$ Γ_{54}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.102±0.010 OUR FIT				Error includes scale factor of 1.1.
0.098±0.021 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.14 ± 0.03 ± 0.02	34	ALBRECHT	95B	ARG $e^+ e^- \approx 10.4$ GeV
0.079±0.013±0.014	60	AVERY	93	CLE2 $e^+ e^- \approx 10.5$ GeV
0.15 ± 0.04 ± 0.03	30	AVERY	91	CLEO $e^+ e^- 10.5$ GeV

WEIGHTED AVERAGE

0.098±0.021 (Error scaled by 1.3)


 $\Gamma(\Xi^- K^+ \pi^+)/\Gamma(p K^- \pi^+)$

$\Gamma(\Xi(1530)^0 K^+)/\Gamma(p K^- \pi^+)$ Γ_{55}/Γ_2 Unseen decay modes of the $\Xi(1530)^0$ are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.052±0.014 OUR AVERAGE				
0.05 ± 0.02 ± 0.01	11	ALBRECHT	95B	ARG $e^+ e^- \approx 10.4$ GeV
0.053 ± 0.016 ± 0.010	24	AVERY	93	CLE2 $e^+ e^- \approx 10.5$ GeV

 $\Gamma(\Xi^- K^+ \pi^+)/\Gamma(\Lambda \pi^+)$ Γ_{54}/Γ_{23}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.47 ± 0.04 OUR FIT				
0.480±0.016±0.039	2665 ± 84	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$

Hadronic modes with a hyperon: $S = 0$ final states $\Gamma(\Lambda K^+)/\Gamma(\Lambda \pi^+)$ Γ_{56}/Γ_{23}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.047±0.009 OUR AVERAGE				
Error includes scale factor of 1.8.				
0.044 ± 0.004 ± 0.003	1162 ± 101	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$
0.074 ± 0.010 ± 0.012	265	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Lambda K^+ \pi^+ \pi^-)/\Gamma(\Lambda \pi^+)$ Γ_{57}/Γ_{23}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.1 × 10⁻²	90	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^0 K^+)/\Gamma(\Sigma^0 \pi^+)$ Γ_{58}/Γ_{39}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.040±0.006 OUR AVERAGE				
0.038 ± 0.005 ± 0.003	366 ± 52	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$
0.056 ± 0.014 ± 0.008	75	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^0 K^+ \pi^+ \pi^-)/\Gamma(\Sigma^0 \pi^+)$ Γ_{59}/Γ_{39}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.0 × 10⁻²	90	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^+ K^+ \pi^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{60}/Γ_{42}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.047±0.011±0.008				
105	ABE	02C	BELL	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^+ K^*(892)^0)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{61}/Γ_{42} Unseen decay modes of the $K^*(892)^0$ are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.078±0.018±0.013				
49	LINK	02G	FOCS	γ nucleus, ≈ 180 GeV

 $\Gamma(\Sigma^- K^+ \pi^+)/\Gamma(\Sigma^+ K^*(892)^0)$ Γ_{62}/Γ_{61}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.35	90	LINK	02G	FOCS γ nucleus, ≈ 180 GeV

Doubly Cabibbo-suppressed modes

$\Gamma(pK^+\pi^-)/\Gamma(pK^-\pi^+)$	Γ_{63}/Γ_2			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0046	90	LINK	05K	FOCS R = (0.05 ± 0.26 ± 0.02)%

Semileptonic modes

$\Gamma(\Lambda\ell^+\nu_\ell)/\Gamma(pK^-\pi^+)$	Γ_{64}/Γ_2	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
0.41 ± 0.05 OUR AVERAGE		
0.42 ± 0.07	PDG	02 Our $\Gamma(\Lambda e^+\nu_e)/\Gamma(pK^-\pi^+)$
0.39 ± 0.08	PDG	02 Our $\Gamma(\Lambda \mu^+\nu_\mu)/\Gamma(pK^-\pi^+)$

$\Gamma(\Lambda e^+\nu_e)/\Gamma(pK^-\pi^+)$	Γ_{65}/Γ_2		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.42 ± 0.07 OUR AVERAGE			
0.43 ± 0.08	11,12 BERGFELD	94 CLE2	$e^+e^- \approx \gamma(4S)$
0.38 ± 0.14	12,13 ALBRECHT	91G ARG	$e^+e^- \approx 10.4 \text{ GeV}$
¹¹ BERGFELD 94 measures $\sigma(e^+e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+\nu_e) = (4.87 \pm 0.28 \pm 0.69) \text{ pb.}$			
¹² To extract $\Gamma(\Lambda_c^+ \rightarrow \Lambda e^+\nu_e)/\Gamma(\Lambda_c^+ \rightarrow pK^-\pi^+)$, we use $\sigma(e^+e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c \rightarrow pK^-\pi^+) = (11.2 \pm 1.3) \text{ pb}$, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (AVERY 91).			
¹³ ALBRECHT 91G measures $\sigma(e^+e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+\nu_e) = (4.20 \pm 1.28 \pm 0.71) \text{ pb.}$			

$\Gamma(\Lambda\mu^+\nu_\mu)/\Gamma(pK^-\pi^+)$	Γ_{66}/Γ_2		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.39 ± 0.08 OUR AVERAGE			
0.40 ± 0.09	14,15 BERGFELD	94 CLE2	$e^+e^- \approx \gamma(4S)$
0.35 ± 0.20	15,16 ALBRECHT	91G ARG	$e^+e^- \approx 10.4 \text{ GeV}$
¹⁴ BERGFELD 94 measures $\sigma(e^+e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda\mu^+\nu_\mu) = (4.43 \pm 0.51 \pm 0.64) \text{ pb.}$			
¹⁵ To extract $\Gamma(\Lambda_c^+ \rightarrow \Lambda\mu^+\nu_\mu)/\Gamma(\Lambda_c^+ \rightarrow pK^-\pi^+)$, we use $\sigma(e^+e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c \rightarrow pK^-\pi^+) = (11.2 \pm 1.3) \text{ pb}$, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (AVERY 91).			
¹⁶ ALBRECHT 91G measures $\sigma(e^+e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda\mu^+\nu_\mu) = (3.91 \pm 2.02 \pm 0.90) \text{ pb.}$			

Inclusive modes

$\Gamma(e^+\text{anything})/\Gamma_{\text{total}}$	Γ_{67}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.045 ± 0.017	VELLA	82 MRK2	$e^+e^- 4.5\text{--}6.8 \text{ GeV}$

$\Gamma(p e^+ \text{anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{68}/Γ
0.018±0.009	17 VELLA	82	MRK2 $e^+ e^-$ 4.5–6.8 GeV	

17 VELLA 82 includes protons from Λ decay.

$\Gamma(\Lambda e^+ \text{anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{69}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.011±0.008	18 VELLA	82	MRK2 $e^+ e^-$ 4.5–6.8 GeV	

18 VELLA 82 includes Λ 's from Σ^0 decay.

$\Gamma(p \text{ anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{70}/Γ
0.50±0.08±0.14	19 CRAWFORD	92	CLEO $e^+ e^-$ 10.5 GeV	

19 This CRAWFORD 92 value includes protons from Λ decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(p \text{ anything (no } \Lambda))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{71}/Γ
0.12±0.10±0.16	CRAWFORD	92	CLEO $e^+ e^-$ 10.5 GeV	

$\Gamma(n \text{ anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{73}/Γ
0.50±0.08±0.14	20 CRAWFORD	92	CLEO $e^+ e^-$ 10.5 GeV	

20 This CRAWFORD 92 value includes neutrons from Λ decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(n \text{ anything (no } \Lambda))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{74}/Γ
0.29±0.09±0.15	CRAWFORD	92	CLEO $e^+ e^-$ 10.5 GeV	


$\Gamma(p \text{ hadrons})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{72}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.41±0.24	ADAMOVICH	87	EMUL γA 20–70 GeV/c	

$\Gamma(\Lambda \text{ anything})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{75}/Γ
0.35±0.11 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.	
0.59±0.10±0.12		CRAWFORD	92	CLEO $e^+ e^-$ 10.5 GeV	
0.49±0.24		ADAMOVICH	87	EMUL γA 20–70 GeV/c	
0.23±0.10	8	21 ABE	86	HYBR 20 GeV γp	

²¹ ABE 86 includes Λ 's from Σ^0 decay.



$\Gamma(\Sigma^\pm \text{anything})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{76}/Γ
0.1±0.05	5	ABE	86	HYBR 20 GeV γp	

$\Gamma(3\text{prongs})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{77}/Γ
0.24±0.07±0.04	KAYIS-TOPAK.03	CHRS	ν_μ emulsion, $\bar{E}=27$ GeV	

Rare or forbidden modes

$\Gamma(p\mu^+\mu^-)/\Gamma_{\text{total}}$

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{78}/Γ
<3.4 × 10⁻⁴	90	0	KODAMA	95	E653 π^- emulsion 600 GeV	

$\Gamma(\Sigma^-\mu^+\mu^+)/\Gamma_{\text{total}}$

A test of lepton-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{79}/Γ
<7.0 × 10⁻⁴	90	0	KODAMA	95	E653 π^- emulsion 600 GeV	

Λ_c^+ DECAY PARAMETERS

See the note on “Baryon Decay Parameters” in the neutron Listings.

α FOR $\Lambda_c^+ \rightarrow \Lambda\pi^+$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.91 ± 0.15 OUR AVERAGE				
-0.78 $\pm 0.16 \pm 0.19$		LINK	06A	FOCS γ A, $\bar{E}_\gamma \approx 180$ GeV
-0.94 $\pm 0.21 \pm 0.12$	414	22 BISHAI	95	CLE2 $e^+ e^- \approx \gamma(4S)$
-0.96 ± 0.42		ALBRECHT	92	ARG $e^+ e^- \approx 10.4$ GeV
-1.1 ± 0.4	86	AVERY	90B	CLEO $e^+ e^- \approx 10.6$ GeV

²² BISHAI 95 actually gives $\alpha = -0.94^{+0.21+0.12}_{-0.06-0.06}$, chopping the errors at the physical limit -1.0 . However, for $\alpha \approx -1.0$, some experiments should get unphysical values ($\alpha < -1.0$), and for averaging with other measurements such values (or errors that extend below -1.0) should *not* be chopped.

α FOR $\Lambda_c^+ \rightarrow \Sigma^+\pi^0$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$-0.45 \pm 0.31 \pm 0.06$	89	BISHAI	95	CLE2 $e^+ e^- \approx \gamma(4S)$

α FOR $\Lambda_c^+ \rightarrow \Lambda\ell^+\nu_\ell$

The experiments don't cover the complete (or same incomplete) $M(\Lambda\ell^+)$ range, but we average them together anyway.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.86 ± 0.04 OUR AVERAGE				
-0.86 $\pm 0.03 \pm 0.02$	3201	23 HINSON	05	CLEO $e^+ e^- \approx \gamma(4S)$
-0.91 $\pm 0.42 \pm 0.25$		24 ALBRECHT	94B	ARG $e^+ e^- \approx 10$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.82 $\pm 0.09 \pm 0.06$	700	25 CRAWFORD	95	CLE2 See HINSON 05
-0.89 $\pm 0.17 \pm 0.09$	350	26 BERGFELD	94	CLE2 See CRAWFORD 95

²³ HINSON 05 measures the form-factor ratio $R \equiv f_2/f_1$ for $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$ events to be $-0.31 \pm 0.05 \pm 0.04$ and the pole mass to be $2.21 \pm 0.08 \pm 0.14$ GeV/c², and from these calculates α , averaged over q^2 , where $\langle q^2 \rangle = 0.67$ (GeV/c)².

²⁴ ALBRECHT 94B uses Λe^+ and $\Lambda \mu^+$ events in the mass range $1.85 < M(\Lambda\ell^+) < 2.20$ GeV.

²⁵ CRAWFORD 95 measures the form-factor ratio $R \equiv f_2/f_1$ for $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$ events to be $-0.25 \pm 0.14 \pm 0.08$ and from this calculates α , averaged over q^2 , to be the above.

²⁶ BERGFELD 94 uses Λe^+ events.

$\Lambda_c^+, \bar{\Lambda}_c^-$ CP-VIOLATING DECAY ASYMMETRIES

$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda\pi^+$, $\bar{\Lambda}_c^- \rightarrow \bar{\Lambda}\pi^-$

This is zero if CP is conserved.

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.07 \pm 0.19 \pm 0.24$	LINK	06A	FOCS γ A, $\bar{E}_\gamma \approx 180$ GeV

$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} e^- \bar{\nu}_e$ This is zero if CP is conserved.

VALUE	DOCUMENT ID	TECN	COMMENT
0.00±0.03±0.02	HINSON	05	CLEO $e^+ e^- \approx \gamma(4S)$

 Λ_c^+ REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1992 edition (Physical Review **D45**, 1 June, Part II) or in earlier editions.

AUBERT	07U	PR D75 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
LINK	06A	PL B634 165	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
AUBERT,B	05S	PR D72 052006	B. Aubert <i>et al.</i>	(BABAR Collab.)
HINSON	05	PRL 94 191801	J.W. Hinson <i>et al.</i>	(CLEO Collab.)
LINK	05F	PL B624 22	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05K	PL B624 166	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
CRONIN-HEN...	03	PR D67 012001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
KAYIS-TOPAK..	03	PL B555 156	A. Kayis-Topaksu <i>et al.</i>	(CERN CHORUS Collab.)
ABE	02C	PL B524 33	K. Abe <i>et al.</i>	(KEK BELLE Collab.)
LINK	02C	PRL 88 161801	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02G	PL B540 25	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	
KUSHNIR...	01	PRL 86 5243	A. Kushnirenko <i>et al.</i>	(FNAL SELEX Collab.)
MAHMOOD	01	PRL 86 2232	A.H. Mahmood <i>et al.</i>	(CLEO Collab.)
AITALA	00	PL B471 449	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
JAFFE	00	PR D62 072005	D.E. Jaffe <i>et al.</i>	(CLEO Collab.)
ALAM	98	PR D57 4467	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	96E	PRPL 276 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEEV	96	JINRRC 3-77 31	A.N. Aleev <i>et al.</i>	(Serpukhov EXCHARM Collab.)
ALEXANDER	96C	PR D53 R1013	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ALBRECHT	95B	PL B342 397	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMMAR	95	PRL 74 3534	R. Ammar <i>et al.</i>	(CLEO Collab.)
BISHAI	95	PL B350 256	M. Bishai <i>et al.</i>	(CLEO Collab.)
CRAWFORD	95	PRL 75 624	G. Crawford <i>et al.</i>	(CLEO Collab.)
KODAMA	95	PL B345 85	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ALBRECHT	94B	PL B326 320	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEEV	94	PAN 57 1370	A.N. Aleev <i>et al.</i>	(Serpukhov BIS-2 Collab.)
		Translated from YF 57 1443.		
AVERY	94	PL B325 257	P. Avery <i>et al.</i>	(CLEO Collab.)
BERGFELD	94	PL B323 219	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
FRAEBETTI	94E	PL B328 193	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AVERY	93	PRL 71 2391	P. Avery <i>et al.</i>	(CLEO Collab.)
BOZEK	93	PL B312 247	A. Bozek <i>et al.</i>	(CERN NA32 Collab.)
FRAEBETTI	93D	PRL 70 1755	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRAEBETTI	93H	PL B314 477	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KUBOTA	93	PRL 71 3255	Y. Kubota <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92	PL B274 239	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92O	ZPHY C56 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARLAG	92	PL B283 465	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
CRAWFORD	92	PR D45 752	G. Crawford <i>et al.</i>	(CLEO Collab.)
JEZABEK	92	PL B286 175	M. Jezabek, K. Rybicki, R. Rylko	(CRAC)
ALBRECHT	91G	PL B269 234	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AVERY	91	PR D43 3599	P. Avery <i>et al.</i>	(CLEO Collab.)
ALVAREZ	90	ZPHY C47 539	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ALVAREZ	90B	PL B246 256	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ANJOS	90	PR D41 801	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
AVERY	90B	PRL 65 2842	P. Avery <i>et al.</i>	(CLEO Collab.)
BARLAG	90D	ZPHY C48 29	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
FRAEBETTI	90	PL B251 639	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
BARLAG	89	PL B218 374	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
AGUILAR-...	88B	ZPHY C40 321	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
Also		PL B189 254	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
Also		PL B199 462	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
Also		SJNP 48 833	M. Begalli <i>et al.</i>	(LEBC-EHS Collab.)
		Translated from YAF 48 1310.		

ALBRECHT	88C	PL B207 109	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	88B	PRL 60 1379	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ADAMOVICH	87	EPL 4 887	M.I. Adamovich <i>et al.</i>	(Photon Emulsion Collab.)
Also		SJNP 46 447	F. Viaggi <i>et al.</i>	(Photon Emulsion Collab.)
		Translated from YAF 46 799.		
AMENDOLIA	87	ZPHY C36 513	S.R. Amendolia <i>et al.</i>	(CERN NA1 Collab.)
JONES	87	ZPHY C36 593	G.T. Jones <i>et al.</i>	(CERN WA21 Collab.)
ABE	86	PR D33 1	K. Abe <i>et al.</i>	
ALEEV	84	ZPHY C23 333	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)
BOSETTI	82	PL 109B 234	P.C. Bosetti <i>et al.</i>	(AACH3, BONN, CERN+)
VELLA	82	PRL 48 1515	E. Vella <i>et al.</i>	(SLAC, LBL, UCB)
BASILE	81B	NC 62A 14	M. Basile <i>et al.</i>	(CERN, BGNA, PGIA, FRAS)
CALICCHIO	80	PL 93B 521	M. Calicchio <i>et al.</i>	(BARI, BIRM, BRUX+)

OTHER RELATED PAPERS

MIGLIOZZI	99	PL B462 217	P. Migliozzi <i>et al.</i>
DUNIETZ	98	PR D58 094010	I. Dunietz