



## Milk products in the dietary management of childhood undernutrition – a historical review

Veronika Scherbaum<sup>1\*</sup> and M. Leila Srour<sup>2</sup>

<sup>1</sup>*Institute for Biological Chemistry and Nutrition, and Food Security Centre, University of Hohenheim, Garbenstrasse 30, 70593 Stuttgart, Germany*

<sup>2</sup>*Health Frontiers, POB 2548, Vientiane, Laos*

### Abstract

The present narrative review outlines the use of milk products in infant and young child feeding from early history until today and illustrates how research findings and technical innovations contributed to the evolution of milk-based strategies to combat undernutrition in children below the age of 5 years. From the onset of social welfare initiatives, dairy products were provided by maternal and child health services to improve nutrition. During the last century, a number of aetiological theories on oedematous forms of undernutrition were developed and until the 1970s the dogma of protein deficiency was dominant. Thereafter, a multifactorial concept gained acceptance and protein quality was emphasised. During the last decades, research findings demonstrated that the inclusion of dairy products in the management of severe acute malnutrition is most effective. For children suffering from moderate acute malnutrition the evidence for the superiority of milk-based diets is less clear. There is an unmet need for evaluating locally produced milk-free alternatives at lower cost, especially in countries that rely on imported dairy products. New strategies for the dietary management of childhood undernutrition need to be developed on the basis of research findings, current child feeding practices, socio-cultural conditions and local resources. Exclusive and continued breast-feeding supported by community-based nutrition programmes using optimal combinations of locally available complementary foods should be compared with milk product-based interventions.

**Key words:** Undernutrition: Infants: Children: Breast-feeding: Dietary management: Dairy products: Ready-to-use therapeutic food

### Introduction

Food crises, nutritional deficiencies and associated illness affecting infants and young children are part of the history of mankind<sup>(1)</sup>. The first records of famines date back to ancient times<sup>(2–4)</sup>, and clinical signs of severe undernutrition are mentioned in the Old Testament<sup>(5,6)</sup>. However, systematic investigations in the field of undernourishment did not exist before the second half of the 19th century, a time when advances in epidemiology, statistics and anthropometry facilitated nutritional assessments<sup>(7,8)</sup>. Concurrently, infant and young child nutrition received more attention within welfare and public health programmes in Europe and the USA<sup>(9)</sup>. After the Second World War, nutrition interventions supported by UN organisations and international non-governmental organisations were increasingly implemented in developing countries, aiming to combat nutritional deficiencies, a major cause of under-five morbidity and mortality. In 2016, an estimated 155 million (22.9%) children under 5 years were stunted, and 52 million (7.7%) were wasted including 16.9 million (2.5%) with severe wasting. Over 41 million children (6%) globally were overweight or obese<sup>(10)</sup>.

In many programmes for the prevention and treatment of childhood undernutrition, dairy products constitute key

components. There is a growing body of evidence supporting the positive effects of milk proteins on the linear growth of healthy children and catch-up growth during recovery from undernutrition<sup>(11,12)</sup>. While potential mechanisms and responsible components of dairy products influencing body composition as well as weight and height gain were extensively studied during recent years, many questions remain unanswered<sup>(12)</sup>.

In the area of infant feeding and childhood undernutrition previous reviews have described developments chronologically<sup>(13–17)</sup>. Other historical reviews focused on breast-feeding, complementary feeding and various types of nutritional deficiencies<sup>(15–28)</sup>. Some were confined to distinct time periods or geographical contexts<sup>(21,29–37)</sup>. Similarly, a great number of reviews regarding milk product utilisation<sup>(38–43)</sup> addressed very specific aspects of prevention and treatment of childhood undernutrition.

The purpose of the present historical review is to address the following questions:

- (1) How did milk-based strategies to combat undernutrition in under-5-year-old children evolve over time?
- (2) What were the impacts of medical doctrines, research findings and technical innovations?

**Abbreviations:** LNS, lipid-based nutrient supplement; MAM, moderate acute malnutrition; RUSF, ready-to-use supplementary food; RUTF, ready-to-use therapeutic food; SAM, severe acute malnutrition.

\* **Corresponding author:** Veronika Scherbaum, email [veronika.scherbaum@uni-hohenheim.de](mailto:veronika.scherbaum@uni-hohenheim.de)

## Terms used in this review

The term 'milk/dairy products' refers to items produced from, or containing, milk of mammals, primarily cattle. These products include infant formula, yoghurt, cheese, condensed milk, skimmed milk, whey protein concentrate, milk-based therapeutic foods, etc., whereas the term 'milk' usually refers to bovine milk.

While the term 'malnutrition' includes forms of overnutrition (i.e. overweight and obesity), the present review focuses on states of undernutrition in children under 5 years, including underweight, wasting, stunting, micronutrient deficiencies and low birth weight<sup>(44)</sup>. In this article, the term 'undernutrition' is used, unless other terms such as kwashiorkor, marasmus, protein–energy malnutrition, severe acute malnutrition (SAM) and moderate acute malnutrition (MAM) were used in publications of that time.

In Table 1<sup>(17,23,25,45–69)</sup>, medical terms used for specific forms of undernutrition during various time periods are summarised.

## Use of milk products in infant and young child feeding until the 20th century

Findings of fatty residues in Neolithic feeding vessels in Europe suggest that products from animal milk were introduced for

feeding young children more than 7000 years ago<sup>(70)</sup>. It is assumed that from very early time preservation of milk was achieved through heating, fermentation and the manufacture of yoghurt, cheese and butter<sup>(38,71–73)</sup>. In ancient civilisations of the Mediterranean region, the Middle East and India breast-feeding was viewed as essential to preserve life and was an obligation of mothers<sup>(15,28)</sup>. When breast milk could not be provided by mothers or wet nurses, animal milk was offered to young children, sometimes directly from the udder of animals<sup>(16)</sup>. Milk was seen as more than solely a source of nutrition. It was often regarded as a heavenly elixir reflecting fertility and the nurturing mother–infant relationship while its whiteness was a symbol of goodness and purity<sup>(38,40)</sup>. Since antiquity, differences in the composition of milk between mammalian species were described and certain qualities were attributed to particular types of animal milk. For example, bovine milk was used as a remedy against specific illnesses during the Roman Empire<sup>(40)</sup>. Concerning infant feeding, animal milk has been used for centuries as the main component of artificial foods including as a substitute for colostrum after delivery<sup>(16)</sup>.

In 1610, O. Gaebelkhovern highlighted that children fed with diluted cows' milk combined with cereal preparations thrived better than those fed solely with unmodified cows' milk<sup>(15,74)</sup>. This observation was endorsed in 1838 by J. F. Simon after his discovery that cows' milk contains more protein and less carbohydrate than human milk<sup>(75,76)</sup>. His findings led to a variety of mixtures with cereal preparations, diluted cows' milk<sup>(15,77)</sup> and often enrichments with sugar and cream<sup>(78)</sup>. In 1884, P. Biedert in Germany and C. D. Meigs in the USA made a precise comparison of the nutrient contents in cows' milk and human breast milk. Based on their metabolic studies in the 1890s, Otto Heubner and M. Rubner calculated the daily energy requirements, using calorimetric methods, for healthy and undernourished infants and young children<sup>(79)</sup>. At the same time, T. M. Rotch published a method for calculating the precise proportions of carbohydrates, proteins and fats required to substitute diluted cows' milk for human milk<sup>(80)</sup>. This so-called 'percentage method' as well as the 'calorie method' introduced by H. Finkelstein were too complicated for the production of artificial formula at the household level<sup>(40)</sup>. On the basis of these research findings, the formula industry used the opportunity to develop artificial infant formulas<sup>(80–82)</sup>.

For centuries, contamination of milk during production, transport and dilution with polluted water has significantly contributed to the high morbidity and mortality of artificially fed infants<sup>(83)</sup>. This feeding practice was particularly common among mothers of lower classes employed in factories, who were forced to wean their babies shortly after delivery<sup>(80,84)</sup>. Besides general improvements in hygiene, by the middle of the 19th century important innovations took place that reduced the risk of milk-borne diseases in Europe and the USA. First, the invention of evaporation enabled the production of condensed milk. Second were the discovery and utilisation of pasteurisation techniques<sup>(21)</sup>. The claims of 'clean milk movements' led to the supply of, often subsidised, 'clean' milk in special dispensaries, milk depots and infant welfare centres. Mandatory pasteurisation laws were adopted in many countries<sup>(85–87)</sup>.

**Table 1.** Terms used for various forms of undernutrition/malnutrition in history\*

First known time of citation	Terms	Reference
About 1300	Wasting†	(46)
1538	Cachexia	(46)
1826	Malnutrition†	(46)
1876	Undernutrition†	(46)
Since 1880	Atrophy, marasmus†	(17,47)
1890s–1930s	Hunger oedema‡, childhood oedema‡ Different regional terms: <i>culebrilla</i> , <i>boufissure d'Annam</i> , <i>syndrome dépigmentation-oedème</i> , <i>syndrome policarencial infantil</i> ‡	(48–50)
1906–1928	<i>Mehlnährschäden</i> ‡	(51)
Since 1935	Kwashiorkor‡	(52)
1940	Infantile pellagra‡	(53)
1944	Malignant malnutrition‡	(54)
1948	Fatty liver disease‡	(55)
1952	Protein malnutrition‡	(56)
1959	Protein–calorie malnutrition‡	(57)
From 1950s–1960s	Low birth weight† Intra-uterine/fetal malnutrition† Growth retardation†/growth restriction†	(58,59) (60,61) (62)
	Small (full) term, small for dates, small for gestational age†	(63,64)
Since 1970	Protein–energy malnutrition‡	(65)
1980–1990	Energy nutrient malnutrition	(23)
1994	Micronutrient malnutrition†, hidden hunger†	(25,66,67)
Since 1999	Oedematous malnutrition‡	(68)
Since 2006	Severe acute malnutrition‡, moderate acute malnutrition†	(69)
Since 2008	Malnutrition† (in all its forms) includes underweight, wasting, stunting, micronutrient deficiencies, low birth weight, overweight and obesity	(44)

\* Modified from Scherbaum<sup>(45)</sup>.

† Term still used today.

‡ Oedematous forms of undernutrition.

The introduction of milk pasteurisation was a milestone in the history of public health. Like other improvements in infection protection, these measures were built on study results in the field of infection epidemiology by J. Snow<sup>(88,89)</sup> and findings in bacteriological research by R. Koch and L. Pasteur in the middle of the 19th century<sup>(86,87)</sup>. However, these advances facilitated the spread of the 'germ theory' which influenced medical thinking that the causes of diseases can be ascribed mainly to microbes<sup>(90)</sup>. Moreover, the 'germ theory' when applied to diseases like beriberi<sup>(90,91)</sup> was a general barrier to the recognition of deficiency diseases<sup>(92)</sup>. Only in the second decade of the 20th century did evidence of specific micronutrient deficiencies lead to the aetiological concept of 'deficiencies'<sup>(93)</sup>.

At the end of the 19th century, it was recognised that young children who were fed with unfortified pasteurised milk, condensed milk or industrially produced infant formula<sup>(94)</sup> were developing infantile scurvy. Based on the 'germ theory', the aetiology of chronic poisoning by absorption of ptomaine toxin, a waste product of bacteria, was suggested. In 1914, the paediatrician A. Hess proved by experiments that through pasteurisation antiscorbutic properties of milk are destroyed, which could be prevented by supplementing fresh fruit or vegetable juices when infants received these formulas<sup>(94)</sup>.

#### Use of milk products in nutrition programmes in the first half of the 20th century

At the beginning of the 20th century, the German paediatricians A. Czerny and A. Keller suggested that overfeeding with cows' milk to young children leads to '*Milchnährschäden*' with symptoms of dyspepsia and failure to thrive. Similarly, a monotonous diet containing mainly cereal flour was suspected to be the main cause of '*Mehlnährschäden*' characterised by undernutrition with oedema, thought to be secondary to protein deficiency<sup>(51,95,96)</sup>. At the same time, the concept of an alimentary toxicosis was proposed by H. Finkelstein. He suspected that the degradation of certain alimentary substrates, through fermentation of carbohydrates, produced toxins in the immature gut of infants leading to food intolerance with diarrhoea and weight faltering<sup>(97)</sup>. The presumed aetiology of the unwholesome effects of an inappropriate composition of children's diets spread worldwide. Attention was distracted from effective measures against essential causes of infant gastroenteritis, namely fundamental improvements in hygiene and promotion of breast-feeding. During the following decades, investigations failed to detect any of the postulated toxins. However, a variety of therapeutic milk-based preparations was developed, for example Finkelstein's 'protein milk' or modified buttermilk<sup>(98)</sup>.

In countries, such as England, cows' milk played a particular role in charitable feeding at the beginning of the 20th century and was most commonly provided to debilitated and severely undernourished children, often together with cod liver oil<sup>(99)</sup>. In the 1920s, when a growing number of 'accessory factors' (vitamins) were discovered, milk was increasingly considered as a 'complete food' and a special nurturing medium for young children<sup>(100,101)</sup>. During that time, intervention studies in the USA and Britain revealed a positive effect of supplementary milk feeding on the nutritional status of school-age

children<sup>(102–105)</sup>. These study results contributed to the expansion of supplementary milk feeding programmes in Britain during the 1930s. Due to major methodological constraints, the results of these studies were questioned by some authors and the influence of the dairy industry on nutrition policies was critically debated<sup>(100,106,107)</sup>.

The high prevalence of child undernutrition between both World Wars led to relief programmes in Austria, Germany, Poland, Russia and other countries, delivered by organisations including Save the Children Fund and the support of philanthropists<sup>(108,109)</sup>. In 1922, Russia, food aid containing milk-based foods, such as mixtures of canned milk with maize, sugar and fats, was implemented by the American Relief Administration for Russian children<sup>(110)</sup>. In Germany, the '*Moro Brei*', a gruel made of whole milk, butter, flour and sugar was frequently offered to undernourished children. To safeguard the quality of these preparations under adequate hygienic conditions, special milk kitchens were established in hospitals<sup>(98)</sup>.

#### Nutritional interventions and protein-role controversies

Experiences in Europe and the USA influenced interventions promoted in overseas territories. Tinned condensed milk was used by the paediatrician Cicely Williams in combination with malt and cod liver oil to treat children suffering from kwashiorkor in the former British colonies on the 'Gold Coast' of West Africa in the 1930s<sup>(52,111)</sup>. Based on her observations of affected children who were fed a monotonous maize diet, she presumed that protein deficiency was the main cause of oedematous forms of undernutrition<sup>(111)</sup>. Her positive view regarding the use of condensed milk in infant feeding rapidly changed after she was transferred to Malaya<sup>(112)</sup> where this product was being used by mothers as a breast milk substitute. As early as 1880 sweetened condensed milk was advertised as '...the food par excellence for delicate infants'<sup>(113)</sup> and was advocated by colonial doctors. Many mothers were convinced by female milk industry employees, dressed as nurses, that this milk product was the best replacement for their own breast milk. In 1939, in her famous speech entitled 'Milk and Murder' held in the Rotary Club in Singapore, Williams named and shamed this practice and its consequences, manifesting in diarrhoea, marasmus and death<sup>(114)</sup>.

During the following decades undernutrition remained a major public health problem in many parts of the world. While protein deficiency was widely regarded as a major cause of oedematous forms of undernutrition, milk was seen as the best source of protein<sup>(56,115)</sup>. Since animal protein was expensive shortly after the Second World War, trials were conducted using plant proteins such as soya, bananas, plantains and maize flour<sup>(116,117)</sup>. However, milk-based diets showed the best results, particularly for treating severely undernourished children<sup>(117–120)</sup>. From the 1950s onward, supplementation with skimmed milk was increasingly practised in nutrition programmes delivered by UNICEF aiming to control 'protein malnutrition'. However, milk powder surpluses decreased in industrialised countries, and achievement of significant increases in local milk production in many overseas regions appeared to be unrealistic<sup>(56,121)</sup>. In 1955, a Protein Advisory

Group was initiated by the WHO and efforts were made to investigate alternative non-milk foods to combat a supposed worldwide 'protein gap'<sup>(122)</sup>. These efforts included the production of 'protein rich food mixtures' based on fish, flour, soya, cottonseed, groundnuts, sesame or coconuts. However, many of these innovative approaches were halted due to difficulties in food technology, food safety considerations and high costs that made these products unaffordable for poor populations<sup>(121)</sup>.

Until the foundation of the World Food Programme in the early 1960s, surpluses primarily of cereals from industrialised countries were distributed to countries suffering from humanitarian crises. Subsequently, fortified blended foods, consisting of maize or wheat, vegetable oil and sugar, were introduced in supplementary feeding programmes. Skimmed milk and soya flour were added as protein sources reflecting the 'protein-calorie deficiency' theory. When US milk surpluses were exhausted and evidence revealed ineffective and unsafe use of milk powder in community feeding programmes, this food aid compound was gradually replaced by maize-soya or wheat-soya blends<sup>(123,124)</sup>. By the 1980s, the daily protein requirements for children were gradually reduced<sup>(125–127)</sup>, and the dogma of protein deficiency was questioned and refuted by research<sup>(128–130)</sup>. Early observations revealed that undernourished children often received too few meals a day and primarily bulky foods with low energy density<sup>(131,132)</sup>. The energy content of complementary foods was readdressed. With respect to oedematous forms of undernutrition, the aetiological concept shifted from 'protein malnutrition' in the 1950s<sup>(56)</sup> to 'protein-calorie malnutrition' in 1959<sup>(57)</sup> to 'protein-energy malnutrition' in the 1970s<sup>(65)</sup>. In addition, the theory of a multifactorial aetiology developed including the potential impact of infections, aflatoxins as well as micronutrient deficiencies, free radicals and most recently alterations of the intestinal microbiome<sup>(128,133–136)</sup>. In the mid-1970s, the focus on a worldwide 'protein gap' faded, but during the next decades, expert committees of the FAO and WHO continued to address protein and amino acid requirements in human nutrition.

Increasing awareness and the 'rethinking protein' with respect to childhood undernutrition are a new development<sup>(126,127,137–140)</sup>. A large proportion of children in low-income countries depend on low-protein diets. These children are frequently affected by chronic energy deficits, repeated infections and stunting, so research has explored the role of protein quality determined by its digestibility and bioavailability of essential amino acids<sup>(11,138,141,142)</sup>. A recent study in Malawi demonstrated a correlation between a high prevalence of stunting and reduced levels of circulating essential amino acids among children below 5 years of age<sup>(137,142–145)</sup>.

### Development of milk products used for dietary interventions

Milk has long been recognised as a well-balanced source of energy with numerous essential nutrients<sup>(12,146)</sup> playing a key role in treating childhood undernutrition both in industrialised and in developing countries<sup>(147)</sup>.

Milk is known to contain high-quality protein with all essential amino acids including lysine which is often deficient in

traditional cereal-based diets of agriculturist populations<sup>(42,148)</sup>. The two main fractions of milk are the water-soluble 'whey' protein and insoluble 'casein' protein. Various authors consider milk to be the best protein source according to the essential and protein-digestibility amino acid scores<sup>(149–151)</sup>. Milk protein positively affects linear growth in healthy children and there is growing evidence of similar effects on recovery from childhood undernutrition<sup>(12,152)</sup>. The beneficial effects of milk protein, commonly in combination with micronutrient supplementation, were demonstrated in a dose-response relationship on catch-up height and weight gain<sup>(153,154)</sup>. Moreover, high-quality milk proteins contribute to effective immune functions by increasing acute-phase protein synthesis in response to infections which often accompany SAM<sup>(11,12,141)</sup>.

The specific effectiveness of high-quality whey protein in the treatment of moderately wasted children (age 6–59 months) was highlighted in a recent intervention study in Malawi and Mozambique. Supplementation with whey-based products resulted in better recovery and growth rates than did supplementation with products based on soya, even though the whey-based supplement provided 33% less total protein and 8% less energy than the soya-based product<sup>(155)</sup>.

Regarding the carbohydrate content of milk, the disaccharide lactose is known to enhance Ca absorption and, like specific oligosaccharides released from milk glycoproteins, induces prebiotic effects on the gut microbiome contributing to enhanced efficiency of food utilisation in the intestine<sup>(12,146,156,157)</sup>.

Milk supplies key micronutrients like Ca, P, Se, Mg, Zn and vitamins A, D, E and B, without the antinutrients, such as phytates and oxalates<sup>(151)</sup>. In addition, milk contains bioactive compounds exhibiting a wide variety of physiological functionalities, including mineral transport and growth-promoting activities<sup>(158)</sup>.

Since 1970, in very severe cases of undernutrition, milk was seen as an excellent vehicle for micronutrient fortification and was valued for its liquid form, enabling nasogastric tube feeding<sup>(159–161)</sup>. On the basis of clinical experience, past research findings and new knowledge about the role of micronutrients, specific therapeutic milk formulas (F-75, F-100) were created. These contain relatively low concentrations of protein and a mixture of specific micronutrients<sup>(162)</sup>. Treatment regimens using these milk formulas for the management of severe undernutrition<sup>(161,163)</sup> were developed and published by the WHO in 1999<sup>(68)</sup>.

During the same year, a paste made of groundnut butter, milk powder, vegetable oil and sugar, fortified with the same mix of micronutrients, was tested in a pilot study of marasmic children treated in a therapeutic feeding centre in Chad<sup>(164)</sup>. The effectiveness of this so-called 'ready-to-use therapeutic food' (RUTF) was demonstrated in Sub-Saharan Africa<sup>(165,166)</sup>. The milk-powder in F-100 was partially replaced by groundnut paste and changed from liquid to a spread, making feasible the community management of children with SAM<sup>(165,167)</sup>, accepted internationally in 2007<sup>(166)</sup>. RUTF do not require cooking and their low moisture content reduces the risk of bacteria and mould growth, allowing for a long shelf life, even without refrigeration.

According to the WHO, there is sufficient evidence of the efficacy of milk products in the dietary management of SAM<sup>(68,167)</sup>. This applies for F-75 and F-100 therapeutic milk



formula for the hospital-based treatment of cases with complications and RUTF for the community-based management of SAM without complications<sup>(168)</sup>.

### Growing commercialisation of ready-to-use foods

The commercial marketing of costly RUTF, patented and primarily produced in industrialised countries, made its usefulness debatable<sup>(169,170)</sup>. This stimulated the development of locally produced therapeutic food, acceptable and cost-effective in most trials<sup>(171–173)</sup>. During the last decade, research shifted to the management of MAM, to prevent the development of severe forms, which is more expensive and management time consuming<sup>(174)</sup>.

Consequently, improved fortified blended foods were developed to provide the energy and nutrient requirements of infants and young children during disasters<sup>(124)</sup>. Ready-to-use supplementary foods (RUSF), with a higher energy density and adjusted micronutrient compositions, were developed for moderately wasted children, as well as pregnant and lactating women<sup>(124,175–177)</sup>. The inclusion of milk protein in RUSF appears beneficial in children recovering from MAM<sup>(155)</sup>. Two recent intervention studies in Guinea-Bissau demonstrated that RUSF with a higher dairy protein content (33%) were superior to RUSF with a lower content (15%) in the community-based management of undernourished preschool children and mothers<sup>(154,178)</sup>.

In 2009, the International Lipid-Based Nutrient Supplements project aimed to develop lipid-based nutrient supplements (LNS) for the prevention of undernutrition in food-insecure settings<sup>(175)</sup>. LNS are special types of RUSF with varying energy densities and micronutrient concentrations, with or without small amounts of milk products. As these industrially manufactured items need to be offered only in small doses, they were added to general food rations for at-risk populations<sup>(175,179)</sup>.

### Strategies to reduce the costs of milk-based ready-to-use foods

The major constraint of using milk-based ready-to-use foods is the high cost. With respect to standard RUTF, of the total expenses per child cured from SAM (US\$ 70–200), about half is spent for the therapeutic product alone<sup>(170,180)</sup>. More than half the cost of the therapeutic formulation is due to the milk powder<sup>(174,181)</sup>.

An attempt to reduce the milk powder content by 25% in standard RUTF by replacing 15% with soya showed that the product with only 10% milk powder was clinically less effective in the treatment of SAM, in weight gain and recovery rates. While both formulations had nearly identical nutrient contents, the content of milk protein, of antinutrients or the impact of unidentified beneficial factors associated with milk supplementation might explain the difference in effectiveness<sup>(153)</sup>.

To reduce the costs of milk-based ready-to-use foods, projects were implemented aiming to replace the skimmed milk content with whey protein concentrate. Whey protein concentrate (34%) is about 25–33% cheaper and can be generated as a surplus product from cheese manufacturing<sup>(148)</sup>. Recent interventions with products based on whey containing high-quality protein, high levels of lactose, micronutrients and

bioactive factors have yielded promising results in the dietary management of MAM and SAM<sup>(182–184)</sup>.

As many countries in Africa and Asia largely rely on imported dairy products, efforts were made during the last 10 years to develop suitable milk-free alternatives based on locally available foods at relatively low cost<sup>(185,186)</sup>. These ready-to-use-food formulations commonly contain cereal flours, pulses, nuts and/or seeds, and vegetable oil and sugar which are often supplemented with a mineral–vitamin premix. While the acceptability of these products has been reported to be generally good<sup>(187,188)</sup>, concern has been raised about the higher content of fibres and antinutrients such as phytates in milk-free formulations, which can impair the bioavailability of micronutrients including Fe and Zn<sup>(189)</sup>. Processing such as dehiscing, soaking, roasting, malting, germination and fermentation have been used to lower the anti-nutrient content<sup>(190)</sup>. Apart from the cost and acceptability considerations of alternative formulations, the optimisation of protein quality is particularly important in the development of milk-free products. Linear programming can contribute to choosing the appropriate ingredients such as foods of animal origin or best combinations of locally available plant proteins<sup>(191)</sup>. The palatability and taste of the improved recipes must be evaluated for acceptance by the target groups<sup>(192)</sup>.

During the last decade, research teams particularly in countries of Sub-Saharan Africa evaluated the efficacy of therapeutic formulations without dairy ingredients compared with milk-based products, including standard RUTF and preparations containing whey. Regarding a formulation containing whey protein concentrate, an equally effective alternative to standard RUTF was demonstrated in the treatment of SAM with lower costs<sup>(184)</sup>.

Compared with the efficacy of milk-based products, certain milk-free formulations have been shown to be equivalent in the management of MAM<sup>(183,192–194)</sup>, whereas in other studies, these formulations were less effective regarding the treatment of MAM<sup>(155,195–200)</sup> as well as SAM<sup>(201)</sup>. While a reduced efficacy of preparations without milk was generally most pronounced in children below the age of 2 years, it has been suggested that milk-free products should be used preferentially in the treatment of undernourished children older than 2 years, whereas the younger age group may depend more on products containing dairy products, especially if breast-feeding has been terminated<sup>(201,202)</sup>. Very recently, a study in Malawi showed that a milk-free formulation containing soya, maize and sorghum, enriched with crystalline amino acids, was as efficacious as standard RUTF with respect to recovery rates of SAM children aged 6–23 months and 24–59 months. Moreover, this milk-free formulation was even better at correcting Fe-deficiency anaemia<sup>(203)</sup>.

The costs of dietary regimens currently used in the treatment of MAM were recently summarised by Suri *et al.*<sup>(39)</sup>. There remains a need to evaluate the cost-effectiveness of treatment by costs per impact or effect<sup>(204)</sup>. The required time period of treatment and nutritional advantages of certain ingredients, the palatability of products and acceptability of interventions and the compliance of the target group should be considered in future programmes<sup>(192,205)</sup>.

### Limitations of current studies, knowledge gaps and research needs

Appraisals of the beneficial role of milk supplementation are often compromised by failing to meet internationally agreed criteria of study design and reporting<sup>(206,207)</sup> and lack relevant study details. Precise information on the amount of dairy products, other sources of protein, the amount of supplementary food offered and actually consumed by the target group, variations in compliance, and the potential impact of educational interventions are lacking<sup>(205,208)</sup>. Similarly, the validity of results from clinical trials could be enhanced by comparing isoenergetic and isoenergetic plus isonitrogenous dietary conditions, but this has been accomplished in very few studies<sup>(209)</sup>.

Additional information is required about the intensity and long-term duration of breast-feeding, the quality of complementary foods as well as family foods including seasonal nutritional insecurities<sup>(210)</sup>.

There is still no scientific evidence about the minimum amount of milk protein required to exert an adequate effect on weight gain and growth among children of different age groups<sup>(68)</sup>, while the impact of other sources of protein and other beneficial compounds needs to be considered<sup>(39)</sup>.

As most estimations of protein, amino acid and other nutrients are based on measurements among healthy individuals, research on nutrient requirements for undernourished children is needed to improve the composition of therapeutic diets to achieve adequate catch-up growth during different stages of treatment, rehabilitation and to minimise potential adverse effects in later life<sup>(205,211,212)</sup>.

Regarding the effectiveness of supplementary feeding programmes, there is still no conclusive evidence with respect to the potential of LNS to achieve adequate weight gain<sup>(213)</sup> and certain developmental outcomes, including the prevention of growth faltering of children<sup>(214–217)</sup>. A Cochrane analysis in 2013 showed no proven benefits of LNS compared with other blended and less costly foods such as fortified maize–soya blend<sup>(218)</sup>.

### Challenges regarding milk-based dietary interventions

The relatively low Fe content and bioavailability of Fe in bovine milk is a particular drawback in offering cows' milk to infants during the complementary feeding period<sup>(219,220)</sup>. In 1992, the American Academy of Pediatrics recommended avoiding whole cows' milk before 1 year of age due to its high renal solute content, which places small children at increased risk of dehydration under conditions of water stress. In addition, the risk of occult intestinal bleeding, which can be prevented by using heat-treated cows' milk<sup>(221,222)</sup>, was taken into consideration<sup>(101,223–226)</sup>. Regarding supplementation of Fe, due to its critical role in catalysing free radical oxidation and susceptibility to infectious diseases, it is generally not recommended to give Fe during the initial stabilisation phase of SAM<sup>(68,163)</sup>. Children treated with nutrient-dense RUTF need to drink sufficient extra water, challenging in many settings with inadequate safe water supplies<sup>(170)</sup>.

Clearly, conditions of undernutrition in the first 2 years of life, such as stunting, severe wasting and intra-uterine growth restriction, are known to cause considerable harm regarding the

health and development of the child<sup>(227)</sup>. However, there are concerns that excessive weight gain during and after rehabilitation may be associated with adverse effects on long-term health<sup>(212,228,229)</sup>. These concerns are based on results from observational studies suggesting that accelerated weight gain and growth in early life enhance the risk for developing obesity and cardiometabolic diseases<sup>(212,230–233)</sup>. As increased growth velocity has been observed among infants who were never or only briefly breast-fed, this effect was explained by the higher milk protein content of artificial infant formulas compared with human breast milk<sup>(234–237)</sup>.

While there is some controversy concerning obesogenic influences of high intake of milk protein in early life<sup>(238)</sup>, critical protein levels in different age groups and contexts have yet to be established<sup>(212)</sup> and pathogenic mechanisms like the suggested induction of insulin and insulin-like growth factor-1 are still hypothetical<sup>(239–241)</sup>.

Apart from the need to support long-term protective effects of breast-feeding against obesity, various gaps in research on the 'growth acceleration hypothesis' must be addressed<sup>(232)</sup>. The short-term benefits of dietary interventions resulting in rapid catch-up weight and growth should be counterbalanced with potential risks of non-communicable diseases in later life<sup>(212,242)</sup>. There is some evidence that the long-term effects of whey protein and casein on linear growth are similar; in contrast, whey protein induces less weight gain as compared with casein<sup>(243–245)</sup>. Consequently, long-term obesogenic complications of catch-up growth may be diminished by the use of whey-based diets<sup>(12,152)</sup>.

The implementation of milk-based interventions in regions with low milk consumption is problematic. Areas with a higher prevalence of lactase deficiency may limit caregiver acceptance of milk in young child feeding<sup>(192)</sup>. As lactase deficiency usually manifests itself among children older than 5 years<sup>(246)</sup>, the American Academy of Pediatrics stated in 1978 that '...it would be inappropriate to discourage supplemental milk feeding programs targeted at children on the basis of primary lactose intolerance'<sup>(247)</sup>. Milk-based diets have not been shown to cause major clinical concern, neither in young children with diarrhoea nor in the treatment of child undernutrition<sup>(248–251)</sup>. In fact, there is evidence that the prebiotic effects of lactose facilitate the absorption of minerals<sup>(156,252)</sup>. It has been suggested that undernourished children with secondary lactase deficiency caused by environmental enteropathy and diarrhoeal diseases might benefit from products with reduced lactose content<sup>(156)</sup>. A meta-analysis of studies in low- and middle-income countries among children with acute diarrhoea demonstrated that liquid feeds with reduced lactose content such as yoghurt were not superior to those containing lactose, whereas liquid lactose-free diets reduced both the duration and risk of treatment failure<sup>(253)</sup>.

For many years the distribution of powdered milk in emergency situations has been a major challenge, particularly when the product has to be reconstituted with unsafe water<sup>(254–256)</sup> or when milk powder is contaminated with *Enterobacter sakazakii*<sup>(257)</sup>. The policy was adopted that humanitarian organisations should not distribute milk powder as take-home rations<sup>(258)</sup>. Similarly, it has been recognised that the high renal solute load of drinks

prepared from skimmed milk powder can be particularly harmful for undernourished infants whose maximum renal concentrating capacity is significantly reduced<sup>(226,256,259)</sup>.

The current focus on 'product-based' approaches to combating childhood undernutrition is seen critically by a variety of authors who point out that ready-to-use foods tend to consume most of the financial resources and are detracting investments from long-term and sustainable programmes<sup>(170,260)</sup>. Comprehensive interventions are needed to address the multifactorial causes of childhood undernutrition, already outlined 30 years ago<sup>(135,169,261,262)</sup>. Due to the specific requirements of undernourished children of different age groups living in various settings, appropriate dietary management cannot be met with industrial products designed in the sense of 'one size that fits all'<sup>(172,263)</sup>.

Feeding commercial foods aiming to prevent MAM during a particularly sensitive time period of early child development might shape the taste preferences of young children towards food items other than those locally available<sup>(170,264)</sup>. Although recent studies did not confirm a decrease in breast-feeding rates and intake of commonly consumed foods during home fortification of complementary foods with LNS<sup>(265–268)</sup>, a negative impact under non-research conditions and uncontrolled promotion of products can never be ruled out<sup>(170)</sup>.

Although breast-feeding offers children and mothers unrivalled health benefits, worldwide deficits in breast-feeding promotion and support are a 'missed opportunity for global health'<sup>(269)</sup>. Aggressive marketing by the formula industry and violations of The International Code of Marketing of Breast-milk Substitutes undermine breast-feeding promotion and contribute to the decline of breast-feeding rates<sup>(270)</sup> with immense health consequences, named 'commerciogenic malnutrition' by the paediatrician D. Jelliffe in 1972<sup>(271)</sup>. A recent study in Laos has shown that parents misperceive coffee creamer products as suitable for infant feeding and use them as breast milk substitutes<sup>(272)</sup>.

## Conclusion

The present historical review describes the changing role and development of milk products in infant and young child feeding and in the dietary management of childhood undernutrition. From the ancient perceptions of milk as a divine living fluid, dairy products have become agro-industrial food items providing key nutrients required for the growth and development of children. Before the availability of pasteurised milk, artificial infant feeding carried the risk of milk-borne diseases and life-threatening consequences on child health. General improvements in hygiene and technical innovations in the middle of the 19th century have facilitated the industrial production of breast milk substitutes.

The historical perspective reveals that dietary interventions were often dominated by unchallenged doctrines and many decades passed before research findings could reject some of them. This applies to the aetiological concept of alimentary toxins contributing to wasting/marasmus, while the 'germ theory' delayed the recognition of diseases caused by nutritional deficiencies. Similarly, the assumption that protein deficiency is the main cause of severe malnutrition contributed

to an increased protein content in therapeutic feeding regimens. However, study results in the 1960s and 1970s challenged this 'protein dogma', the focus of research shifted to the role of energy and micronutrient deficiencies and with respect to protein from quantitative to qualitative considerations.

Regarding dietary therapies of children affected by SAM, research supports improved efficacy with inclusion of milk powder. This has led to the development of RUTF allowing outpatient management of severely undernourished children without medical complications. However, the following strategy of using imported expensive milk products to treat moderate forms of MAM is critically viewed as a supply- rather than a demand-driven approach. Studies revealed the community production of ready-to-use foods with locally available food leads to better acceptance, lower costs and offers opportunities for local employment.

Current nutritional interventions favour the inclusion of milk products due to their high density and bioavailability of nutrients such as high-quality protein and key micronutrients. Regarding negative long-term consequences of rapid catch-up weight associated with the use of cows' milk products there is some evidence that these risks may be minimised by using whey protein products.

However, many challenges and research gaps remain in the use of milk-based products applied in the dietary management of childhood undernutrition. Comprehensive interventions are needed to address the multifactorial causes of undernutrition and the specific requirements of children of different age groups living in various settings. Finally, the high costs of commercial milk products need to be compared with equally funded community-based child care and nutrition programmes supporting and promoting breast-feeding and healthy complementary foods that are locally available.

## Acknowledgements

There are no conflicts of interest.

## References

1. Cohen MN (1995) Prehistoric patterns of hunger. In *Hunger in History: Food Shortage, Poverty, and Development*, pp. 52–97 [LF Newman, editor]. Oxford: Blackwell.
2. Dando WA (1983) Biblical famines, 1850 B.C.–A.D. 46: insights for modern mankind. *Ecol Food Nutr* **13**, 231–249.
3. Murton B (2000) Famine. In *The Cambridge World History of Food*, vol. 2, pp. 1411–1426 [KFE Kiple and KC Ornelas, editors]. Cambridge: Cambridge University Press.
4. Gill RB (editor) (2000) *The Great Maya Droughts: Water, Life, and Death*. Albuquerque, NM: University of New Mexico Press.
5. BibelHub (2017) Nehemiah 9:21. <http://biblehub.com/nehemiah/9-21.htm>
6. BibelHub (2017) Lamentations 5:10. <http://biblehub.com/lamentations/5-10.htm>
7. Komlos J & Meermann L (2004) *The Introduction of Anthropometrics into Development and Labor Economics. Discussion Papers in Economics 381*. Munich: University of Munich, Department of Economics.
8. Rotberg RI (1983) Nutrition in history. *J Interdiscipl Hist* **14**, 199–204.



9. McGill N (1921) Infant welfare work in Europe. An account of recent experience in Great Britain, Austria, Belgium, France, Germany, Italy. In *Community Child-Welfares Series*, vol. 76. Washington, DC: Government Printing Office, Bureau Publication.
10. United Nations Children's Fund, World Health Organization & World Bank Group (2017) Levels and Trends in Child Malnutrition. Joint Child Malnutrition Estimates. Key Findings of the 2017 Edition. [http://www.who.int/nutgrowthdb/jme\\_brochure2017.pdf](http://www.who.int/nutgrowthdb/jme_brochure2017.pdf)
11. Manary M, Callaghan M, Singh L, *et al.* (2016) Protein quality and growth in malnourished children. *Food Nutr Bull* **37**, S29–S36.
12. Yackobovitch-Gavan M, Phillip M & Gat-Yablonski G (2017) How milk and its proteins affect growth, bone health, and weight. *Horm Res Paediatr* **88**, 63–69.
13. Quandt SA (2000) Infant and child nutrition. In *The Cambridge World History of Food*, vol. 2, pp. 1444–1453 [KF Kiple and KC Ornelas, editors]. Cambridge: Cambridge University Press.
14. Wickes IG (1953) A history of infant feeding. I. Primitive peoples; ancient works; Renaissance writers. *Arch Dis Child* **28**, 151–158.
15. Fildes VA (1986) *Breasts, Bottles and Babies: A History of Infant Feeding*. Edinburgh: Edinburgh University Press.
16. Castilho SD & Barros Filho AA (2010) The history of infant nutrition. *J Pediatr (Rio J)* **86**, 179–188.
17. Hansen JDL (2000) Protein–energy malnutrition. In *The Cambridge World History of Food*, pp. 977–988 [KFE Kiple and KC Ornelas, editors]. Cambridge: Cambridge University Press.
18. Tönz O (2003) Stillpraxis im Wandel der Zeit (Breastfeeding practices over the course of time). In *Stillen Frühkindliche Ernährung und reproduktive Gesundheit*, pp. 1–6 [V Scherbaum, FM Perl and U Kretschmer, editors]. Cologne: Deutscher Ärzte-Verlag.
19. Bryder L (2009) From breast to bottle: a history of modern infant feeding. *Endeavour* **33**, 54–59.
20. Droese W (1985) Zur Geschichte der Beikost in der Säuglingsernährung (History of complementary foods during the infant feeding period). In *Beikost in der Säuglingsernährung*, pp. 111–117 [H Ewerbeck, editor]. Berlin: Springer.
21. Obladen M (2014) Technical inventions that enabled artificial infant feeding. *Neonatology* **106**, 62–68.
22. Stevens EE, Patrick TE & Pickler R (2009) A history of infant feeding. *J Perinat Educ* **18**, 32–39.
23. Waterlow JC (editor) (2006) *Protein–Energy Malnutrition*. New Barnet: Smith-Gordon.
24. Golden MH (2002) The development of concepts of malnutrition. *J Nutr* **132**, 2117S–2122S.
25. Allen LH (2000) Ending hidden hunger: the history of micronutrient deficiency control. In *Background Paper of the World Bank–UNICEF Nutrition Assessment Project*. Washington, DC: World Bank.
26. Semba RD (2012) The historical evolution of thought regarding multiple micronutrient nutrition. *J Nutr* **142**, 143S–156S.
27. Waterlow JC (1961) The rate of recovery of malnourished infants in relation to the protein and calorie levels of the diet. *J Trop Pediatr* **7**, 16–22.
28. Papastavrou M, Genitsaridi SM, Komodiki E, *et al.* (2015) Breastfeeding in the course of history. *J Pediatr Neo Care* **2**, 00096.
29. Weaver LT (2011) How did babies grow 100 years ago? *Eur J Clin Nutr* **65**, 3–9.
30. Weaver LT (2006) The emergence of our modern understanding of infant nutrition and feeding 1750–1900. *Curr Paediatr* **16**, 342–347.
31. Fomon S (2001) Infant feeding in the 20th century: formula and beikost. *J Nutr* **131**, 409S–420S.
32. Anderson S (1996) Then and now: infant feeding in Britain, 1900–1914. *Prof Care Mother Child* **6**, 167, 170.
33. Forsyth D (1911) The history of infant-feeding from Elizabethan times. *Proc R Soc Med* **4**, 110–141.
34. Fulminante F (2015) Infant feeding practices in Europe and the Mediterranean from prehistory to the middle ages: a comparison between the historical sources and bio-archaeology. *Child Past* **8**, 24–47.
35. Keusch GT (2003) The history of nutrition: malnutrition, infection and immunity. *J Nutr* **133**, 336S–340S.
36. Rijpma S (1996) Malnutrition in the history of tropical Africa. *Civilisations* **43**, 45–63.
37. Heikens GT & Manary M (2009) 75 Years of Kwashiorkor in Africa. *Malawi Med J* **21**, 96–98.
38. Vernon K (2001) Milk and dairy products. In *The Cambridge Food History*, vol. 1, pp. 692–701 [KF Kiple and KC Ornelas, editors]. Cambridge: Cambridge University Press.
39. Suri DJ, Moorthy D & Rosenberg IH (2016) The role of dairy in effectiveness and cost of treatment of children with moderate acute malnutrition: a narrative review. *Food Nutr Bull* **37**, 176–185.
40. Obladen M (2014) Milk demystified by chemistry. *J Perinat Med* **42**, 641–647.
41. Muehlhoff E, Bennett A & McMahon D (2013) *Milk and Dairy Products in Human Nutrition*. Rome: FAO.
42. Sadler K, Kerven C, Calo M, *et al.* (2009) *Milk Matters: A Literature Review of Pastoralist Nutrition and Programming Responses*. Addis Ababa: Feinstein International Center, Tufts University and Save the Children.
43. Valenze D (2011) *Milk – A Local and Global History*. New Haven and London: Yale University Press.
44. Corvalan C, Dangour AD & Uauy R (2008) Need to address all forms of childhood malnutrition with a common agenda. *Arch Dis Child* **93**, 361–362.
45. Scherbaum V (2013) Entwicklung und Erprobung ernährungstherapeutischer Interventionen zur Bekämpfung kindlicher Mangelernährung (Development and testing of nutritional interventions to combat childhood malnutrition). Assistant Professor Cumulative Habilitation, University of Hohenheim.
46. Merriam-Webster (2013) *Merriam-Webster's Dictionary Unabridged*. Springfield, MA: Merriam-Webster Inc.
47. Davidson T (1908) *Chambers's Twentieth Century Dictionary of the English Language*. Edinburgh: W. R. Chambers Ltd.
48. Normet L (1926) La bouffissure d'Annam (Annam's puffiness). *Bull Soc Pathol Exotique* **3**, 207–213.
49. Trowell HC, Davies JN & Dean RFA (1954) *Kwashiorkor: Part 1. Reports of Kwashiorkor in Children and a Discussion of Terminology*. London: Edward Arnold. (Reprinted in 1982 by the Nutrition Foundation, Academic Press, New York.)
50. Autret M & Behar M (1954) Infantile multiple deficiency syndrome in Central America (Kwashiorkor). *Bull World Health Organ* **11**, 891–966.
51. Czerny A & Keller A (1913, 1928) *Des Kindes Ernährung, Ernährungsstörungen und Ernährungstherapie: ein Handbuch für Ärzte (Of Child Nutrition, Nutritional Disorders and Nutritional Therapy: A Manual for Doctors)*. Leipzig, Vienna: F. Deuticke.



52. Williams CD (1935) Kwashiorkor: a nutritional disease of children associated with a maize diet. *Lancet* **226**, 1151–1152.
53. Trowell HC (1940) Infantile pellagra. *Trans R Soc Trop Med Hyg* **33**, 389–390.
54. Trowell HC & Muwazi EM (1945) A contribution to the study of malnutrition in Central Africa; a syndrome of malignant malnutrition. *Trans R Soc Trop Med Hyg* **39**, 229–243.
55. Waterlow JC (1948) *Fatty Liver Disease in Infants in the British West Indies*. Medical Research Council Special Report Series. London: His Majesty's Stationery Office.
56. Brock JF & Autret M (1952) Kwashiorkor in Africa. *Bull World Health Organ* **5**, 1–71.
57. Jelliffe DB (1959) Protein–calorie malnutrition in tropical preschool children; a review of recent knowledge. *J Pediatr* **54**, 227–256.
58. Tanner JM (editor) (1981) *A History of the Study of Human Growth*. Cambridge: Cambridge University Press.
59. Drillien CM (1958) Growth and development in a group of children of very low birth weight. *Arch Dis Child* **33**, 10–18.
60. Anonymous (1965) Intrauterine malnutrition. *JAMA* **191**, 1077–1078.
61. Warkany J, Monroe BB & Sutherland BS (1961) Intrauterine growth retardation. *Am J Dis Child* **102**, 249–279.
62. Schulte FJ, Michaelis R & Nolte R (1967) Meinhard von Pfaundler and the history of small-for-dates infants. *Dev Med Child Neurol* **9**, 511.
63. Rumbolz WL & McGoogan LS (1953) Placental insufficiency and the small undernourished full-term infant. *Obstet Gynecol* **1**, 294–301.
64. Walker J (1967) 'Small for dates' – clinical aspects. *Proc R Soc Med* **60**, 877–879.
65. Wellcome Trust Working Party (1970) Classification of infantile malnutrition. *Lancet* **ii**, 302–303.
66. World Health Organization (1992) *National Strategies for Overcoming Micronutrient Malnutrition. Thirteenth Meeting, 27 January 1992*. Geneva: WHO.
67. Maberly GF, Trowbridge FL, Yip R, *et al.* (1994) Programs against micronutrient malnutrition: ending hidden hunger. *Annu Rev Public Health* **15**, 277–301.
68. World Health Organization (1999) *Management of Severe Malnutrition: a Manual for Physicians and Other Senior Health Workers*. Geneva: WHO.
69. Grobler-Tanner C & Collins S (2004) *Community Therapeutic Care (CTC): A New Approach to Managing Acute Malnutrition in Emergencies and Beyond. Technical Notes*. Washington, DC: FANTA.
70. Lacaille AD (1950) Infant feeding-bottles in prehistoric times. *Proc R Soc Med* **43**, 565–568.
71. Tamime AY (2002) Fermented milks: a historical food with modern applications – a review. *Eur J Clin Nutr* **56**, Suppl. 4, S2–S15.
72. Holsinger VH, Rajkovski KH & Stabel JR (1997) Milk pasteurisation and safety: a brief history and update. *Rev Sci Tech* **16**, 441–451.
73. Salque M, Bogucki PI, Pyzel J, *et al.* (2013) Earliest evidence for cheese making in the sixth millennium BC in northern Europe. *Nature* **493**, 522–525.
74. Radbill SX (1981) Infant feeding through the ages. *Clin Pediatr (Phila)* **20**, 613–621.
75. Cone TE (1981) History of infant feeding. From the earliest years through the development of scientific concepts. In *Infant and Child Feeding*, pp. 4–34 [JT Bond, editor]. New York: Academic Press.
76. Schuman AJ (2003) A concise history of infant formula (twists and turns included). *Contemp Pediatr* **20**, 91–103.
77. Grabmayr S & Scherbaum V (2003) Ernährungsformen in den ersten Lebenstagen (Diets during the first days of life). In *Stillen Fröhkindliche Ernährung und reproduktive Gesundheit (Breastfeeding Nutrition in Early Childhood and Reproductive Health)*, pp. 71–74 [V Scherbaum, U Kretschmer and FM Perl, editors]. Cologne: Deutscher Ärzte-Verlag.
78. Barness LA (1987) History of infant feeding practices. *Am J Clin Nutr* **46**, 168–170.
79. Heubner O (1913) *Festschrift Dr. Otto L. Heubner zum 70. Geburtstag und zum Andenken an den Abschluss seiner Lehrtätigkeit. Gewidmet von seinen Schülern (A Collection of Writings Published in Honour of a Scholar Dr. Otto L. Heubner's 70th Birthday and to Commemorate the Completion of His Teaching. Dedicated to His Students)*. Berlin and Heidelberg: Springer.
80. Mephram TB (1993) "Humanizing" milk: the formulation of artificial feeds for infants (1850–1910). *Med Hist* **37**, 225–249.
81. Ploss HH (1853) Über das aufziehen der kinder ohne brust (Raising children without the breast). *J Kinderkrankheiten* **20**, 217–225.
82. Wood AL (1955) The history of artificial feeding of infants. *J Am Diet Assoc* **31**, 474–482.
83. Obladen M (2014) From swill milk to certified milk: progress in cow's milk quality in the 19th century. *Ann Nutr Metab* **64**, 80–87.
84. Wickes JG (1953) A history of infant feeding. IV. Nineteenth century continued. *Arch Dis Child* **28**, 416–422.
85. Lee KS (2007) Infant mortality decline in the late 19th and early 20th centuries: the role of market milk. *Perspect Biol Med* **50**, 585–602.
86. Exner M, Hartemann P & Kistemann T (2001) Hygiene and health – the need for a holistic approach. *Am J Infect Control* **29**, 228–231.
87. Bloomfield SF & Scott EA (2003) Developing an effective policy for home hygiene: a risk-based approach. *Int J Environ Health Res* **13**, Suppl. 1, S57–S66.
88. Winkelstein W Jr (1995) A new perspective on John Snow's communicable disease theory. *Am J Epidemiol* **142**, S3–S9.
89. Paneth N & Fine P (2013) The art of medicine: the singular science of John Snow. *Lancet* **381**, 1267–1269.
90. Carter KC (1977) The germ theory, beriberi, and the deficiency theory of disease. *Med Hist* **21**, 119–136.
91. Bennett JA (2001) Germs or rations? Beriberi and the Japanese labor experiment in colonial Fiji and Queensland. *Pac Stud* **24**, 1–17.
92. Ihde AJ & Becker SL (1971) Conflict of concepts in early vitamin studies. *J Hist Biol* **4**, 1–33.
93. Funk C (1912) The etiology of deficiency diseases. *J State Med* **20**, 341–368.
94. Rajakumar K (2001) Infantile scurvy: a historical perspective. *Pediatrics* **108**, E76.
95. Kraus F, Meyer E, Minkowski O, *et al.* (1920) *Ergebnisse der Inneren Medizin und Kinderheilkunde: Achtzehnter Band (Results of Internal Medicine and Paediatrics: Eighteenth Volume)*. Berlin: J. Springer.
96. Salge B (1910) *Einführung in die moderne Kinderheilkunde (Introduction to Modern Paediatrics)*, 2nd ed. Berlin: Springer.
97. Finkelstein H (1907) *Über alimentäre Intoxikation. Jahrbuch für Kinderheilkunde und physische Gesundheit*. 67. Berlin: S. Karger. <http://www.archive.org/stream/jahrbuchfuerkin00>

- unkngoog/jahrbuchfuerkin00unkngoog\_djvu.txt (accessed October 2017).
98. Nützenadel W (2010) *Des Kindes Ernährung – Ein Rückblick. Entwicklungen und Perspektiven der Kinder- und Jugendmedizin 150 Jahre Pädiatrie in Heidelberg (Of Child Nutrition – A Review. Developments and Perspectives of Child and Adolescent Medicine, 150 Years of Paediatrics in Heidelberg)*. Mainz: Kirchheim.
  99. Atkins P (2007) School milk in Britain, 1900–1934. *J Pol Hist* **19**, 395–427.
  100. Pollock J (2006) Two controlled trials of supplementary feeding of British school children in the 1920s. *J R Soc Med* **99**, 323–327.
  101. Allen LH & Dror DK (2011) Effects of animal source foods, with emphasis on milk, in the diet of children in low-income countries. *Nestle Nutr Workshop Ser Pediatr Program* **67**, 113–130.
  102. McCollum EV (1924) The nutritional value of milk. In *World's Dairy Congress, Washington DC, 2–10 October 1923*, pp. 421–437 [LA Rogers and RD Lenoir, editors]. Washington, DC: US Government Printing Office.
  103. Mann HCC (1926) *Diets for Boys During the School Age*. Special Report Series 105. London: Medical Research Council.
  104. Orr JB (1928) Influence of amount of milk consumption on the rate of growth of school children. *Br Med J* **1**, 140–141.
  105. Leighton G & Clark ML (1929) Milk consumption and the growth of school children: second preliminary report on tests to the Scottish Board of Health. *Br Med J* **1**, 23–25.
  106. Student (1931) The Lanarkshire milk experiment. *Biometrika* **23**, 398–406.
  107. Atkins P (2005) Fattening children or fattening farmers? School milk in Britain, 1921–1941. *Econ Hist Rev* **58**, 57–78.
  108. Sellick P (2001) Responding to children affected by armed conflict. a case study of Save the Children Fund (1919–1999). PhD Thesis, University of Bradford.
  109. Roberts SL (2010) Place, life histories, and the politics of relief: episodes in the life of Francesca Wilson, humanitarian educator activist. PhD Thesis, University of Birmingham.
  110. Rhodes BD (2001) *United States Foreign Policy in the Interwar Period 1918–1941*. Westport: Praeger.
  111. Williams CD (1933) A nutritional disease of childhood associated with a maize diet. *Arch Dis Child* **8**, 423–433.
  112. Stanton J (2001) Listening to the Ga: Cicely Williams' discovery of kwashiorkor on the Gold Coast. *Clio Med* **61**, 149–171.
  113. Richter J (2001) International regulation of transnational corporations: the infant food debate. PhD Thesis, Amsterdam School of Communication Research.
  114. Williams CD (1939) *Milk and Murder. Speech to the Singapore Rotary Club*. Penang: Malaysia International Organization of Consumers Union.
  115. Gopalan C (1967) Malnutrition in childhood in the tropics. *Br Med J* **4**, 603–607.
  116. Dean RF (1951) The nutritional adequacy of a vegetable substitute for milk. *Br J Nutr* **5**, 269–274.
  117. Dean RF (1952) The treatment of kwashiorkor with milk and vegetable proteins. *Br Med J* **2**, 791–796.
  118. Altmann A (1948) The syndrome of malignant malnutrition (kwashiorkor; infantile pellagra). Its conception as a protein deficiency and its treatment with skimmed lactic acid milk. *Clin Proc* **7**, 32–53.
  119. Walt F & Wills L (1950) Malignant malnutrition. *S Afr Med J* **24**, 920–925.
  120. Spies TD, Dreizen S, Snodgrass RM, *et al.* (1959) Effect of dietary supplement of nonfat milk on human growth failure; comparative response in undernourished children and in undernourished adolescents. *AMA J Dis Child* **98**, 187–197.
  121. Carpenter KJ (1994) *Protein and Energy. A Study of Changing Ideas in Nutrition*. Cambridge: Cambridge University Press.
  122. United Nations (1968) *Feeding the Expanding World Population: International Action to Avert the Impending Protein Crisis/Report to the Economic and Social Council of the Advisory Committee on the Application of Science and Technology to Development. Special Report Series*. New York: Economic and Social Council. Advisory Committee on the Application of Science and Technology to Development.
  123. Marchione TJ (2002) Foods provided through U.S. Government Emergency Food Aid Programs: policies and customs governing their formulation, selection and distribution. *J Nutr* **132**, 2104S–2111S.
  124. Fleige LE, Moore WR, Garlick PJ, *et al.* (2010) Recommendations for optimization of fortified and blended food aid products from the United States. *Nutr Rev* **68**, 290–315.
  125. Food and Agriculture Organization (1965) *Protein Requirements. Report of a Joint FAO/WHO Expert Group*. FAO Nutrition Meetings Report Series No. 37. Rome: FAO.
  126. Food and Agriculture Organization & World Health Organization (1973) *Energy and Protein Requirements. Report of a Joint FAO/WHO Ad Hoc Expert Committee*. Geneva: WHO and FAO.
  127. Food and Agriculture Organization, World Health Organization & United Nations University (1985) *Energy and Protein Requirements. Report of a Joint FAO/WHO/UNU Expert Consultation*. WHO Technical Report Series no. 724. Geneva: WHO.
  128. Scherbaum V & Furst P (2000) New concepts on nutritional management of severe malnutrition: the role of protein. *Curr Opin Clin Nutr Metab Care* **3**, 31–38.
  129. McLaren DS (1974) The great protein fiasco. *Lancet* **ii**, 93–96.
  130. Golden MH (1982) Protein deficiency, energy deficiency, and the oedema of malnutrition. *Lancet* **i**, 1261–1265.
  131. Welbourne H (1955) The danger period during weaning: a study of Baganda children who were attending child welfare clinics near Kampala, Uganda. *J Trop Pediatr* **1**, 34–46.
  132. Welbourne H (1955) The danger period during weaning. A study of Baganda children who were attending child welfare clinics near Kampala, Uganda [Parts II and III]. *J Trop Pediatr* **1**, 98–111, 161–173.
  133. Hendrickse RG, Coulter JB, Lamplugh SM, *et al.* (1982) Aflatoxins and kwashiorkor: a study in Sudanese children. *Br Med J (Clin Res Ed)* **285**, 843–846.
  134. Golden MH & Ramdath D (1987) Free radicals in the pathogenesis of kwashiorkor. *Proc Nutr Soc* **46**, 53–68.
  135. McLellan A (2014) Does the distribution of ready to use food products for the prevention of undernutrition meet the ultimate needs of the beneficiary? *Afr J Food Agric Nutr Dev* **14**, 8956–8962.
  136. Kane AV, Dinh DM & Ward HD (2015) Childhood malnutrition and the intestinal microbiome. *Pediatr Res* **77**, 256–262.
  137. Uauy R (2013) Keynote: rethinking protein. *Food Nutr Bull* **34**, 228–231.
  138. Anonymous (2013) Dietary protein quality evaluation in human nutrition. Report of an FAO Expert Consultation. *FAO Food Nutr Pap* **92**, 1–66.
  139. World Health Organization, Food and Agriculture Organization & United Nations University (2007) *Protein and*

*Amino Acid Requirements in Human Nutrition. Report of a Joint FAO/WHO/UNU Expert Consultation.* WHO Technical Report Series no. 935. Geneva: WHO.

140. Semba RD (2016) The rise and fall of protein malnutrition in global health. *Ann Nutr Metab* **69**, 79–88.
141. Ghosh S (2016) Protein quality in the first thousand days of life. *Food Nutr Bull* **37**, S14–S21.
142. Ghosh S, Suri D & Uauy R (2012) Assessment of protein adequacy in developing countries: quality matters. *Br J Nutr* **108**, Suppl. 2, S77–S87.
143. Schonfeldt HC & Gibson Hall N (2012) Dietary protein quality and malnutrition in Africa. *Br J Nutr* **108**, Suppl. 2, S69–S76.
144. Semba RD, Shardell M, Sakr Ashour FA, *et al.* (2016) Child stunting is associated with low circulating essential amino acids. *EBioMedicine* **6**, 246–252.
145. Uauy R, Suri DJ, Ghosh S, *et al.* (2016) Low circulating amino acids and protein quality: an interesting piece in the puzzle of early childhood stunting. *EBioMedicine* **8**, 28–29.
146. Haug A, Hostmark AT & Harstad OM (2007) Bovine milk in human nutrition – a review. *Lipids Health Dis* **6**, 25.
147. Weaver C, Wijesinha-Bettoni R, McMahon D, *et al.* (2013) Milk and dairy products as part of the diet. In *Milk and Dairy Products in Human Nutrition*, pp. 103–206 [E Muehlhoff, A Bennett and D McMahon, editors]. Rome: FAO.
148. Wijesinha-Bettoni R & Burlingame B (2013) Milk and dairy product composition. In *Milk and Dairy Products in Human Nutrition*, pp. 43–52 [E Muehlhoff, A Bennett and D McMahon, editors]. Rome: FAO.
149. Reeds P, Schaafsma G, Tome D, *et al.* (2000) Criteria and significance of dietary protein sources in humans. Summary of the workshop with recommendations. *J Nutr* **130**, 1874S–1876S.
150. Boye J, Wijesinha-Bettoni R & Burlingame B (2012) Protein quality evaluation twenty years after the introduction of the protein digestibility corrected amino acid score method. *Br J Nutr* **108**, Suppl. 2, S183–S211.
151. Pereira PC (2014) Milk nutritional composition and its role in human health. *Nutrition* **30**, 619–627.
152. Gat-Yablonski G, Yackobovitch-Gavan M & Phillip M (2017) Which dietary components modulate longitudinal growth? *Curr Opin Clin Nutr Metab Care* **20**, 211–216.
153. Oakley E, Reinking J, Sandige H, *et al.* (2010) A ready-to-use therapeutic food containing 10% milk is less effective than one with 25% milk in the treatment of severely malnourished children. *J Nutr* **140**, 2248–2252.
154. Batra P, Schlossman N, Balan I, *et al.* (2016) A randomized controlled trial offering higher- compared with lower-dairy second meals daily in preschools in Guinea-Bissau demonstrates an attendance-dependent increase in weight gain for both meal types and an increase in mid-upper arm circumference for the higher-dairy meal. *J Nutr* **146**, 124–132.
155. Stobaugh HC, Ryan KN, Kennedy JA, *et al.* (2016) Including whey protein and whey permeate in ready-to-use supplementary food improves recovery rates in children with moderate acute malnutrition: a randomized, double-blind clinical trial. *Am J Clin Nutr* **103**, 926–933.
156. Grenov B, Briend A, Sangild PT, *et al.* (2016) Undernourished children and milk lactose. *Food Nutr Bull* **37**, 85–99.
157. Karav S, Le Parc A, Leite Nobrega de Moura Bell JM, *et al.* (2016) Oligosaccharides released from milk glycoproteins are selective growth substrates for infant-associated bifidobacteria. *Appl Environ Microbiol* **82**, 3622–3630.
158. Park YW & Nam MS (2015) Bioactive peptides in milk and dairy products: a review. *Korean J Food Sci Anim Resour* **35**, 831–840.
159. Ashworth A (1980) Practical aspects of dietary management during rehabilitation from severe protein–energy malnutrition. *J Hum Nutr* **34**, 360–369.
160. Ashworth A (1979) Progress in the treatment of protein–energy malnutrition. *Proc Nutr Soc* **38**, 89–97.
161. Ashworth A, Jackson A, Khanum S, *et al.* (1996) Ten steps to recovery. *Child Health Dialogue* **1996**, 10–12.
162. Briend A & Golden MH (1993) Treatment of severe child malnutrition in refugee camps. *Eur J Clin Nutr* **47**, 750–754.
163. Ashworth A & Burgess A (2003) *Caring for Severely Malnourished Children*. Oxford: Macmillan Education Ltd.
164. Briend A, Lacsala R, Prudhon C, *et al.* (1999) Ready-to-use therapeutic food for treatment of marasmus. *Lancet* **353**, 1767–1768.
165. Collins S & Sadler K (2002) Outpatient care for severely malnourished children in emergency relief programmes: a retrospective cohort study. *Lancet* **360**, 1824–1830.
166. World Health Organization, World Food Programme, United Nations System Standing Committee on Nutrition, *et al.* (2007) *Community-Based Management of Severe Acute Malnutrition. A Joint Statement by the World Health Organization, the World Food Programme, the United Nations System Standing Committee on Nutrition and the United Nations Children's Fund*. Geneva, New York and Rome: WHO, UNICEF, WFP, UN-SCN.
167. World Health Organization (2013) *Guideline: Updates on the Management of Severe Acute Malnutrition in Infants and Children*. Geneva: WHO.
168. Collins S, Dent N, Binns P, *et al.* (2006) Management of severe acute malnutrition in children. *Lancet* **368**, 1992–2000.
169. Latham MC, Jonsson U, Sterken E, *et al.* (2011) RUTF stuff. Can the children be saved with fortified peanut paste? (Correspondence). *World Nutr* **2**, 62–85.
170. Greiner T (2014) *The Advantages, Disadvantages and Risks of Ready-to-Use Foods*. IBFAN breastfeeding briefs, no. 56/57. Geneva: GIFA.
171. Sandige H, Ndekha MJ, Briend A, *et al.* (2004) Home-based treatment of malnourished Malawian children with locally produced or imported ready-to-use food. *J Pediatr Gastroenterol Nutr* **39**, 141–146.
172. Krumbein T, Scherbaum V & Biesalski HK (2006) Locally produced ready-to-use therapeutic food (RUTF) in an inpatient setting in Uganda. *Field Exchange* **28**, 21.
173. Weber JM, Ryan KN, Tandon R, *et al.* (2017) Acceptability of locally produced ready-to-use therapeutic foods in Ethiopia, Ghana, Pakistan and India. *Matern Child Nutr* **13**, 12250.
174. Purwestri RC, Scherbaum V, Inayati DA, *et al.* (2012) Cost analysis of community-based daily and weekly programs for treatment of moderate and mild wasting among children on Nias Island, Indonesia. *Food Nutr Bull* **33**, 207–216.
175. LNS Network (2009) LNS Research Network Meeting Report, Rome, February 6, 2009. <http://www.unhcr.org/4b7532529.pdf> (accessed October 2017).
176. Shoham J & Duffield A (2009) Proceedings of the World Health Organization/UNICEF/World Food Programme/United Nations High Commissioner for Refugees Consultation on the management of moderate malnutrition in children under 5 years of age. *Food Nutr Bull* **30**, S464–S474.

177. de Pee S & Bloem MW (2009) Current and potential role of specially formulated foods and food supplements for preventing malnutrition among 6- to 23-month-old children and for treating moderate malnutrition among 6- to 59-month-old children. *Food Nutr Bull* **30**, S434–S463.
178. Schlossman N, Brown C, Batra P, *et al.* (2017) A randomized controlled trial of two ready-to-use supplementary foods demonstrates benefit of the higher dairy supplement for reduced wasting in mothers, and differential impact in infants and children associated with maternal supplement response. *Food Nutr Bull* **38**, 275–290.
179. Chaparro CM & Dewey KG (2010) Use of lipid-based nutrient supplements (LNS) to improve the nutrient adequacy of general food distribution rations for vulnerable sub-groups in emergency settings. *Matern Child Nutr* **6**, Suppl. 1, 1–69.
180. Emergency Nutrition Network (2013) *The Management of Acute Malnutrition: A Review of Donor and Government Financing Arrangements*. Geneva: Emergency Nutrition Network (ENN).
181. Manary MJ (2006) Local production and provision of ready-to-use therapeutic food (RUTF) spread for the treatment of severe childhood malnutrition. *Food Nutr Bull* **27**, S83–S89.
182. Hoppe C, Andersen GS, Jacobsen S, *et al.* (2008) The use of whey or skimmed milk powder in fortified blended foods for vulnerable groups. *J Nutr* **138**, 145S–161S.
183. LaGrone LN, Trehan I, Meuli GJ, *et al.* (2012) A novel fortified blended flour, corn–soy blend “plus-plus,” is not inferior to lipid-based ready-to-use supplementary foods for the treatment of moderate acute malnutrition in Malawian children. *Am J Clin Nutr* **95**, 212–219.
184. Bahwere P, Banda T, Sadler K, *et al.* (2014) Effectiveness of milk whey protein-based ready-to-use therapeutic food in treatment of severe acute malnutrition in Malawian under-5 children: a randomised, double-blind, controlled non-inferiority clinical trial. *Matern Child Nutr* **10**, 436–451.
185. Michaelsen KF, Hoppe C, Roos N, *et al.* (2009) Choice of foods and ingredients for moderately malnourished children 6 months to 5 years of age. *Food Nutr Bull* **30**, S343–S404.
186. Lhotska L, Scherbaum V & Bellows AC (2015) Maternal, infant, and young child feeding: intertwined subjectivities and corporate accountability. In *Gender, Nutrition, and the Human Right to Adequate Food Toward an Inclusive Framework*, pp. 162–253 [AC Bellows, FLS Valente, S Lemke and MD Nunez Burbano de Lara, editors]. New York and Abingdon: Taylor and Francis Routledge.
187. Purwestri RC, Scherbaum V, Inayati DA, *et al.* (2013) Impact of daily *versus* weekly supply of locally produced ready-to-use food on growth of moderately wasted children on Nias Island, Indonesia. *ISRN Nutr* **2013**, 412145.
188. Purwestri RC, Scherbaum V, Inayati DA, *et al.* (2012) Supplementary feeding with locally-produced ready-to-use food (RUF) for mildly wasted children on Nias Island, Indonesia: comparison of daily and weekly program outcomes. *Asia Pac J Clin Nutr* **21**, 374–379.
189. Kana Sop MM, Gouado I, Mananga MJ, *et al.* (2012) Trace elements in foods of children from Cameroon: a focus on zinc and phytate content. *J Trace Elem Med Biol* **26**, 201–204.
190. Hotz C & Gibson RS (2007) Traditional food-processing and preparation practices to enhance the bioavailability of micronutrients in plant-based diets. *J Nutr* **137**, 1097–1100.
191. Weber J & Callaghan M (2016) Optimizing ready-to-use therapeutic foods for protein quality, cost, and acceptability. *Food Nutr Bull* **37**, S37–S46.
192. Scherbaum V, Purwestri RC, Stuetz W, *et al.* (2015) Locally produced cereal/nut/legume-based biscuits *versus* peanut/milk-based spread for treatment of moderately to mildly wasted children in daily programmes on Nias Island, Indonesia: an issue of acceptance and compliance? *Asia Pac J Clin Nutr* **24**, 152–161.
193. Delchevalerie P, Van Herp M, Degroot N, *et al.* (2015) Ready-to-use supplementary food *versus* corn soya blend with oil premix to treat moderate acute child malnutrition: a community-based cluster randomized trial. <https://doi.org/10.13140/rg.2.1.1367.5366> (accessed October 2017).
194. Nikiema L, Huybregts L, Kolsteren P, *et al.* (2012) Treating moderate acute malnutrition in first-line health services: an effectiveness cluster-randomized trial in Burkina Faso. *Am J Clin Nutr* **100**, 241–249.
195. Ackatia-Armah RS, McDonald CM, Doumbia S, *et al.* (2015) Malian children with moderate acute malnutrition who are treated with lipid-based dietary supplements have greater weight gains and recovery rates than those treated with locally produced cereal–legume products: a community-based, cluster-randomized trial. *Am J Clin Nutr* **101**, 632–645.
196. Karakochuk C, van den Briel T, Stephens D, *et al.* (2012) Treatment of moderate acute malnutrition with ready-to-use supplementary food results in higher overall recovery rates compared with a corn–soya blend in children in southern Ethiopia: an operations research trial. *Am J Clin Nutr* **96**, 911–916.
197. Nackers F, Broillet F, Oumarou D, *et al.* (2010) Effectiveness of ready-to-use therapeutic food compared to a corn/soy-blend-based pre-mix for the treatment of childhood moderate acute malnutrition in Niger. *J Trop Pediatr* **56**, 407–413.
198. Thakwalakwa C, Ashorn P, Phuka J, *et al.* (2010) A lipid-based nutrient supplement but not corn–soy blend modestly increases weight gain among 6- to 18-month-old moderately underweight children in rural Malawi. *J Nutr* **140**, 2008–2013.
199. Matilsky DK, Maleta K, Castleman T, *et al.* (2009) Supplementary feeding with fortified spreads results in higher recovery rates than with a corn/soy blend in moderately wasted children. *J Nutr* **139**, 773–778.
200. Patel MP, Sandige HL, Ndekha MJ, *et al.* (2005) Supplemental feeding with ready-to-use therapeutic food in Malawian children at risk of malnutrition. *J Health Popul Nutr* **23**, 351–357.
201. Irena AH, Bahwere P, Owino VO, *et al.* (2013) Comparison of the effectiveness of a milk-free soy–maize–sorghum-based ready-to-use therapeutic food to standard ready-to-use therapeutic food with 25% milk in nutrition management of severely acutely malnourished Zambian children: an equivalence non-blinded cluster randomised controlled trial. *Matern Child Nutr* **11** Suppl. 4, 105–119.
202. Bahwere P, Balaluka B, Wells JC, *et al.* (2016) Cereals and pulse-based ready-to-use therapeutic food as an alternative to the standard milk- and peanut paste-based formulation for treating severe acute malnutrition: a noninferiority, individually randomized controlled efficacy clinical trial. *Am J Clin Nutr* **103**, 1145–1161.
203. Bahwere P, Akomo P, Mwale M, *et al.* (2017) Soya, maize, and sorghum-based ready-to-use therapeutic food with amino acid is as efficacious as the standard milk and peanut paste-based formulation for the treatment of severe acute

- malnutrition in children: a noninferiority individually randomized controlled efficacy clinical trial in Malawi. *Am J Clin Nutr* **106**, 1100–1112.
204. Rosenberg I, Rogers B, Webb P, *et al.* (2012) Enhancements in food aid quality need to be seen as a process, not as a one-off event. *J Nutr* **142**, 1781.
205. DiRienzo D (2016) Research gaps in the use of dairy ingredients in food aid products. *Food Nutr Bull* **37**, S51–S57.
206. Schulz KF, Altman DG & Moher D (2010) CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *BMJ* **340**, c332.
207. Save the Children & Humanitarian Innovation Fund (2015) Standardised Indicators and Categories for Better cMAM Reporting, April 2015 edition. <http://www.cmamreport.com/sites/all/themes/stc/cmam-assets/STANDARDISED%20CATEGORIES%20AND%20INDICATORS%20FOR%20BETTER%20CMAM%20REPORTING%20%20FINAL%20Apr%202015.pdf>
208. Inayati DA, Scherbaum V, Purwestri RC, *et al.* (2012) Combined intensive nutrition education and micronutrient powder supplementation improved nutritional status of mildly wasted children on Nias Island, Indonesia. *Asia Pac J Clin Nutr* **21**, 361–373.
209. Noriega KE & Lindshield BL (2014) Is the inclusion of animal source foods in fortified blended foods justified? *Nutrients* **6**, 3516–3535.
210. Scherbaum V & Srouf ML (2016) The role of breastfeeding in the prevention of childhood malnutrition. In *Hidden Hunger Malnutrition in the First 1,000 Days of Life, Consequences and Solutions*, vol. 115, pp. 75–90 [HK Biesalski and RE Black, editors]. Basel: Karger.
211. Pencharz P, Jahoor F, Kurpad A, *et al.* (2014) Current issues in determining dietary protein and amino-acid requirements. *Eur J Clin Nutr* **68**, 285–286.
212. Singhal A (2017) Long-term adverse effects of early growth acceleration or catch-up growth. *Ann Nutr Metab* **70**, 236–240.
213. Thakwalakwa CM, Ashorn P, Jawati M, *et al.* (2012) An effectiveness trial showed lipid-based nutrient supplementation but not corn-soya blend offered a modest benefit in weight gain among 6- to 18-month-old underweight children in rural Malawi. *Public Health Nutr* **15**, 1755–1762.
214. Phuka JC, Gladstone M, Maleta K, *et al.* (2012) Developmental outcomes among 18-month-old Malawians after a year of complementary feeding with lipid-based nutrient supplements or corn-soy flour. *Matern Child Nutr* **8**, 239–248.
215. Maleta KM, Phuka J, Alho L, *et al.* (2015) Provision of 10–40 g/d lipid-based nutrient supplements from 6 to 18 months of age does not prevent linear growth faltering in Malawi. *J Nutr* **145**, 1909–1915.
216. Ashorn P, Alho L, Ashorn U, *et al.* (2015) Supplementation of maternal diets during pregnancy and for 6 months postpartum and infant diets thereafter with small-quantity lipid-based nutrient supplements does not promote child growth by 18 months of age in rural Malawi: a randomized controlled trial. *J Nutr* **145**, 1345–1353.
217. Mangani C, Maleta K, Phuka J, *et al.* (2013) Effect of complementary feeding with lipid-based nutrient supplements and corn-soy blend on the incidence of stunting and linear growth among 6- to 18-month-old infants and children in rural Malawi. *Matern Child Nutr* **11**, Suppl. 4, 132–143.
218. Lazzarini M, Rubert I & Pani P (2013) Specially formulated foods for treating children with moderate acute malnutrition in low- and middle-income countries. *Cochrane Database Syst Rev*, issue 6, CD009584.
219. Committee on Nutrition American Academy of Pediatrics (1971) Iron fortified formulas. *Pediatrics* **47**, 786.
220. Dror DK & Allen LH (2011) The importance of milk and other animal-source foods for children in low-income countries. *Food Nutr Bull* **32**, 227–243.
221. Fomon SJ, Ziegler EE, Nelson SE, *et al.* (1981) Cow milk feeding in infancy: gastrointestinal blood loss and iron nutritional status. *J Pediatr* **98**, 540–545.
222. Dewey KG (2005) *Guiding Principles for Feeding Non-Breastfed Children 6–24 months of Age*. Geneva: World Health Organization.
223. Anonymous (1992) American Academy of Pediatrics Committee on Nutrition: the use of whole cow's milk in infancy. *Pediatrics* **89**, 1105–1109.
224. Ziegler EE (2011) Consumption of cow's milk as a cause of iron deficiency in infants and toddlers. *Nutr Rev* **69**, Suppl. 1, S37–S42.
225. Domellof M, Braegger C, Campoy C, *et al.* (2014) Iron requirements of infants and toddlers. *J Pediatr Gastroenterol Nutr* **58**, 119–129.
226. Ziegler EE (2007) Adverse effects of cow's milk in infants. *Nestle Nutr Workshop Ser Pediatr Program* **60**, 185–196; discussion 196–199.
227. Black RE, Bhutta SZ, Bryce J, *et al.* (2008) *The Lancet's Series on Maternal and Child Undernutrition Executive Summary*. London: The Lancet.
228. Weaver LT (2006) Rapid growth in infancy: balancing the interests of the child. *J Pediatr Gastroenterol Nutr* **43**, 428–432.
229. Bhutta ZA, Ahmed T, Black RE, *et al.* (2008) What works? Interventions for maternal and child undernutrition and survival. *Lancet* **371**, 417–440.
230. Eid EE (1970) Follow-up study of physical growth of children who had excessive weight gain in first six months of life. *Br Med J* **2**, 74–76.
231. Druet C, Stettler N, Sharp S, *et al.* (2012) Prediction of childhood obesity by infancy weight gain: an individual-level meta-analysis. *Paediatr Perinat Epidemiol* **26**, 19–26.
232. Singhal A (2016) The role of infant nutrition in the global epidemic of non-communicable disease. *Proc Nutr Soc* **75**, 162–168.
233. Marinkovic T, Toemen L, Kruithof CJ, *et al.* (2017) Early infant growth velocity patterns and cardiovascular and metabolic outcomes in childhood. *J Pediatr* **186**, 57–63.e4.
234. Mameli C, Mazzantini S & Zuccotti GV (2016) Nutrition in the first 1000 days: the origin of childhood obesity. *Int J Environ Res Public Health* **13**, E838.
235. Woo Baidal JA, Locks LM, Cheng ER, *et al.* (2016) Risk factors for childhood obesity in the first 1,000 days: a systematic review. *Am J Prev Med* **50**, 761–779.
236. Patro-Golab B, Zalewski BM, Kolodziej M, *et al.* (2016) Nutritional interventions or exposures in infants and children aged up to 3 years and their effects on subsequent risk of overweight, obesity and body fat: a systematic review of systematic reviews. *Obes Rev* **17**, 1245–1257.
237. Koletzko B, von Kries R, Closa R, *et al.* (2009) Can infant feeding choices modulate later obesity risk? *Am J Clin Nutr* **89**, 1502S–1508S.
238. Lu L, Xun P, Wan Y, *et al.* (2016) Long-term association between dairy consumption and risk of childhood obesity: a systematic review and meta-analysis of prospective cohort studies. *Eur J Clin Nutr* **70**, 414–423.
239. Hoppe C, Molgaard C, Juul A, *et al.* (2004) High intakes of skimmed milk, but not meat, increase serum IGF-I

- and IGFBP-3 in eight-year-old boys. *Eur J Clin Nutr* **58**, 1211–1216.
240. Hoppe C, Molgaard C, Vaag A, *et al.* (2005) High intakes of milk, but not meat, increase s-insulin and insulin resistance in 8-year-old boys. *Eur J Clin Nutr* **59**, 393–398.
  241. Socha P, Grote V, Gruszfeld D, *et al.* (2011) Milk protein intake, the metabolic–endocrine response, and growth in infancy: data from a randomized clinical trial. *Am J Clin Nutr* **94**, 1776S–1784S.
  242. Jain V & Singhal A (2012) Catch up growth in low birth weight infants: striking a healthy balance. *Rev Endocr Metab Disord* **13**, 141–147.
  243. Lebenthal Y, Yackobovitch-Gavan M, Lazar L, *et al.* (2014) Effect of a nutritional supplement on growth in short and lean prepubertal children: a prospective, randomized, double-blind, placebo-controlled study. *J Pediatr* **165**, 1190–1193.e1.
  244. Masarwi M, Gabet Y, Dolkart O, *et al.* (2016) Skeletal effect of casein and whey protein intake during catch-up growth in young male Sprague–Dawley rats. *Br J Nutr* **116**, 59–69.
  245. Yackobovitch-Gavan M, Lebenthal Y, Lazar L, *et al.* (2016) Effect of nutritional supplementation on growth in short and lean prepubertal children after 1 year of intervention. *J Pediatr* **179**, 154–159.e1.
  246. Vandenplas Y (2015) Lactose intolerance. *Asia Pac J Clin Nutr* **24**, Suppl. 1, S9–S13.
  247. American Academy of Pediatrics, Committee on Nutrition (1990) Practical significance of lactose intolerance in children: supplement. *Pediatrics* **86**, 643–644.
  248. Solomons NW, Torun B, Caballero B, *et al.* (1984) The effect of dietary lactose on the early recovery from protein–energy malnutrition. I. Clinical and anthropometric indices. *Am J Clin Nutr* **40**, 591–600.
  249. Savaiano DA, Boushey CJ & McCabe GP (2006) Lactose intolerance symptoms assessed by meta-analysis: a grain of truth that leads to exaggeration. *J Nutr* **136**, 1107–1113.
  250. Torun B, Solomons NW, Caballero B, *et al.* (1984) The effect of dietary lactose on the early recovery from protein–energy malnutrition. II. Indices of nutrient absorption. *Am J Clin Nutr* **40**, 601–610.
  251. Brown KH, Peerson JM & Fontaine O (1994) Use of nonhuman milks in the dietary management of young children with acute diarrhea: a meta-analysis of clinical trials. *Pediatrics* **93**, 17–27.
  252. Ziegler EE & Fomon SJ (1983) Lactose enhances mineral absorption in infancy. *J Pediatr Gastroenterol Nutr* **2**, 288–294.
  253. Gaffey MF, Wazny K, Bassani DG, *et al.* (2013) Dietary management of childhood diarrhea in low- and middle-income countries: a systematic review. *BMC Public Health* **13**, Suppl. 3, S17.
  254. Scherbaum V (2003) Infant formula distribution in Northern Iraq. Summary of assessment. Field Exchange 20, November 2003. p5. [www.enonline.net/fex/20/infant](http://www.enonline.net/fex/20/infant)
  255. Scherbaum V (2003) Säuglingsernährung in Nordirak (Infant feeding in northern Iraq). *Ernährungs-Umschau* **50**, 476–480.
  256. Michaelsen KF, Nielsen AL, Roos N, *et al.* (2011) Cow's milk in treatment of moderate and severe undernutrition in low-income countries. *Nestle Nutr Workshop Ser Pediatr Program* **67**, 99–111.
  257. Jacobs C, Braun P & Hammer P (2011) Reservoir and routes of transmission of *Enterobacter sakazakii* (*Cronobacter* spp.) in a milk powder-producing plant. *J Dairy Sci* **94**, 3801–3810.
  258. Mourey A (2015) *Nutrition Manual for Humanitarian Action*. Geneva: International Committee of the Red Cross.
  259. Benabe JE & Martinez-Maldonado M (1998) The impact of malnutrition on kidney function. *Miner Electrolyte Metab* **24**, 20–26.
  260. Mason JB & Margetts BM (2017) Magic bullets vs community action: the trade-offs are real. *World Nutr* **8**, 5–25.
  261. Taylor CE & Taylor EM (1976) Multi-factorial causation of malnutrition. In *Nutrition in the Community*, pp. 75–85 [DS McLaren, editor]. Chichester: Wiley.
  262. United Nations Children's Fund (1990) *Strategy for Improved Nutrition of Children and Women in Developing Countries, UNICEF Policy Review*. New York: UNICEF.
  263. Schweitzer C (2016) Ready-to-use supplementary foods and ready-to-use therapeutic foods: developing product standards. *Food Nutr Bull* **37**, S47–S50.
  264. Mennella JA (2014) Ontogeny of taste preferences: basic biology and implications for health. *Am J Clin Nutr* **99**, 704S–711S.
  265. Galpin L, Thakwalakwa C, Phuka J, *et al.* (2007) Breast milk intake is not reduced more by the introduction of energy dense complementary food than by typical infant porridge. *J Nutr* **137**, 1828–1833.
  266. Owino VO, Bahwere P, Bisimwa G, *et al.* (2011) Breast-milk intake of 9-10-month-old rural infants given a ready-to-use complementary food in South Kivu, Democratic Republic of Congo. *Am J Clin Nutr* **93**, 1300–1304.
  267. Kumwenda C, Dewey KG, Hemsworth J, *et al.* (2014) Lipid-based nutrient supplements do not decrease breast milk intake of Malawian infants. *Am J Clin Nutr* **99**, 617–623.
  268. Flax VL, Siega-Riz AM, Reinhart GA, *et al.* (2015) Provision of lipid-based nutrient supplements to Honduran children increases their dietary macro- and micronutrient intake without displacing other foods. *Matern Child Nutr* **11**, Suppl. 4, 203–213.
  269. Anonymous (2017) Breastfeeding: a missed opportunity for global health. *Lancet* **390**, 532.
  270. Barennes H, Slesak G, Goyet S, *et al.* (2016) Enforcing the international code of marketing of breast-milk substitutes for better promotion of exclusive breastfeeding: can lessons be learned? *J Hum Lact* **32**, 20–27.
  271. Jelliffe DB (1972) Commercially induced malnutrition? Time for a dialogue. *Nutr Rev* **30**, 199–205.
  272. Barennes H, Andriatahina T, Latthaphasavang V, *et al.* (2008) Misperceptions and misuse of Bear Brand coffee creamer as infant food: national cross sectional survey of consumers and paediatricians in Laos. *BMJ* **337**, a1379.