

The 14 November 2001, $M_w = 7.8$ Kokoxili Earthquake in Northern Tibet (Qinghai Province, China)

J. Van der Woerd

Institut de Physique du Globe, Strasbourg

A.-S. Mériaux, Y. Klinger

Laboratoire de Tectonique, Institut de Physique du Globe, Paris

F.J. Ryerson

Institute of Geophysics and Planetary Physics, Lawrence Livermore National Laboratory

Y. Gaudemer, and P. Tapponnier

Laboratoire de Tectonique, Institut de Physique du Globe, Paris

INTRODUCTION

On 14 November 2001, at 5:26 PM local time (9:26:10.3 GMT) a great earthquake ($M_s=8.1$) struck the Kokoxili region of the northwest part of Qinghai province, China (Figure 1). Although this earthquake was felt in Sichuan province, more than 1,000 km away from the epicentral area, it did not attract much media attention because it shook a thinly populated area on the northern margin of the Tibetan Plateau. The preliminary epicenter (U.S. Geological Survey-National Earthquake Information Center [NEIC]) is located near 90.5°E, about 300 km west of the Golmud-Lhasa road (Figures 2 and 3), across which surface rupture has been reported near the Kunlun Pass. As mentioned in several Chinese newspapers, the famous milestone statue at the Kunlun Pass broke and fell because of the earthquake and the construction of the Golmud-Lhasa railroad has been put on hold since the occurrence of this unexpected event.


The earthquake appears to have ruptured the left-lateral Kunlun Fault (Tapponnier and Molnar, 1977) or Kusai-Dongxi-Maqu Fault (Cui and Jiang, 1979; Jia *et al.*, 1988), which extends roughly east-west for about 1,600 km in north Tibet. This event follows, nearly four years later, the 8 November 1997, $M_w=7.6$ Manyi earthquake, which broke another segment of the same active fault system farther west (Figure 2).

The centroid (Harvard CMT Catalog), at 35.5°N 92.7°E, lies about 200 km east of the USGS (NEIC) and

China Seismological Bureau (CSB) preliminary epicentral locations, and about 35 km south of the fault, consistent with a south-dipping fault-plane (Figure 3). Eighteen of the twenty-two aftershocks with magnitude $M_b \geq 3.8$ that occurred during the first week are located another 100 km farther east, between 93°E and 94.7°E. The easternmost ones are located east of the Kunlun Pass. Two focal mechanisms, one from the Harvard CMT Catalog, and the other, from the Earthquake Information Center in Tokyo (Earthquake Research Institute ERI, Tokyo), indicate pure left-lateral to normal-left-lateral motion on a N90-96°E striking plane, compatible with the average N95-100°E strike of the Kunlun Fault in the area. The USGS fault-plane solution, on the other hand, appears to be radically different, but would not be inconsistent with slip on a steep, N20°E-trending normal fault plane (figure 3).

Shortly after the earthquake, the Chinese Seismological Bureau deployed about ten temporary stations at the eastern end of the rupture, along the Golmud to Lhasa road. The earthquake was also recorded by a set of ten temporary stations deployed since June 2001 for a passive seismologic experiment in western Tibet, about 1,000 km to the west (IT program, INSU, CNRS France). Unfortunately, these stations had to be moved away four days after the main shock.

In this report, after describing the Kunlun Fault and what is known of previous events along it, we discuss the rupture length, the positions of the largest aftershocks and possible triggering scenarios for future earthquake occurrence.




▲ Figure 1. Map of the active faults of Tibet, after Tapponnier et al. (2001a).

TECTONIC SETTING


Since about 50 Ma, the continued northward convergence of India relative to Siberia has created the largest continental collision zone on the planet. At present, out of the ~ 5 cm/yr shortening between the two continents, ~ 2 cm/yr are absorbed by the Himalayan frontal thrust (Lavé and Avouac, 2000; Bilham et al., 1997) resulting in the growth of the world's highest mountain range. Most of the remaining convergence appears to be absorbed along Tibet's northern margin, by extrusion of lithospheric blocks along great strike-slip faults combined with oblique thrusting and crustal thickening (Meyer et al., 1998; Tapponnier et al., 2001a). The Altyn Tagh and Haiyuan faults, north of the Qaidam basin, as well as the Karakorum, and the Kunlun and Xian Shui He faults, on the western and eastern side of Tibet, respectively, allow the blocks to move past each other at long-term rates on order of a few centimeters per year (Figure 1) (Avouac and Tapponnier, 1993; Lasserre et al., 1999; Mériaux et al., 2000b; Van der Woerd et al., 2000).

The Kunlun Fault is not only one of the faults that takes part in the extrusion of Tibet but also an example of large-scale slip-partitioning in the continental crust (e.g., Meyer et al., 1998; Tapponnier et al., 2001a). Northeastward convergence between Tibet and the Gobi, guided by left-lateral slip on the Altyn Tagh Fault, has led to crustal-shortening of the Qaidam basin and Qilian ranges, apparently above a décollement that roots underneath the Kunlun range, where the lithospheric mantle of the latter regions subducts southwards (Tapponnier et al., 1990; Meyer et al., 1998; Chen et al., 1999). The orientation of the Kunlun fault, which is parallel to the Kunlun range, and the strong east-west shear-wave splitting anisotropy across it (McNamara et al., 1994; Lavé, 1996; Herquel et al., 1995) suggest that the fault extends deeper in the lithosphere as a shear zone, as does the Altyn Tagh Fault (Wittlinger et al., 1998).

For more than 1,000 km, the Kunlun Fault runs south of the Kunlun range along and north of the most recent (12 Ma to present), calcalkaline volcanic belt of Tibet (Turner et al., 1996; Mériaux et al., 2000b; Jolivet et al., 2001). The range, an active margin in the Permo-Triassic, separates the folded



▲ **Figure 2.** Simplified large-scale map of the Kunlun Fault and of its first-order segmentation (after Van der Woerd, 1998).



▲ **Figure 3.** Map of Kusai Hu segment of the Kunlun Fault with location of main shock epicenters and centroid, and of aftershocks with magnitude 4. Fault planes solutions of main shock and of M 5.5 aftershocks and previous events since 1980 are also shown. Maximum extent of surface rupture is outlined. Inset displays source modelling results redrawn from Kikuchi and Yamanaka (2001) (ERI, Tokyo), centered on Harvard CMT. Scale for slip is in meters; maximum slip is 5 meters.

turbidites and phyllites of the Songpan Ganze terrane of eastern Tibet (Chang *et al.*, 1986) from the Paleozoic and lower Proterozoic basement of the Qaidam to the north. The timing of Cenozoic motion along the fault is not well known. It has probably been slipping since at least the Mid-Miocene (12 Ma) coevally with shortening to the north, which started also in the Miocene (*e.g.*, Meyer *et al.*, 1998; Métivier *et al.*, 1998), and with volcanism to the south (Deng, 1991; Arnaud *et al.*, 1992; Turner *et al.*, 1996). However, it may have started to slip earlier, in the late Oligocene, as shortening driven by the collision propagated northeastwards into Asia (Tapponnier *et al.*, 2001a).

The fault stretches from 86°E to 105°E (about 1,600 km), from the borders of Xinjiang, Xizang and Qinghai provinces, across the dry permafrost plateau (~5000 m. a.s.l.) down to the humid, forested and deeply incised mountainous rim of the Sichuan basin (Figures 1 and 2). From cosmogenic dating (¹⁴C, ¹⁰Be, ²⁶Al) of offset alluvial terraces along the fault, the long-term slip-rate along much of the Kunlun Fault has been determined to be 11.5 ± 2.0 mm/yr (Van der Woerd *et al.*, 1998; 2000; 2002). East of the Kunlun Pass, coseismic offsets of channels and terraces risers have been interpreted to result from the occurrence of large earthquakes (7.5 M 8) with recurrence times of about 900 years in the west, and 450 years in the east (Van der Woerd *et al.*, 2000; 2002).

KUSAI HU SEGMENT OF THE KUNLUN FAULT

The main trace of the Kunlun fault may be divided into six principal segments separated by first-order bends, jogs or junctions (Figure 2). The 14 November 2001 earthquake ruptured the western, Kusai Hu segment of the fault, named from the large lake that it crosses in its central part (Figure 3). This segment, though arguably the most remote and least well known along the main stretch of the fault, is one of its clearest and longest (270 km) (Figures 3 and 4). Between ~91°E and 94°E it strikes N100E, which is close to the average strike of the entire fault system. West of 91°E, the fault splays into an extensional horsetail with several oblique, normal-sinistral strands whose strikes range between N20°E and N120°E (Mériaux *et al.*, 2000b). The fault traces, which are a few ten to a few hundred kilometers long, are well marked across the Quaternary bajadas of the plateau. One of the longest is the Manyi fault, which extends for 270 km between 86.5°E and 89.5°E with a N80°E average strike (Tapponnier and Molnar, 1977; Xu *et al.*, 1999).

Although the Kusai Hu segment of the Kunlun fault is remarkably straight between 91°E and 94°E, a progressive change in strike, from N100°E in the west to N95°E in the east, induces slightly different styles of faulting, from oblique normal-strike-slip, to nearly pure strike-slip and oblique reverse-strike-slip faulting.

Near 91°E, one oblique normal strand appears to be chiefly responsible for the uplift of the Buka Daban Feng, highest peak of the eastern Kunlun range, at 6,800 m asl. This


strand bounds the steep, southwest-trending southern flank of Buka Daban, at the foot of which large glaciers pond between faceted spurs (Figure 4A). East of 91°E, slip partitioning occurs between two parallel, normal and pure left-lateral, strands. Normal faulting takes place at the base of south facing triangular facets, along the Kunlun range front, while sinistral slip occurs 1 to 2 km southwards, across the coalescent fans of the piedmont bajadas (Figures 4C and 4D), a geometry often found along oblique strike-slip faults in Tibet and adjacent areas (Armijo *et al.*, 1986; Van der Woerd, 1992; Allen *et al.*, 1984; Replumaz *et al.*, 2001). This partitioning implies that the Kunlun Fault dips southwards. At a more detailed level, the bajada trace is divided into three right-stepping sub-segments that have lengths of ~30, 75 and 60 km. Some of the most spectacular, clean-cut sinistral offsets of terrace risers and channels incised into post-glacial fans and terraces are visible on SPOT images of the fault trace, west of Kusai Hu (Figure 4C). These offsets typically range between 25 and 60 meters, implying that, given the ~11 mm/yr rate, they postdate the end of the early Holocene climatic optimum (~6,000 yr). East of Kusai Hu, the fault follows the range front more closely. Yet farther east, past a left-step at 93.6°E, the fault splays into two distinct strands (Figure 4B). The northern one continues eastwards as the Xidatan-Dongdatan segment of the main fault. The southern one connects with the oblique, sinistral-reverse Kunlun Pass Fault (Kidd and Molnar, 1988; Van der Woerd *et al.*, 2002) which bounds the steep southern face of the Burhan Budai range.

HISTORICAL AND RECENT SEISMICITY


The north central Tibetan plateau is such a remote and sparsely inhabited area that information about earthquakes is nonexistent, to our knowledge, before 1900. No paleoseismological evidence has been published yet in the western literature. The long-term seismic behavior of the Kunlun Fault is thus unknown. Since 1900, however, several earthquakes with M_s 7 have occurred along the fault and related surface structures have been described (Gu *et al.*, 1989).

Far east of Burhan Budai Shan, a large event (35.5°E-97.5°E) with an estimated magnitude $M \sim 7.5$ ruptured most of the Dongxi Co segment of the fault on 7 January 1937 (Jia *et al.*, 1988). The epicenter is located approximately at the western end of this segment and the rupture clearly extended eastwards. We mapped and measured fresh free faces, large mole tracks, and sinistral offset rills (~4 m) east of Dongxi Co that are unambiguously due to this event (Van der Woerd, 1998; 2002). The 1937 rupture, which crosses the Dongxi Co pull-apart is thus at least 150 km long. Our field observations, however, indicate that it did not cross the Qinglong river, east of Xiadawu. The length obtained is in good agreement with the estimated magnitude of 7.5, according to magnitude/rupture length scaling laws (Wells and Coppersmith, 1995). No focal mechanism has been determined for the 1937 event, but it is predominantly sinistral strike-slip in the field (Figure 5).



A



B




▲ **Figure 4.** Satellite image views of specific sites along the Kusai Hu segment of the fault.
 (A) Landsat view of western part of Kusai Hu segment, showing splaying of Kunlun fault west of Buka Daban Feng.
 (B) SPOT mosaic of eastern part of Kusai Hu segment and of splay between Xidatan segment and Kunlun Pass fault.



▲ **Figure 4.** Satellite image views of specific sites along the Kusai Hu segment of the fault.

(C) SPOT image enlargement of terrace and stream offsets due to cumulative left-lateral displacement along the fault. This is the area shown on 15 August 2000, *Geophysical Research Letters* cover.

(D) SPOT image enlargement of pull-apart and large offset alluvial fan along left-lateral strand of fault, south of normal fault bounding faceted spurs of East Kunlun range. Both faults, only 1-2 km apart, probably merge at depth, testifying to local transtensional kinematics on south-dipping, oblique fault in basement beneath sediments.



▲ **Figure 5.** View, looking west, of eroded moles tracks of 1937 earthquake east of Dongxi lake.

A second earthquake with a magnitude of about $M \sim 7$ occurred about 100 km westward of the 1937 event epicenter, on 19 April 1963. It probably ruptured part of the Alag Hu segment of the fault. Fitch (1970) determined the focal mechanism to be left-lateral strike-slip. Molnar and Deng (1984), quoting reported displacements of several meters (Li and Jia, 1981), suggested that the total rupture length for the two events of 1937 and 1963 might have reached about 300 km, but this is likely overestimated. In 1971, a smaller earthquake with magnitude $M \sim 6.4$ occurred at the junction between the Alag Hu and Dongxi Co segment of the fault. The focal mechanism is nearly pure left-lateral strike-slip.

Prior to 14 November 2001, no large earthquake had been documented along the Kusai segment of the fault. One significant event, however, was the $M \sim 5.8$, 6 March 1980 earthquake north of the western stretch of that segment, apparently on the central Kunlun fault. The focal mechanism of this event shows left-lateral strike-slip with a thrust component (Harvard CMT). Another significant event was the $M \sim 5.8$, 3 January 1994 earthquake, located southwest of Buka Daban Feng, hence southwest of the western termination of the Kusai segment. This latter event had a normal fault plane solution with NE-striking nodal planes, consistent with the oblique normal fault strands observed in the area.

Farther west of Buka Daban Feng, the Manyi Fault, and faults west of it, have produced two large earthquakes during the latter part of the 20th century. On 14 July 1973, two

earthquakes of magnitude $M \sim 7.3$ and $M \sim 6.1$ (NEIC), at the western end of the Manyi segment, near $35.25^{\circ}\text{N} - 86.5^{\circ}\text{E}$, ruptured an oblique, normal sinistral fault striking northeast-southwest (Tapponnier and Molnar, 1977) (Figure 2). On 8 November 1997, the Manyi earthquake, of magnitude $M_w \sim 7.6$, produced a 170-km-long surface break with maximum offsets on order of 7 m (Peltzer *et al.*, 1999). Both the InSAR results and the focal mechanism (NEIC, Harvard CMT) indicate pure left-lateral strike-slip on a Quaternary fault mapped on Landsat imagery (Tapponnier and Molnar, 1977). The mainshock was followed by a series of aftershocks that roughly lined up with the surface rupture (Velasco *et al.*, 1998; Xu *et al.*, 1999).

MAIN SHOCK AND AFTERSHOCK SEQUENCE

Length of the 14 November 2001 Earthquake Rupture

Preliminary locations of the epicenter of the 14 November 2001 earthquake, of estimated magnitude M_s 7.9 to 8.1 (NEIC; Chinese Seismological Bureau, 2001, Li Haibing, personal comm.) place it at $36^{\circ}\text{N} 90.5^{\circ}\text{E}$, near the highest summits of the Eastern Kunlun Range (Buka Daban Feng and Wei Xue Shan, $\sim 6,800$ m and $\sim 6,000$ m, respectively) at the western termination of the Kusai Hu segment of the fault. The centroid, on the other hand, is located at $35.5^{\circ}\text{N} 92.7^{\circ}\text{E}$, about 200 km eastward near lake Kusai. The source duration of the event was ~ 120 seconds. Following a weak initial pulse in the west the bulk of the energy was dissipated eastward in



▲ **Figure 6.** View of fresh mole tracks due to 14 November 2001 $M_w=7.8$ earthquake near Kunlun Pass, at eastern end of rupture (courtesy of Xinhua press). Several meters of sinistral slip are required to produce the cracks and pressure ridges that tear the frozen ground.

about 80 seconds (ERI, Tokyo). It is thus likely that the rupture started in the west, possibly on an oblique normal fault near Buka Daban Feng and propagated eastward along much of the length of the Kusai segment (~ 270 km) with most of the seismic energy radiated along the main strike-slip strand of the fault and a maximum left-slip in excess of 5 m (Figure 3, inset). Since surface breaks are seen east of the Golmud-Lhasa road near the Kunlun Pass, with probably still meters of sinistral slip (Figure 6) apparently along the Kunlun Pass Fault, the rupture length may have exceeded that of the

Kusai Hu segment, unless the westernmost part of that segment was spared by the 14 November 2001 earthquake. If the rupture length was in excess of 270 - 300 km, then it would rank second only to that of the 1905 Bolnai earthquake in Mongolia (> 350 km) (Khilko *et al.*, 1985), and make the Kokoxili earthquake one of the largest strike-slip event in Asia during the last ~ 100 yr, ahead of even the 1920 Haiyuan earthquake (230 km) in Gansu province (Deng *et al.*, 1986; Gaudemer *et al.*, 1995), or the 1957 Gobi Altai earthquake (260 km) in Mongolia (Baljinnyam *et al.*, 1993).

Aftershock sequence

Between 14 November and 8 December, twenty-two aftershocks with magnitude greater than 3.8 occurred, eleven of them in the day (15 hours) following the main shock, and seven in the next five days. Most of the aftershocks occurred east of 93°E, extending as far east as 94.8°E (NEIC), close to the Kunlun Pass splay, in the general area where the rupture likely stopped (Figure 3). Two of the fault-plane solutions of the eastern aftershocks have thrust mechanisms, which is consistent with an east-propagating rupture stopping near 94°N. The distance from the NEIC epicenter to the easternmost aftershocks reaches 350 km. Even with a maximum uncertainty of 50 km in the longitudinal location of these events, this distance would still be 250 km. This corroborates the inference that the entire Kusai Hu segment ruptured during the 14 November, 2001 main shock. Preliminary field investigations (Xu Xiwei, personal comm.), confirm that the rupture crosses the Kunlun Pass road at 94.05°E, to continue eastward, south of the Burhan Budai Shan. No breaks, on the other hand, have been described so far along the trace of the Kunlun Fault in Xidatan, implying that the next 160-km-long segment of the fault toward the east (Van der Woerd *et al.*, 1998; 2002) was not involved in the recent event.

POSSIBLE TRIGGERING OF FUTURE LARGE EARTHQUAKES

The seismicity in the last 100 years is certainly not representative of the long-term behavior of the Kunlun Fault. However, the fact that since 1937 several segments of the fault system have ruptured with earthquakes of magnitude greater than 7 implies that the whole fault is in an active period of its seismic cycle. Other segments, along which no large earthquake has occurred in the last 100 years, may thus be ready to break in the next future (Xidatan Dongdatan, Alag Hu, or Maqen segment farther east, Figure 2).

Based on the 11.5 mm/yr slip-rate and co-seismic, apparently characteristic, displacements of 8-12 m determined along the Xidatan segment, we deduced a recurrence time for large earthquakes rupturing that segment of about 900 years (Van der Woerd *et al.*, 2000; 2002; Tapponnier *et al.*, 2001b). At one site, the last large earthquake mole tracks on the lowest terrace were clearly washed out by a flood dated (with cosmogenic ¹⁰Be and ²⁶Al) to be 278 ± 87 years old (Van der Woerd *et al.*, 1998; 2002). Trenching at different sites along the segment (Zhao *et al.*, 1994; Zhao, 1996), further corroborates the inference that no large earthquake ruptured the youngest alluvial terrace and fan surfaces along the Xidatan segment in the last 500 years.

The clustering of aftershocks, some of them with thrust fault plane solutions, at the eastern end of the rupture, near and east of the Kunlun Pass, implies that the occurrence of the earthquake induced fairly large stress changes in the region encompassing the western end of the Xidatan Dongdatan segment of the fault. This segment has not broken in the last 500 years, and only two large earthquakes have offset, by ~24 ± 3 m,

a riser dated at 1,788 ± 388 years at one site in the middle of Xidatan, near 35.4°N 94.4°E (Van der Woerd *et al.*, 1998, 2002). Such observations are not inconsistent with the inference that we might be close to the end of the average recurrence interval of large earthquakes on that part of the Kunlun fault. If this were the case, then the Coulomb stress concentration induced by the 14 November 2001 event in the Kunlun Pass area could trigger another large earthquake with magnitude greater than 7.5 on the Xidatan Dongdatan segment of the Kunlun Fault in the next months or years. Such sequential triggering of earthquakes would not be surprising. It has been observed along other major strike slip-faults, most remarkably the North Anatolian Fault in Turkey. Since 1939, nearly all the principal segments of that fault have ruptured successively from East to West, each $M > 7$ earthquake triggering the next, months or decades later, in cascade mode (e.g., Stein *et al.*, 1997). As a result, the whole length of the North Anatolian Fault between Erzinjan and Izmit (~1200 km) slipped in 60 years.

The possibility of similar earthquake triggering along the Kunlun fault, possibly all the way to Alag Hu, makes it particularly urgent to instrument and study the region located east of the Kunlun Pass in detail, with the various techniques that are now commonly used to understand post- and preseismic deformation and related phenomena. ■

ACKNOWLEDGMENTS

The results of the research reported here was supported by Institut National des Sciences de l'Univers (CNRS, France) and the Ministry of Geology and Mineral Resources (Chinese Academy of Geological Sciences, Beijing, China). We thank Xu Xiwei for sharing some observations gathered in the field a few days after the earthquake, and Li Haibing for retrieving further information in Chinese. This is EOST-IPGS contribution No. 2002-001 and IGP contribution No. 1802.

REFERENCES

- Allen, C.R., Luo Zhuoli, Qian Hong, Wen Xueze, Zhou Huawei et Huang Weishi (1991). Field study of a highly active fault zone: The Xianshuihe fault of southwestern China, *Geol. Soc. Am. Bull.*, **103**, 1,178-1,199.
- Armijo, R., P. Tapponnier, J.L. Mercier et Han Tong Lin (1986). Quaternary extension in southern Tibet: Field observations and tectonic implications, *J. Geophys. Res.*, **91**, 13,803-13,872.
- Arnaud, N.O., P. Vidal, P. Tapponnier, P. Matte and W.M. Deng (1992). The high K²O volcanism of northwestern Tibet; geochemistry and tectonic implications, *Earth planet. Sci. Lett.*, **111**, 351-367.
- Avouac J.P., and P. Tapponnier (1993). Kinematic model of active deformation in central Asia, *Geophys. Res. Lett.*, **20**, 895-898.
- Bilham, R., K. Larson, J. Freymueller and Project Idylhim members (1997). GPS measurements of present-day convergence across the Nepal Himalaya, *Nature(Lond.)*, **386**, 61-64.
- Baljinnyam, I., Bayasgalan, A., Borisov, B.A., Cisternas, A., Dem'yanovich, M.G., Ganbaatar, L., Kochetov, V.M., Kurushin, R.A., Molnar, P., Philip, H., and Vasilov, Y.Y. (1993). *Ruptures of Major Earthquakes and Active Deformation in Mongolia and its Surroundings*, *Geol. Soc. Am. Memoir* 181, 62 pp.
- Cui Zhengyuan and Yang Bin (1979). On the Tuosuohu-Maqu active fault zone, *Northwest. Seism. J.*, **1**, 57-61 (in Chinese).
- Chang, C., N. Chen, M.P. Coward, W. Deng, J.F. Dewey, A. Gansser, N.B.W. Harris, C. Jin, W.S.F. Kidd, M.R. Leeder, H. Li, J. Lin, C.

- Liu, H. Mei, P. Molnar, Y. Pan, Y. Pan, J.A. Pearce, R.M. Shackleton, A.B. Smith, Y. Sun, M. Ward, D.R. Watts, J. Xu, R. Xu, J. Yin, and Y. Zhang (1986). Preliminary conclusions of the Royal Society and Academia Sinica 1985 geotraverse of Tibet, *Nature*, **323** (6,088), 501-507.
- Chen W.P., C.Y. Chen, and J. Nábelek (1999). Present-day deformation of the Qaidam basin with implications for intra-continental tectonics, *Tectonophysics*, **305**, 165-181.
- Coward, M.P., W.S.F. Kidd, Pan Yun, R.M. Shackleton, F.R.S. and Zhang Hu (1988). The structure of the 1985 Tibet Geotraverse, Lhasa to Golmud, *Phil. Trans. R. Soc. London*, **A327**, 307-336.
- Deng, Q., S. Chen, S. Zhu, Y. Wang, W. Zhang, D. Jiao, B.C. Buchfield, P. Molnar, L.H. Royden, and P. Zhang (1986). Variation in the geometry and amount of slip on the Haiyuan fault zone, China, and the surface rupture of the 1920 Haiyuan earthquake, in *Earthquake Source Mechanics*, *Geophys. Monogr. Ser.* **37**, 169-182.
- Deng, W. (1991). Cenozoic volcanism and intraplate subduction in northern margin of the Tibet plateau, *Chin. J. Geochem.*, **10**, 140-152.
- Fitch, T.J. (1970). Earthquake mechanisms in the Himalayan, Burmese, and Adaman regions and continental tectonics in central Asia, *J. Geophys. Res.*, **75**, 2699-2709.
- Gaudemer Y., P. Tapponnier, B. Meyer, G. Peltzer, Guo Shunmin, Chen Zhitai, Dai Huagang and I. Cifuentes (1995). Partitioning of crustal slip between linked, active faults in the eastern Qilian Shan, and evidence for a major seismic gap, the « Tianzhu gap », on the Western Haiyuan fault, Gansu (China), *Geophys. J. Int.*, **120**, 599-645.
- Gu, G., L. Ting Hong, and S. Zhen Jiliang (1989). *Catalogue of Chinese Earthquakes (1831 BC-1969 AD)*, Science Press, Beijing, China.
- Herquel, G., G. Wittlinger and J. Guilbert (1995). Anisotropy and crustal thickness of Northern-Tibet. New constraints for tectonic modelling, *Geophys. Res. Lett.*, **22**, 1,925-1,928.
- Jia, Y., H. Dai and X. Su (1988). Tuosuo Lake Earthquake Fault in Qinghai Province, in *Research on Earthquake Faults in China*, Xinjiang Seismological Bureau, Xinjiang Press, Wulumuqi, 66-71.
- Jolivet, M., Brunel, M., Seward, D., Xu, Z., Yang, J., Tapponnier, P., Malavieille, J., Roger, F., Leyrelop, A., Mériaux, A.-S., Crambes, C., Arnaud, N. and Wu, C., (2001). Neogene extensional features and Quaternary volcanism associated with the Kunlun fault, *Geophys. Res. Abs.* **3**, EGS 26th General Assembly, Nice.
- Kidd, W.S.F. and P. Molnar (1988). Quaternary and active faulting observed on the 1985 Academia Sinica-Royal Geotraverse of Tibet, *Phil. Trans. R. Soc. London*, **A327**, 337-363.
- Kikuchi, M., and Y. Yamanaka (2001). Earthquake Information Center, Earthquake Research Institute, University of Tokyo, Japan, [website url: http://www.eri.u-tokyo.ac.jp/topics/200111140926/](http://www.eri.u-tokyo.ac.jp/topics/200111140926/)
- Khilko, S.D., R.A. Kurushin, V.M. Kochetkov, L.A. Misharina, V.I. Melnikova, N.A. Gilyova, S.V. Lastochkin, I. Balzhinnyam, and D. Monhoo (1985). Earthquakes and the bases of the seismic zoning of Mongolia, vol. 41 of the joint Soviet-Mongolian scientific - Research Geological Expedition.
- Lasserre, C., P. H. Morel, Y. Gaudemer, P. Tapponnier, F. J. Ryerson, G. King, F. Métivier, M. Kasser, M. Kashgarian, B. Liu, T. Lu, and D. Yuan (1999). Post-glacial left slip-rate and past occurrence of M 8 earthquakes on the western Haiyuan fault (Gansu, China), *J. Geophys. Res.*, **104**, 17,633-17,651.
- Lavé, J., J.P. Avouac, R. Lacassin, P. Tapponnier and J.P. Montagner (1996). Seismic anisotropy beneath Tibet: evidence for eastward extrusion of the Tibetan lithosphere, *Earth planet. Sci. Lett.*, **140**, 83-96.
- Lave J. and J.P. Avouac (2000). Active folding of fluvial terraces across the Siwaliks Hills, Himalayas of central Nepal, *J. Geophys. Res.*, **105**, 5,735-5,770.
- Li Longhai and Jia Yunhong (1981). Characteristics of the deformation band of the 1937 Tuosuo Lake earthquake (M=7.5) in Qinghai, *Northwest. Seism. J.*, **3**, 61-65.
- McNamara, D.E., T.J. Owens, P.G. Silver and F.T. Wu (1994). Shear wave anisotropy beneath the Tibetan Plateau, *J. Geophys. Res.*, **99**, 13,655-13,665.
- Mériaux A.-S., F.J. Ryerson, P. Tapponnier, J. Van der Woerd, R.C. Finkel, M.W. Caffee, Xu Zhiqin, (2000a). Tectonique et Volcanisme actifs au Nord Tibet, *18e Réunion des Sciences de la Terre*, Soc Géol. Fr. édit. Paris, 294.
- Mériaux, A.-S., P. Tapponnier, F.J. Ryerson, J. Van der Woerd, C. Lasserre, Xu Xiwei, Li Haibing, Xu Zhiqin, R.C. Finkel and M.W. Caffee (2000b). Fast Extrusion of the Tibet Plateau: a 3cm/yr, 100Kyr Slip-Rate on the Altyn Tagh Fault, *Eos Trans. Am. Geophys. U.*, **81**, 1,137.
- Métivier, F., Y. Gaudemer, P. Tapponnier and B. Meyer (1998). Northeastward growth of the Tibet plateau deduced from balanced reconstruction of two depositional areas: The Qaidam and Hexi Corridor basins, China, *Tectonics*, **17**, 823-842.
- Meyer, B., P. Tapponnier, L. Bourjot, F. Métivier, Y. Gaudemer, G. Peltzer, G. Shunmin and C. Zhitai (1998). Mechanisms of active crustal thickening in Gansu-Qinghai, and oblique, strike-slip controlled, northeastward growth of the Tibet plateau, *Geophys. J. Int.*, **135**, 1-47.
- Molnar, P. and Deng Qidong (1984). Faulting associated with large earthquakes and the average rate of deformation in central and eastern Asia, *J. Geophys. Res.*, **89**, 6203-6227.
- Peltzer, G., F. Crampé, and G. King (1999). Evidence of Nonlinear Elasticity of the Crust from the Mw 7.6 Manyi (Tibet) Earthquake, *Science*, **286**, 272.
- Replumaz A., R. Lacassin, P. Tapponnier and P.H. Leloup (2001). Large river offset and Plio-Quaternary dextral slip rate on the Red River fault (Yunnan, China), *J. Geophys. Res.*, **106**, 819-836.
- Stein R., A. Barka, and J. Dietrich (1997). Progressive failure on the North Anatolian fault since 1939 by earthquake stress triggering, *Geophys. J. Int.*, **128**, 594-604.
- Tapponnier, P. and P. Molnar (1977). Active faulting and tectonics in China, *J. Geophys. Res.*, **82**, 2,905-2,930.
- Tapponnier, P., B. Meyer, J.P. Avouac, G. Peltzer, Y. Gaudemer, Guo Shunmin, Xiang Hongfa, Yin Kelun, Chen Zhitai, Cai Shuahua and Dai Huagang (1990). Active thrusting and folding in the Qilian Shan, and decoupling between upper crust and mantle in northeastern Tibet, *Earth planet. Sci. Lett.*, **97**, 382-403.
- Tapponnier, P., Xu Zhiqin, F. Roger, B. Meyer, N. Arnaud, G. Wittlinger, and Yang Jingsui, (2001a). Oblique stepwise rise and growth of the Tibet plateau, *Science*, **294**, 1,671.
- Tapponnier, P., F.J. Ryerson, J. Van der Woerd, A.S. Mériaux, C. Lasserre (2001b). Long-term slip rates and characteristic slip: keys to active fault behavior and earthquake hazard, *C. R. Acad. Sci. Paris*, **333**, 483-494.
- Turner, S., N. Arnaud, J. Liu, N. Rogers, C. Hawkesworth, N. Harris, S. Kelley, P. Van Calsteren and W. Deng (1996). Post-collision, shoshonitic volcanism on the Tibetan Plateau : Implications for convective thinning of the lithosphere and the source of ocean island basalts, *J. Petrol.*, **37**, 45-71.
- Van der Woerd, J. (1992). Décalages de la morphologie glaciaire et post-glaciaire observés sur les images du satellite SPOT le long de la faille du Kunlun, DEA report, Institut de Physique du Globe de Paris, 24pp.
- Van Der Woerd, J., F.J. Ryerson, P. Tapponnier, Y. Gaudemer, R. Finkel, A.S. Mériaux, M. Caffee, Zhao Guoguang and He Qunlu (1998). Holocene left slip-rate determined by cosmogenic surface dating on the Xidatan segment of the Kunlun fault (Qinghai, China), *Geology*, **26**, 695-698.
- Van der Woerd, J., F.J. Ryerson, P. Tapponnier, A.-S. Mériaux, Y. Gaudemer, B. Meyer, R.C. Finkel, M.W. Caffee, Zhao Guoguang, Xu Zhiqin (2000). Uniform Slip-Rate long the Kunlun Fault: Implications for seismic behaviour and large-scale Tectonics, *Geophys. Res. Lett.*, **27**, 2,353-2,356.

- Van der Woerd, J., P. Tapponnier, F.J. Ryerson, A.-S. Mériaux, B. Meyer, Y. Gaudemer, R.C. Finkel, M.W. Caffee, Zhao Guoguang, X. Zhiqin (2002). Uniform Post-Glacial slip-rate along the central 600 km of the Kunlun Fault (Tibet), from ^{26}Al , ^{10}Be and ^{14}C dating of riser offsets, and climatic origin of the regional morphology, *Geophys. J. Int.*, (in press).
- Velasco, A.A., C.J. Amon and S. Beck (2000). Broadband Source Modeling of the November 8, 1997 Tibet, (Mw=7.5) Earthquake and its Tectonic Implications, *J. geophys. Res.*, **105**, 28,065-28,028.
- Wells, D. and K. Coppersmith, (1995). New empirical relationship among magnitude, rupture width, rupture area, and surface displacement, *Bull. Seism. Soc. Am.*, **84**, 974-1,002.
- Wittlinger, P., Tapponnier, G. Poupinet (1998). Tomographic evidence for lithospheric shear along the Altyn Tagh Fault (China), *Science*, **282**, 74.
- Xu Xiwei et al. (1999). *Manyi earthquake*, report from China Seismological Bureau.
- Zhao Guoguang (1996). Quaternary faulting in north Qinghai-Tibet plateau, *Continental dynamics*, **1**, 30-37.
- Zhao Guoguang, He Qunlu, P. Tapponnier and Y. Gaudemer (1994). Preliminary investigation on the Holocene paleoearthquake features along the mid-section of the Ku-ma

active fault, East Kunlun, in Proceedings of First Symposium on Geology and Geophysics of East Kunlun and its adjacent areas, *Continental Geodynamics*, Beijing, China.

*Institut de Physique du Globe
Strasbourg
France
jerome.vanderwoerd@east.u-strasbg.fr
(J.V.D.W.)*

*Laboratoire de Tectonique
Institut de Physique du Globe
Paris
France
(A.-S.M., Y.K., Y.G., P.T.)*

*Institute of Geophysics and Planetary Physics
Lawrence Livermore National Laboratory
Livermore, CA
USA
(F.J.R.)*