



HAL
open science

New analytical planetary theories VSOP2013 and TOP2013

J.-L. Simon, G. Francou, A. Fienga, H. Manche

► **To cite this version:**

J.-L. Simon, G. Francou, A. Fienga, H. Manche. New analytical planetary theories VSOP2013 and TOP2013. *Astronomy and Astrophysics - A&A*, EDP Sciences, 2013, 557, pp.A49. 10.1051/0004-6361/201321843 . hal-01856071

HAL Id: hal-01856071

<https://hal.archives-ouvertes.fr/hal-01856071>

Submitted on 2 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License

New analytical planetary theories VSOP2013 and TOP2013[★]

J.-L. Simon¹, G. Francou², A. Fienga^{3,1}, and H. Manche¹

¹ IMCCE, Observatoire de Paris, UMR CNRS 8028, UPMC, 75014 Paris, France
e-mail: jlsimon@imcce.fr

² SYRTE, Observatoire de Paris, UMR CNRS 8630, UPMC, 75014 Paris, France
e-mail: gerard.francou@obspm.fr

³ GéoAzur, Observatoire de la Côte d'Azur, UNR CNRS 7329, 04870 Côte d'Azur, France
e-mail: agnes.fienga@oca.eu

Received 6 May 2013 / Accepted 19 June 2013

ABSTRACT

Context. The development of precise numerical integrations of the motion of the planets, taking into account the most recent observations, lead us to improve the two families of analytical planetary theories built in the Institut de mécanique céleste et de calcul des éphémérides (IMCCE), the Variations Séculaires des Orbites Planétaires (VSOP) and the Theory of the Outer Planets (TOP) theories. **Aims.** We have built the solutions VSOP2010 and TOP2010 fitted to the Jet Propulsion Laboratory (JPL) numerical integration DE405 and the solutions VSOP2013 and TOP2013 fitted to the European recent numerical integration INPOP10a. This paper specifically considers VSOP2013 and TOP2013.

Methods. We have improved the construction of VSOP by analytically computing the perturbations due to the asteroids and to Pluto. We have increased the precision of the VSOP solutions of Jupiter and Saturn by using TOP solutions. We have also improved the construction of TOP by computing the perturbations due to the telluric planets from VSOP solutions. Moreover, TOP contains a solution of the motion of the Pluto-Charon barycenter.

Results. From 1890 to 2000, the precision of VSOP2013 goes from a few 0.01 mas (planets except Mars and Uranus) up to 0.7 mas (Mars and Uranus). Compared to the previous solution (VSOP2000), this represents an improvement of a factor of 2 to 24, depending on the planet. From -4000 to 8000, the precision is of a few 0.1'' for the telluric planets (1.6'' for Mars), i.e. an improvement of about a factor of 5 compared to VSOP2000. The TOP2013 solution is the best for the motion of the major planets from -4000 to 8000. Its precision is of a few 0.1'' for the four planets, i.e. a gain between 1.5 and 15, depending on the planet compared to VSOP2013. The precision of the theory of Pluto remains valid up to the time span from 0 to 4000. The VSOP2013 and TOP2013 data are available on the WEB server of the IMCCE.

Key words. celestial mechanics – ephemerides

1. Introduction

For the last 30 years, two kinds of analytical planetary theories¹ have been built in the Institut de mécanique céleste et de calcul des éphémérides (IMCCE): the Variations Séculaires des Orbites Planétaires (VSOP) theories, essentially issued from the research works of P. Bretagnon, and the Theory of Outer Planets (TOP) theories, derived from the works of J.-L. Simon. The VSOP theories are solutions for the motion of the eight planets in the solar system; they are very precise over a time span of about 1000 years and the ephemerides published by the IMCCE from 1984 to 2006 were derived from VSOP. The TOP theories are solutions for the motion of the four major planets and are accurate over several thousands of years. Currently, the ephemerides of the planets are obtained mainly from very precise numerical integrations fitted to the most recent observations: the European ephemerides Intégration Numérique Planétaire de l'Observatoire de Paris (INPOP, Fienga et al. 2008, 2009, 2011), the American ephemerides Development Ephemeris (DE; e.g. Standish 1998; or more

recently Folkner et al. 2009; Folkner 2010) and the Russian ephemerides Ephemerides of Planets and the Moon (EPM; Pitjeva 2005, 2010). Nevertheless, the construction of analytical planetary theories remains useful: (i) even if the aim of analytical theories is not to compete with numerical integrations for space engineering, they can provide precise ephemerides for the concern of most astronomers; (ii) their precision slowly decrease with time and they remain accurate over several thousand years; (iii) they allow a precise analysis of the perturbations; (iv) they are useful for some problems such as the study of the Earth's rotation; and (v) from these theories it is possible to obtain compact solutions of good precision.

We first built the analytical solutions VSOP2010 and TOP2010, fitted to DE405 (Standish 1998). We then built VSOP2013 and TOP2013, fitted to INPOP10a (Fienga et al. 2011). This paper particularly deals with VSOP2013 and TOP2013. After giving the common characteristics of these two kinds of solutions in Sect. 2, we will present the construction of the VSOP theories (Sect. 3) and of the TOP theories (Sect. 4). A method to build an analytical solution of the motion of Pluto is given in Sect. 5. Perturbations by asteroids are studied in Sect. 6. Lastly, the results are presented and discussed in Sect. 7.

2. Common characteristics

The two solutions VSOP and TOP have several characteristics in common that are given here.

[★] VSOP2013 and TOP2013 are available by ftp on: [ftp.imcce.fr/pub/ephem/planets/vsop2013](ftp://imcce.fr/pub/ephem/planets/vsop2013) and [ftp.imcce.fr/pub/ephem/planets/top2013](ftp://imcce.fr/pub/ephem/planets/top2013).

¹ The word "theory" is a traditional term used in celestial mechanics for the analytical solution of the equations of the motion of solar system bodies.

2.1. Variables and equations

- The variables used are the elliptic variables a , λ , $k = e \cos \varpi$, $h = e \sin \varpi$, $q = \sin i/2 \cos \Omega$, and $p = \sin i/2 \sin \Omega$ where a is the semi-major axis, λ the mean longitude, e the eccentricity of the orbit, ϖ the longitude of perihelion, i the inclination to the ecliptic J2000, and Ω the longitude of the ascending node.
- The equations of the motion of the planets are the Lagrange differential equations (for the computing of the second members of the Lagrange equations, see [Chapront et al. 1975](#)).

2.2. Form of the solutions

There are two kinds of analytical planetary theories: general planetary theories and classical planetary theories. In general planetary theories, the variables are developed under the form of Fourier series, with arguments being linear combinations of short-period arguments (mean longitudes of the planets) and long-period arguments (longitudes of nodes and perihelia). Developing the long-period arguments with respect to time, we obtain the classical planetary theories under the form of Poisson series, with arguments being linear combinations of short-period arguments. The VSOP and TOP solutions are classical planetary theories. For a planet j , we note that x_j are the elements other than mean longitudes λ_j ; x_j and λ_j have the form

$$\begin{aligned} x_j &= x_j^0 + x_j^1 t + \dots + x_j^p t^p + X_j^0 + t X_j^1 + \dots + t^p X_j^p \\ \lambda_j &= \lambda_j^0 + n_j t + l_j^2 t^2 + \dots + l_j^p t^p + L_j^0 + t L_j^1 + \dots + t^p L_j^p, \end{aligned} \quad (1)$$

where t is the time (the timescale will be explained in the following sections), x_j^0, \dots, x_j^p and $\lambda_j^0, n_j, \dots, l_j^p$ are numerical coefficients, and X_j^q and L_j^q are Fourier series of short-period arguments the form of which will be explained in the following sections.

2.3. Integration of the equations

- Results of the integration are obtained in the following way. First, the perturbations are built order by order up to the third order with respect to the masses. An iterative method is then used for the following computations.
- The integration constants x_j^0 and λ_j^0 and the mean motions n_j are determined by fitting, over the time span from 1890 to 2000, the solutions to the numerical integrations DE405 ([Standish 1998](#)) for the solutions VSOP2010 and TOP2010 or INPOP10a ([Fienga et al. 2011](#)) for VSOP2013 and TOP2013.

We note that the numerical formulae given in the following sections will generally relate to the solutions fitted to INPOP10a.

3. The VSOP theories

The VSOP theories are analytical theories for the eight planets: Mercury, Venus, the Earth-Moon barycenter (EMB), Mars, Jupiter, Saturn, Uranus, and Neptune.

3.1. Historical review

The first VSOP solutions were:

- VSOP82 ([Bretagnon 1982](#)). In this solution, the iterative method was developed up to the fifth order with respect to the masses. The relativity was introduced by the

Schwarzschild problem and the perturbations of the Earth-Moon barycenter by the Moon were taken into account. So, the arguments of the Poisson series were the mean mean longitudes of the planets and the Delaunay arguments of the Moon D, F, and L. The mean mean longitudes are linear functions of time defined for each planet j by

$$\bar{\lambda}_j = \lambda_j^0 + n_j t, \quad (2)$$

where n_j is the mean motion of the planet j . VSOP82 was fitted to the numerical integration of the JPL, DE200 ([Standish 1982](#)). The timescale is barycentric dynamical time (TDB) obtained from [Fairhead & Bretagnon \(1990\)](#).

- VSOP87 ([Bretagnon & Francou 1988](#)). This solution was an extension of VSOP82, so this solution was also fitted to DE200. The solutions were expressed with elliptic elements and also with rectangular (X, Y, Z) or spherical (longitude, latitude, and radius vector) variables. The reference frames were the ecliptic and equinox J2000 or the ecliptic and equinox of the date. The coordinates were heliocentric or barycentric.

VSOP82 and VSOP87 were used to compute the ephemerides of the IMCCE up to the construction of the INPOP numerical integrations ([Fienga et al. 2009, 2011](#)).

- VSOP2000 ([Moisson & Bretagnon 2001](#)). This solution was an improvement of VSOP82. The iterative method was developed up to the eighth order of the masses. The perturbations by the five big asteroids Ceres, Pallas, Vesta, Iris, and Bamberga were introduced during the iterations. Second order perturbations by the Moon on Mercury, Venus, the Earth-Moon barycenter, and Mars were computed. Relativistic corrections were also introduced in the iterations. The arguments of the series were linear combinations of 16 angles, the mean mean longitudes of the eight major planets and of the five big asteroids, and the three Delaunay angles of the Moon. The solution was fitted to the numerical integration of the JPL DE403 ([Standish et al. 1995](#)). This solution was 10 to 100 times more precise than VSOP82 and VSOP87.

3.2. New VSOP theories

3.2.1. The works of P. Bretagnon

Starting in VSOP2000, Bretagnon performed 15 additional iterations with a numerical precision 10 times better than VSOP2000, with the following characteristics:

- The Poisson series were developed up to the 12th degree with respect to time.
- The eight major planets and the five big asteroids had their orbits analytically computed during the same process.
- Perturbations of the Moon issued from ELP2000 ([Chapront-Touze & Chapront 1983](#)) on the eight major planets were introduced during the iteration process.
- Relativistic corrections were introduced in the iterations.
- The solution was fitted to DE403 ([Standish et al. 1995](#)) using the inertial mean ecliptic of J2000 defined by [Chapront et al. \(2002\)](#). Starting from the equatorial rectangular variables $(x, y, z)_{\text{equ}}$ given by DE403, we computed the ecliptic rectangular variables $(x, y, z)_{\text{ecl}}$ by

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix}_{\text{ecl}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \epsilon & \sin \epsilon \\ 0 & -\sin \epsilon & \cos \epsilon \end{pmatrix} \begin{pmatrix} \cos \phi & \sin \phi & 0 \\ -\sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}_{\text{equ}}, \quad (3)$$

where ϵ and ϕ are issued from [Chapront et al. \(2002\)](#) as

$$\epsilon = 23^\circ 26' 21''.40928 \quad \text{and} \quad \phi = -0.05294''. \quad (4)$$

From these ecliptic rectangular coordinates, we computed elliptic elements issued from DE403 and we computed the integration constants by fitting elliptic elements of VSOP to the elements issued from DE403 over the time span from 1890 to 2000. We note that the inertial mean ecliptic J2000 that was used was very close to but not exactly the same as the dynamic ecliptic of VSOP. So, the integration constants of the variables q and p of the EMB were very small but not equal to zero.

3.2.2. The construction of the VSOP2010 and VSOP2013 solutions

Unfortunately, Bretagnon did not have time to complete his work. We took his work up again introducing various changes and supplements

- The integration constants have been computed, first by fitting VSOP2010 to the numerical integration of the JPL DE405, then to different versions of the INPOP numerical integrations INPOP06 ([Fienga et al. 2008](#)), INPOP08 ([Fienga et al. 2009](#)), and finally INPOP10a ([Fienga et al. 2011](#)) for VSOP2013. These fits were done over the time span from 1890 to 2000.
- We have computed the perturbations due to the J_2 of the Sun on the planets Mercury, Venus, EMB, and Mars. The values used for the J_2 of the Sun are 2×10^{-7} for DE405 and 2.4×10^{-7} for INPOP10a.
- We have introduced the perturbations induced by the asteroids as explained in Sect. 6.
- We have added the perturbations of Pluto on the outer planets in the form of Poisson series as explained in Sect. 5.4.5.
- Starting from the TOP solutions, we have improved the precision of the theories of Jupiter and Saturn over a large time span as explained in Sect. 4.3.
- The arguments of the series are linear combinations of 17 angles: the 16 angles of VSOP2000 and the linear function of time μ defined below.
- The timescale is TDB and the relationship with terrestrial time (TT) depends on the ephemeris. For DE405, one can use TE405 ([Irwin & Fukushima 1999](#)). For INPOP10a, the transformation between TT and TDB is numerically integrated with the equations of the motion of the bodies, leading to a 4D-ephemeris (see [Fienga et al. 2009](#)).
- The solutions are fitted with DE405 or INPOP10a as explained in Sect. 3.2.1. For the fit to DE405, we use the values given by [Chapront et al. \(2002\)](#)

$$\epsilon = 23^\circ 26' 21''.40960 \quad \text{and} \quad \phi = -0.05028''. \quad (5)$$

The fit to INPOP10a is made starting from the equatorial rectangular coordinates of INPOP10a and using the angles given in Eq. (5) to compute the ecliptic rectangular coordinates, then the elliptic elements. The comparison, from 1890 to 2000, with the elliptic elements issued from DE405 gives small corrections Δq and Δp for p and q of EMB. From these corrections, we deduce the corrections $\Delta\epsilon$ and $\Delta\phi$ by

$$\Delta\epsilon = 2\Delta q; \Delta\phi = \frac{2\Delta p}{\sin \epsilon}. \quad (6)$$

We finally found the values corresponding to INPOP10a:

$$\epsilon = 23^\circ 26' 21''.41136 \quad \text{and} \quad \phi = -0.05188''. \quad (7)$$

Table 1. GM of the Sun, the planets, and the five big asteroids used for the construction of INPOP10a.

Celestial body	GM (m ³ s ⁻²)
Sun	1.327 124 4006 × 10 ²⁰
Mercury	2.203 208 0486 × 10 ¹³
Venus	3.248 585 9883 × 10 ¹⁴
EMB	4.035 032 4500 × 10 ¹⁴
Mars	4.282 831 4258 × 10 ¹³
Vesta	1.766 887 8029 × 10 ¹⁰
Iris	1.031 426 8890 × 10 ⁹
Bamberg	6.227 510 9355 × 10 ⁸
Ceres	6.314 933 5465 × 10 ¹⁰
Pallas	1.478 332 7322 × 10 ¹⁰
Jupiter	1.267 127 6483 × 10 ¹⁷
Saturn	3.794 062 6061 × 10 ¹⁶
Uranus	5.794 549 0071 × 10 ¹⁵
Neptune	6.836 534 0639 × 10 ¹⁵
Pluto	9.816 008 8771 × 10 ¹¹

We note that if, starting from the values in Eq. (5), we apply this method replacing INPOP10a by DE403, we find $\epsilon = 23^\circ 26' 21''.40926$ and $\phi = -0.05296''$, values very close to the Chapront values given in Eq. (4), that validates our method. The values found in Eq. (7) are consistent with the ICRF (International Celestial Reference Frame) definition. Indeed, by the fit of INPOP10a to planet positions obtained by VLBI (Very Long Baseline Interferometry) tracking of spacecraft, INPOP10a is directly tied to ICRF at the precision of these observations, about a few mas ([Fienga et al. 2011](#)). In consequence, VSOP2013 being fitted to INPOP10a can then be seen as linked to ICRF by means of INPOP10a.

We use the systems of planetary masses of DE405 for VSOP2010 and of INPOP10a for VSOP2013. We note G the newtonian gravitational constant and M the mass of a celestial body. Table 1 gives the GM of the Sun, the planets, and the five big asteroids used for the construction of INPOP10a.

4. The TOP theories

The TOP (Theory of the Outer Planets) theories are analytical theories for the motion of the four large planets Jupiter, Saturn, Uranus, and Neptune.

4.1. Historical review

The first TOP solutions were:

- TOP82 ([Simon 1983](#)). In this solution, the perturbations were Poisson series of the mean mean longitudes of the four large Planets. A supplementary input was brought to the theory for the couple Jupiter-Saturn by computing, with an iterative method and using harmonic analysis, the mutual perturbations Jupiter-Saturn up to the seventh order with respect to the masses. The relativity was introduced as in VSOP82. The solution also contained the perturbations of the four large planets by the telluric planets at the third order of masses from VSOP82. TOP82 was fitted to DE200.
- An extension of TOP82: JASON84 (Jupiter And Saturn Orbits from Neolithic). JASON84 ([Simon & Bretagnon 1984](#)) was an extension of TOP82 where the mutual perturbations Jupiter-Saturn were computed up to the 20th order

of masses. Before the construction of the new TOP solutions, JASON84 was the most precise theory of the motion of Jupiter and Saturn over the time span [J2000-6000, J2000+6000] and was used to build the tables for the motion of those planets in the book *Planetary Programs and Tables* from -4000 to +2800 (Bretagnon & Simon 1986).

4.2. New TOP theories

The aim of TOP is not to compete with VSOP over a short time span but to build very precise theories of the four large planets over several thousands of years. For that purpose we use a completely different representation.

4.2.1. The representation

The solutions are also Poisson series of the form of Eqs. (1), but X_j^q and L_j^q are Fourier series of only one argument μ

$$X_j^q/L_j^q = \sum_r A_r \cos r\mu + B_r \sin r\mu, \quad (8)$$

where r is a positive integer. The maximum value of r is $2^{19}(524\,288)$ and μ is linked to the mean motions of Jupiter and Saturn n_5 and n_6 by:

$$\mu = (n_5 - n_6)t/880 = 0.359\,536\,23\,t, \quad (9)$$

where t is measured in thousands of years from J2000 and with the values of n_5 and n_6 given in Table 6. The period of μ is about 17 485 years. The mean motions of the four major planets are linked to μ by relations of the form

$$\begin{aligned} n_5t &= 1473\mu + \sigma_5t = 1473\mu + 0.094\,095\,55t \\ n_6t &= 593\mu + \sigma_6t = 593\mu + 0.094\,095\,55t \\ n_7t &= 208\mu + \sigma_7t = 208\mu - 0.001\,873\,89t \\ n_8t &= 106\mu + \sigma_8t = 106\mu + 0.022\,132\,13t. \end{aligned} \quad (10)$$

This representation was chosen because perturbations are more convergent using this form than the classical form of Poisson series of the mean mean longitudes for Jupiter and Saturn because the choice of μ allows us to take in account an important part of the development with respect to the time of the long periods of a general theory of the couple Jupiter-Saturn (Simon et al. 1992). For Uranus and Neptune, the convergence is the same as in the classical form.

4.2.2. Construction of the solution

The solution is computed by the iterative method of Simon & Joutel (1988). The Lagrange equations are $\frac{dx}{dt} = F(\mu, t)$, with $F(\mu, t_i)$ computed by harmonical analysis, for 13 values of the time $t_0, t_0 + \Delta t, t_0 - \Delta t, \dots, t_0 + 6\Delta t, t_0 - 6\Delta t, t_0$ corresponding to J2000 and $\Delta t = 1200$ yrs. After interpolation and integration, the solution has the form of Eqs. (1) with $p = 12$. The integration constants have been computed by fitting TOP2010 to DE405 and TOP2013 to INPOP10a, over the time span from 1890 to 2000.

4.2.3. Perturbations by the telluric planets

For TOP2010, we used the perturbations by the telluric planets developed up to the third order of the masses extracted from

VSOP82. They are written as μ series and are introduced in the iterations. For TOP2013, we obtain these perturbations by computing the differences between the last iteration of the iterative process used for VSOP2013 and the last iteration computed without taking in account the telluric planets. This method gives a better estimation of the perturbations by the telluric planets as will be seen in Sect. 7.3.2.

4.2.4. Heliocentric spherical and rectangular variables

It is easy to compute by harmonic analysis the heliocentric spherical variables (longitude, latitude, radius vector) and the heliocentric rectangular variables (X, Y, Z) from the elliptic elements, under the form of Poisson series of μ . These variables have been introduced in the TOP solutions.

4.3. Amelioration of VSOP theories using TOP theories

The good convergence of the mutual perturbations Jupiter-Saturn in the TOP solution can be used to improve the perturbations corresponding to some periodic terms in the VSOP representation of the semi-major axis and the mean longitude of these two planets. As an example of how the VSOP improvement is obtained for the Jupiter mean longitude development we take the argument $4\lambda_5 - 10\lambda_6$.

4.3.1. VSOP and TOP developments

In VSOP, this development has the form

$$L_v = \sum_{p=0,11} t^p [s_{4-10}^p \sin(4\lambda_5 - 10\lambda_6) + c_{4-10}^p \cos(4\lambda_5 - 10\lambda_6)]. \quad (11)$$

This argument corresponds to the TOP argument 38μ . The TOP development has the form

$$L_t = \sum_{q=0,12} t^q [s_{38}^q \sin(38\mu) + c_{38}^q \cos(38\mu)]. \quad (12)$$

The values of s_{4-10}^p and c_{4-10}^p are given in Cols. 2 and 3 of Table 2, and the values of s_{38}^q and c_{38}^q in Cols. 4 and 5. In Table 2, to estimate the amplitude of the Poisson terms for $t = \pm 6000$ yrs, the coefficients are given in arcseconds for the periodic terms and in arcseconds per 6000 years for the Poisson terms.

4.3.2. TOP developed in VSOP form

From Eqs. (10), we find

$$38\mu = -(4\lambda_5 - 10\lambda_6) - 0.564\,573\,30\,t, \quad (13)$$

where t is measured in thousands of years from J2000. Substituting the value of 38μ given by Eq. (13) in Eq. (12), we obtain a Poisson development issued from TOP but under the VSOP representation. Up to t^{20} this development has the form

$$L_{tv} = \sum_{r=0,20} t^r [s_{4-10}^{r'} \sin(4\lambda_5 - 10\lambda_6) + c_{4-10}^{r'} \cos(4\lambda_5 - 10\lambda_6)]; \quad (14)$$

the values of $s_{4-10}^{r'}$ and $c_{4-10}^{r'}$ are given in Cols. 6 and 7 of Table 2.

Table 2. Poisson developments of the mean longitude of Jupiter for the argument $4\lambda_5 - 10\lambda_6$ for VSOP2013, TOP2013, TOP2013 in VSOP form, and corrections to VSOP2013.

	VSOP2013		TOP2013		TOP2013 in VSOP form		Corrections to VSOP2013	
	s_{4-10}^p	c_{4-10}^p	s_{38}^q	c_{38}^q	$s_{4-10}^{r'}$	$c_{4-10}^{r'}$	$\delta s_{4-10}^{r'}$	$\delta c_{4-10}^{r'}$
t^0	10.370	3.476	-10.556	2.860	10.370	3.476	0	0
t	-22.920	47.143	10.418	12.650	-22.919	47.144	0	0
t^2	-106.469	-70.446	7.007	-12.358	-106.471	-70.444	0	0
t^3	138.263	-156.792	-8.624	0.062	138.254	-156.791	0	0
t^4	166.715	197.399	1.854	3.923	166.699	197.380	0	0
t^5	-220.086	133.030	1.223	-1.711	-220.056	132.897	0	0
t^6	-78.488	-200.287	-0.726	-0.218	-78.421	-200.435	0	0
t^7	153.283	-29.450	-0.021	0.294	153.211	-29.432	0	0
t^8	-0.706	100.680	0.066	0.026	-0.551	100.738	0	0
t^9	-56.991	-12.332	0.025	-0.031	-57.910	-12.219	-0.919	0.113
t^{10}	11.877	-26.174	-0.012	-0.021	12.823	-29.575	0.945	-3.401
t^{11}	-0.018	-0.011	-0.006	0.005	13.666	9.282	13.685	9.292
t^{12}			-0.002	-0.003	-5.526	5.856	-5.526	5.856
t^{13}					-2.405	-2.866	-2.405	-2.866
t^{14}					1.372	-0.983	1.372	-0.983
t^{15}					0.411	0.607	0.411	0.607
t^{16}					-0.253	0.175	-0.253	0.175
t^{17}					-0.074	-0.100	-0.074	-0.100
t^{18}					0.038	-0.030	0.038	-0.030
t^{19}					0.011	0.013	0.011	0.013
t^{20}					-0.004	0.004	-0.004	0.004

Notes. Units are in arcseconds for the periodic terms and arcseconds per 6000 years for the Poisson terms.

4.3.3. Comparison between these developments

The comparison between the coefficients of the Cols. 2 and 3 of Table 2 with the coefficients of the Cols. 4 and 5 shows the best convergence of the TOP representation. For instance, for $t = \pm 6000$ yrs the amplitude of the Poisson terms in t^9 is about $58''$, $\sqrt{(s_{4-10}^p)^2 + (c_{4-10}^p)^2}$, for VSOP2013 and only $0.04''$, $\sqrt{(s_{38}^q)^2 + (c_{38}^q)^2}$, for TOP2013. The comparison between the coefficients of the Cols. 2 and 3 with the coefficients of the Cols. 6 and 7 shows the inaccuracy of the VSOP representation from Poisson terms in t^{11} .

4.3.4. Correction of the VSOP development

The VSOP and TOP theories have been built in different ways and their integration constants are not the same. So, the change of all VSOP coefficients of Eq. (11) by the coefficients of Eq. (13) does not give good results. After some tests, we have estimated that the best correction for this argument is to add to the VSOP coefficients the differences between the coefficients of Eq. (13) and the coefficients of Eq. (11) starting from Poisson terms in t^9 . So, these corrections have the form

$$\delta L_v = \sum_{r=9,20} t^r [\delta s_{4-10}^{r'} \sin(4\lambda_5 - 10\lambda_6) + \delta c_{4-10}^{r'} \cos(4\lambda_5 - 10\lambda_6)]. \quad (15)$$

The values of $\delta s_{4-10}^{r'}$ and $\delta c_{4-10}^{r'}$ are given in Cols. 8 and 9 of Table 2.

4.3.5. Application of the method

This method has been applied to the Poisson developments of the semi-major axis and the mean longitude of Jupiter and Saturn for

the 12 arguments

$$\begin{aligned} 4\lambda_5 - 11\lambda_6 + 3\lambda_7 \quad (7\mu) & & 2\lambda_5 - 6\lambda_6 + 3\lambda_7 \quad (12\mu) & & (16) \\ 2\lambda_5 - 5\lambda_6 \quad (19\mu) & & 6\lambda_5 - 16\lambda_6 + 3\lambda_7 \quad (26\mu) & & \\ 4\lambda_5 - 10\lambda_6 \quad (38\mu) & & 8\lambda_5 - 21\lambda_6 + 3\lambda_7 \quad (45\mu) & & \\ 6\lambda_5 - 15\lambda_6 \quad (57\mu) & & 8\lambda_5 - 20\lambda_6 \quad (76\mu) & & \\ 7\lambda_5 - 17\lambda_6 \quad (230\mu) & & 4\lambda_5 - 12\lambda_6 \quad (249\mu) & & \\ 6\lambda_5 - 14\lambda_6 \quad (536\mu) & & 4\lambda_5 - 9\lambda_6 \quad (555\mu) & & \end{aligned}$$

Though not rigorous, this method significantly improves the precision of VSOP for large time spans, as we shall see in Sect. 7.4.2.

4.3.6. Degree of the Poisson development in VSOP

With these complements, the Poisson series of VSOP2013 and VSOP2010 are developed up to the 20th degree with respect to time.

5. Theory of the motion of Pluto

5.1. Perturbations by Pluto on the major planets

To obtain a precise theory of the major planets, especially for Uranus and Neptune, it is necessary to include the perturbations due to Pluto (or, more exactly, of the Pluto-Charon barycenter) and, therefore, to build an analytical theory of Pluto at least at the first order of masses.

5.2. Construction of the theory of Pluto in the classical form

A very close Neptune-Pluto resonance is the argument $2\bar{\lambda}_8 - 3\bar{\lambda}_9$ with a period of about 20 000 years, $\bar{\lambda}_8$ and $\bar{\lambda}_9$ being the mean longitudes of Neptune and Pluto, respectively. Looking at

Table 3, we can understand why the construction of a theory of Pluto in the classical form is a complex problem. This table gives a few terms of the perturbations at the first order with respect to the masses for the mean longitude, computed with a precision of 5^{-11} rad (0.01 mas), for two couples of planets:

- Saturn perturbed by Jupiter (which represents the most important first order perturbations obtained in the construction of an analytical theory of the eight planets),
- Pluto perturbed by Neptune.

We can see that the amplitude of the most important perturbation of the mean longitude of Pluto, 4.56 rad (9.4×10^5 arcsec), which corresponds to the resonance $2\bar{\lambda}_8 - 3\bar{\lambda}_9$, is greater than the most important perturbation of the mean longitude of Saturn, 0.013 rad (2.7×10^3 arcsec), by a factor of 350. Moreover, the number of terms greater than 5^{-11} rad (0.01 mas) is 329 for Saturn perturbed by Jupiter and 15 539 for Pluto perturbed by Neptune. From the second iteration, the number of terms of the perturbations of the mean longitude of Pluto soars and it becomes impossible to build an analytical theory of Pluto in the classical form of series of the mean mean longitudes. In our representation using μ , we obtain

$$n_8 t = 106\mu + 0.022t \quad n_9 t = 71\mu - 0.176t, \quad (17)$$

where the coefficients of t are in rad/1000 yrs. The mean motion of Neptune n_8 is given in Table 6. For the mean motion of Pluto n_9 , we used a preliminary value (25.350 505 rad/1000 yrs) given by Chapront (2012, priv. comm.). The resonance $2\bar{\lambda}_8 - 3\bar{\lambda}_9$ corresponds to the argument μ and we have exactly the same difficulties as in the classical representation.

5.3. Choice of a new argument for the representation

We chose a representation with an argument ν such that the resonance should correspond to 0ν . So, the perturbations corresponding to the resonance should be expanded in polynoms of time. We then take

$$\nu = n_8 t / 105 = 0.363\ 171\ 17t, \quad (18)$$

where t is measured in thousands of years from J2000 and with n_8 given in Table 6. This argument is very near to μ given by Eq. (9) and we obtain $n_9 t = 70\nu - 0.086\ 321\ 37t$. The resonance corresponds then to 0ν .

5.4. Construction of the analytical theory of Pluto

We built our theory of Pluto as explained in Simon (2004). As for the VSOP and TOP solutions, we built two solutions fitted to DE405 and INPOP10a, respectively.

5.4.1. First order theory

We first computed the mutual perturbations Jupiter-Pluto, Saturn-Pluto, Uranus-Pluto, and Neptune-Pluto in Poisson series of ν . The mutual perturbation Neptune-Pluto are computed without difficulties. For instance, the mean longitude of Pluto contains, for this couple, 44 terms with an amplitude greater than 0.01 mas (the amplitude of the most important term being about 30'') and 250 Poisson terms giving contributions greater than 0.01 mas over 1000 yrs. As an example, in the first order Saturn-Jupiter, computed in Poisson series of μ , the mean longitude of Saturn contains 329 terms with an amplitude greater than 0.01 mas and 496 Poisson terms giving contributions greater than 0.01 mas over 1000 yrs.

Table 3. Perturbations at the first order of the masses for the mean longitudes greater than 5×10^{-11} rad: comparison between Saturn perturbed by Jupiter and Pluto perturbed by Uranus.

Rank	Saturn/Jupiter		Pluto/Neptune	
	Argument	Amplitude	Argument	Amplitude
1	$2\lambda_5 - 5\lambda_6$	0.013	$2\lambda_8 - 3\lambda_9$	4.56
2	$\lambda_5 - \lambda_6$	3×10^{-3}	$4\lambda_8 - 6\lambda_9$	1.08
3	$\lambda_5 - 2\lambda_6$	2×10^{-3}	$6\lambda_8 - 9\lambda_9$	0.40
10	$3\lambda_5 - 4\lambda_6$	1×10^{-4}	$20\lambda_8 - 30\lambda_9$	1×10^{-2}
100	$14\lambda_5 - 16\lambda_6$	8×10^{-8}	$13\lambda_8 - 18\lambda_9$	4×10^{-5}
329	$17\lambda_5 - 25\lambda_6$	5×10^{-11}	$82\lambda_8 - 23\lambda_9$	5×10^{-6}
1000			$24\lambda_8 - 42\lambda_9$	8×10^{-7}
15 539			$155\lambda_8 - 220\lambda_9$	5×10^{-11}

Notes. Units of the amplitudes are in rad.

Table 4. Secular terms of the mean longitude of Pluto in ''/1000 yrs.

t^2	t^3	t^4	t^5	t^6	t^7	t^8	t^9
3770	397	20	7	0.7	0.1	0.03	0.0005

5.4.2. Construction of the theory

We applied the method used for building the TOP solutions for the combined and simultaneous resolution of two systems of equations in using the argument ν . The first system corresponds to the five planets Jupiter, Saturn, Uranus, Neptune, and Pluto, the second to the four major planets only. For the first iteration, we started from the first order theory for Pluto and from the TOP Poisson series of μ given in Poisson series of ν for the major planets. We also introduced the perturbations due to the telluric planets at the second order of masses. The resolution of the first system of equations gives the theory of the motion of Pluto. The differences between the solutions of the two systems give the perturbations of Pluto on the major planets.

5.4.3. The resonance $2\bar{\lambda}_8 - 3\bar{\lambda}_9$ (0ν)

The secular terms corresponding to the resonance converge perfectly and decrease with time as can be seen in Table 4 which gives the values of the secular terms of the mean longitude of Pluto in ''/1000 yrs.

5.4.4. The argument $\bar{\lambda}_7 - 3\bar{\lambda}_9$

This argument has a period of 4000 yrs and corresponds to the argument 4ν like the great inequality Uranus-Neptune $\bar{\lambda}_7 - 2\bar{\lambda}_8$. This argument is known as a long short-period argument, i.e. an argument with a period similar to the periods of the long-period arguments of general theories. As shown by Joutel (1990), in a classical planetary theory, the perturbations corresponding to this type of argument do not converge well. Here, the amplitude is about 700'' for the mean longitude of Pluto with an error of about 10''. We shall discuss the consequences of this error in Sect. 7.3.2.

Table 5. Integration constants and mean motions of VSOP2013.

Planet	a^0 (au)	λ^0 (rad)	k^0	h^0	q^0	p^0	n (rad/1000 yrs)
Mercury	0.387 098 3099	4.402 608 6317	0.044 660 6294	0.200 723 3087	0.040 615 6406	0.045 635 4933	26 087.903 140 6855
Venus	0.723 329 8199	3.176 134 4616	-0.004 492 8210	0.005 066 8515	0.006 824 1139	0.028 822 8192	10 213.285 547 4344
EMB	1.000 001 0176	1.753 470 3694	-0.003 740 8181	0.016 284 4892	-0.000 000 0014	-0.000 000 0010	6 283.075 850 3532
Mars	1.523 679 3402	6.203 500 0141	0.085 365 5932	-0.037 899 7092	0.010 470 4280	0.012 284 4865	3 340.612 434 1455
Jupiter	5.202 603 2063	0.599 546 1070	0.046 985 8470	0.012 003 7197	-0.002 065 6227	0.011 183 8646	529.690 961 5623
Saturn	9.554 910 3860	0.874 018 5101	-0.002 959 9134	0.055 429 6361	-0.008 717 4559	0.019 891 4362	213.299 086 1085
Uranus	19.218 438 5555	5.481 225 3957	-0.045 953 0748	0.005 648 3402	0.001 859 2408	0.006 486 0185	74.781 659 0308
Neptune	30.110 415 9870	5.311 897 9332	0.005 998 8382	0.006 691 8100	-0.010 291 4751	0.011 516 7667	38.132 972 2261

5.4.5. Introduction of the theories of Pluto in TOP and VSOP

The Poisson series of ν do not have a convergence as good as the series of μ for Jupiter and Saturn. For that reason, we kept μ as the argument of the TOP solutions and transformed in Poisson series of μ the Poisson series of ν corresponding to the complete solution of Pluto. Moreover, we transformed in Poisson series of μ the perturbations of Pluto on the major planets and we added them to the VSOP and TOP solutions of the major planets. So μ becomes an argument of the Poisson series of VSOP. At last, we have introduced the theory of the motion of Pluto in both TOP and VSOP solutions.

6. Perturbations by the asteroids

6.1. Representation of the perturbations

We introduced the analytical perturbations of the asteroids considered by numerical planetary ephemerides on the eight major planets. These perturbations are computed at the first order of masses and have the form of Poisson series of μ as explained in [Fienga & Simon \(2005\)](#). For each variable and each planet-asteroid couple, the first order perturbations are computed by harmonical analysis ([Simon 1986](#)) in the form of Fourier series. The arguments of the series have the form $i\bar{\lambda}_p + j\bar{\lambda}_a$, where i and j are integers and $\bar{\lambda}_p$ and $\bar{\lambda}_a$ are the mean mean longitudes of the planet and the asteroid, respectively. Then, writing $in_p + jn_a = q\mu + \epsilon t$ where q is an integer and ϵ is as small as possible, we transform the Fourier series of mean mean longitudes in Poisson series of μ . By addition of the Poisson series given by all the asteroids, we obtain, finally, for each variable of each planet, the perturbations under the form of only one Poisson series of μ .

6.2. The asteroids in DE405 and INPOP10a

The integration of the motion of the asteroids is not done in the same way for DE405 and INPOP10a. In DE405, the orbits of Ceres, Pallas, and Vesta under their own gravitational forces, those of the Sun, the planets, and the Moon are integrated separately from the planet integration. The orbits of the 297 asteroids (Iris and Bamberga included) are also integrated separately under the gravitational forces of the Sun, the planets, the Moon, Ceres, Pallas, and Vesta, but only the action of these 297 asteroids upon Mars, the Earth, and the Moon, as well as their contributions to the solar system barycenter, are included in the planetary integration. In INPOP10a, 165 asteroids are integrated with the planets and their perturbations are taken into account for all the planets. The analytical first order perturbations by the asteroids is a first approach of the complete perturbations of the asteroids on the planets. So the model of the VSOP solutions is

closer from INPOP10a model than the DE405 one. We shall talk about this point again in Sect. 7.3.1.

We note that the motion of the five big asteroids are included in the iterative process of VSOP. So, the first order perturbations are computed for 295 asteroids in VSOP2010 and for 160 asteroids in VSOP2013. For TOP2010 and TOP2013, the perturbations are computed for the 300 asteroids of DE405 and for the 165 asteroids of INPOP10a, respectively.

7. Results

7.1. Integration constants and mean elements

Mean elements of the variables x_j and λ_j , referred to J2000, are the secular parts of the Eqs. (1)

$$\langle x_j \rangle = x_j^0 + x_j^1 t + \dots + x_j^p t^p \quad (19)$$

$$\langle \lambda_j \rangle = \lambda_j^0 + n_j t + l_j^2 t^2 + \dots + l_p t^p.$$

The mean elements are useful for

- the determination of the starting integration constants in building classical planetary theories,
- the determination of the integration constants of general planetary theories ([Laskar 1988](#)),
- the amelioration of general planetary theories by fitting the long-period terms of these theories to the mean elements of classical theories ([Bretagnon & Simon 1990](#)).

The purpose of the mean elements is not to compute ephemerides. Nevertheless, it is possible to obtain approximate ephemerides of the planets by adding the small Poisson series of μ given by [Simon et al. \(1994\)](#) to the mean elements. For the heliocentric longitudes, the precision of these approximate ephemerides is about a few arcseconds for the telluric planets and Neptune and about a few tens of arcseconds for the other planets over the time span from 1000 to 3000. Mean elements are too voluminous to be published in this paper, but they are available on the WEB server of the IMCCE. We give here only the integration constants x_0^j and the mean motions n_j of the solutions VSOP2013 (Table 5) and TOP2013 (Table 6).

7.2. Number of terms of the series

We give the number of terms of the solutions VSOP2013 and TOP2013, for different levels of truncation in the series, in Table 7. For the small levels of truncation, the TOP solutions are more compact than the VSOP solutions for the major planets. This is due both to the best convergence of the Poisson series in the TOP representation and because numerous VSOP terms of a period superior to 1000 yrs are represented in TOP by the same small multiples of μ (smaller than 17μ).

Table 6. Integration constants and mean motions of TOP2013.

Planet	a^0 (au)	λ^0 (rad)	k^0	h^0	q^0	p^0	n (rad/1000 yrs)
Jupiter	5.202 603 2025	0.599 544 6520	0.046 985 8464	0.012 003 7085	-0.002 065 6227	0.011 183 8645	529.690 962 2786
Saturn	9.554 910 4300	0.874 020 9500	-0.002 959 8987	0.055 429 6608	-0.008 717 4558	0.019 891 4362	213.299 081 1942
Uranus	19.218 438 2726	5.481 221 8694	-0.045 953 1057	0.005 648 4158	0.001 859 2404	0.006 486 0177	74.781 661 6318
Neptune	30.110 415 8724	5.311 899 0423	0.005 998 8612	0.006 691 7075	-0.010 291 4756	0.011 516 7670	38.132 972 3622
Pluto	39.544 617 1440	4.165 471 1248	-0.178 738 9594	-0.173 404 7186	-0.051 702 3078	0.139 779 9252	25.335 660 2044

Table 7. Number of terms of the solutions VSOP2013 and TOP2013 for several levels of truncation.

Planet	Complete solution	10^{-14}	10^{-12}	10^{-10}	10^{-8}
Mercury	272 360	138 226	21 399	3625	533
Venus	289 647	225 161	33 674	5135	704
EMB	294 426	281 337	51 498	7505	984
Mars	309 140	309 107	118 169	15 136	1880
Jupiter	324 596	323 738	104 894	15 889	2348
	37 918	37 418	31 233	13 763	2103
Saturn	350 513	350 167	190 036	31 382	4368
	62 124	56 365	49 801	26 832	4071
Uranus	330 581	330 100	211 837	36 997	5045
	81 272	71 563	61 560	35 515	5890
Neptune	322 572	318 401	113 009	18 793	2712
	41 026	40 475	34 441	16 647	2993
Pluto	114 088	114 088	92 500	51 886	11 421

Notes. Units used for the levels of truncation are in: au for the semi-major axis, rad for the mean longitude, and without dimension for k , h , q , p . For Jupiter, Saturn, Uranus and Neptune, the first line corresponds to VSOP2013, the second line corresponds to TOP2013.

7.3. Precision over the time span from 1890 to 2000

7.3.1. VSOP solutions

We have estimated the precision of VSOP2013 and VSOP2010 by computing their maximum differences to INPOP10a and DE405 from 1890 to 2000. Table 8 gives these differences for the elliptic elements of the eight planets. This table also gives the differences between VSOP2000 and DE403 given by [Moisson & Bretagnon \(2001\)](#). The accuracy of VSOP2013 is very good. The precision of the mean longitudes is about a few 0.01 mas for Mercury, Venus, EMB, Saturn, and Neptune, 0.2 mas for Jupiter, and 0.7 mas for Mars and Uranus. Related to VSOP2000, the improvement is about a factor of 2 for Uranus and up to 24 for EMB and Neptune. This is especially the result of the better inclusion of the perturbations by the asteroids and Pluto. Moreover, we note that the VSOP2013-INPOP10a differences are smaller than the VSOP2010-DE405 differences by a factor of 2.5 to 5 for the planets from Mercury to Jupiter. The reason is that the computation of the perturbations by the asteroids is closer between VSOP2013 and INPOP10a than in the case of VSOP2010 and DE405 as explained in Sect. 6.2. Another estimation of the precision of our solutions is given in Table 9 which gives the accuracy of VSOP2013, VSOP2010, and VSOP2000 compared to the first version VSOP82 ([Bretagnon 1982](#)) for the mean longitudes of planets from 1890 to 2000. We can see that the improvement is very important for the new VSOP solutions, especially for VSOP2013 (by a factor between 15 and 2121).

Table 8. Maximum differences from 1890 to 2000 between VSOP2013, VSOP2010, VSOP2000, and the numerical integrations for the elliptic variables.

Planet	VSOP	a	λ	k	h	q	p
Mercury	2013	0.003	0.03	1.0	1.2	0.1	0.1
	2010	0.006	0.07	3.1	1.0	0.5	0.9
	2000	0.006	0.27	8.0	4.0	0.5	0.8
Venus	2013	0.002	0.02	0.3	0.2	0.1	1.1
	2010	0.003	0.08	1.0	1.4	0.8	0.7
	2000	0.012	0.29	10.0	3.0	1.1	1.5
EMB	2013	0.003	0.01	0.8	0.5	0.1	1.9
	2010	0.003	0.06	0.8	0.6	0.1	2.0
	2000	0.021	0.35	3.0	17.0	3.4	4.2
Mars	2013	0.078	0.74	4.6	5.4	0.3	1.1
	2010	0.073	2.01	7.0	6.4	0.7	1.6
	2000	0.134	2.88	12.0	30.0	2.6	7.2
Jupiter	2013	0.099	0.19	2.9	3.3	0.5	0.4
	2010	0.683	0.72	13.3	13.6	0.6	0.4
	2000	0.910	0.47	15.0	16.0	5.0	4.3
Saturn	2013	0.173	0.09	4.8	4.3	0.9	0.9
	2010	0.816	0.19	5.5	5.8	1.0	0.9
	2000	6.774	1.75	19.0	35.0	7.8	8.1
Uranus	2013	15.120	0.76	40.0	39.1	3.0	2.3
	2010	24.467	0.88	60.6	68.5	2.8	2.2
	2000	23.600	1.49	66.0	57.0	38.2	11.0
Neptune	2013	3.432	0.08	9.1	3.8	1.5	1.0
	2010	7.747	0.27	10.6	7.8	1.5	1.1
	2000	47.032	1.86	61.0	68.0	4.1	11.3

Notes. Units are in: km (a), mas (λ), and 10^{-10} (k , h , q , p). The numerical integrations of reference are INPOP10a for VSOP2013, DE405 for VSOP2010, and DE403 for VSOP2000. The values corresponding to VSOP2000 are issued from [Moisson & Bretagnon \(2001\)](#).

The differences between VSOP2013 and INPOP2010a for the mean longitudes of the planets from 1890 to 2000 are illustrated by Fig. 1.

7.3.2. TOP solutions

The Table 10 gives the differences between TOP2013 and INPOP10a, and between TOP2010 and DE405 for the elliptic elements of the four major planets and Pluto from 1890 to 2000. The orbits of the telluric planets are not integrated with the major planets and the TOP theories cannot be as accurate as the VSOP theories over short time spans. Nevertheless, we can see that TOP2013 is more precise than TOP2010 thanks to the method used for computing the perturbations by the telluric planets in TOP2013 as explained in Sect. 4.2.3. The precision of TOP2013 is of the same order as the precision of VSOP2000. For Pluto, over the time span from 1890 to 2000, the accuracy




Fig. 1. Mean longitudes of the planets: VSOP2013 – INPOP10a differences from 1890 to 2000. Units are in mas.

Table 9. Mean longitudes of the planets: gain in precision with respect to VSOP82 for VSOP2013, VSOP2010, and VSOP2000 from 1890 to 2000.

Planet	VSOP2013	VSOP2010	VSOP2000
Mercury	21	9	2
Venus	228	50	15
EMB	233	54	10
Mars	15	6	4
Jupiter	212	53	81
Saturn	962	496	53
Uranus	59	51	30
Neptune	2121	633	90

Table 10. Maximum differences from 1890 to 2000 between TOP2013, TOP2010, and the numerical integrations for the elliptic variables.

Planet	TOP	a	λ	k	h	q	p
Jupiter	2013	0.47	0.84	20.7	28.0	16.7	10.8
	2010	4.94	1.39	78.3	88.4	36.6	31.3
Saturn	2013	1.02	1.86	31.2	41.0	18.6	15.1
	2010	17.10	2.89	99.9	69.1	38.7	33.2
Uranus	2013	16.55	1.70	50.9	48.5	18.5	15.3
	2010	29.79	2.58	122.8	93.2	39.3	43.0
Neptune	2013	6.96	0.67	21.5	17.7	12.4	14.7
	2010	43.66	1.52	70.6	67.5	42.0	43.0
Pluto	2013	124.71	2.85	198.1	185.8	56.9	23.3
	2010	130.49	3.42	161.6	149.7	26.5	23.9




Fig. 2. Mean longitude of Pluto: TOP2013 – INPOP10a differences from 1890 to 2000. Units are in mas.

is about 3 mas for the mean longitude, about 125 km for a , and 2×10^{-8} for k and h . This accuracy is far below the actual uncertainties of the Pluto observations. The differences between TOP2013 and INPOP2010A for the mean longitude of Pluto from 1890 to 2000 are illustrated in Fig. 2. We see that these differences are short-period terms. Although, the short-period terms perfectly converge in the iterative process, the bad convergence of the perturbations corresponding to the argument $\lambda_7 - 3\lambda_9$ (see Sect. 5.3.4) involves an uncertainty in the determination of the integration constants explaining these short-period differences.

7.4. Precision over large time spans

The accuracy of our solutions over large time spans has been estimated by computing the differences with two numerical

Notes. Units are in: km (a), mas (λ), 10^{-10} (k, h, q, p). The numerical integrations of reference are INPOP10a for TOP2013 and DE405 for TOP2010.

integrations from -4000 to 8000 . The solutions VSOP2010 and TOP2010 have been compared with an internal numerical integration, the initial values of which are issued from the solutions. The solutions VSOP2013 and TOP2013 have been compared with the extension of INPOP10a from -4000 to 8000 (Manche 2012, priv. comm.). The comparisons with these two different integrations are similar. Tables 11 and 12 refer only to the comparisons with the extension of INPOP10a.

7.4.1. Telluric planets

Table 11 gives the differences between VSOP2013 and the extension of INPOP10a from -4000 to 8000 (Manche 2012, priv. comm.) for the mean longitude (λ) and the heliocentric coordinates (L, B, R) of the four telluric planets over three time spans: from 900 to 3100, from 0 to 4000, and from -4000 to 8000.

We can see that the precision remains good for large time spans. For instance, from 0 to 4000, the accuracy of the heliocentric longitude is better than $0.02''$ for Venus and $0.05''$ for Mercury and EMB. From -4000 to 8000 , the precision is better

Table 11. Telluric planets: maximum differences over large time spans between VSOP2013 and the extension of INPOP10a from -4000 to 8000 for the mean longitude and the heliocentric coordinates.

Planet	Time span	λ	L	B	R
Mercury	[900, 3100]	0.01	0.02	0.002	0.9
	[0, 4000]	0.03	0.05	0.006	2.6
	$[-4000, 8000]$	0.12	0.20	0.021	9.2
Venus	[900, 3100]	0.002	0.002	0.001	0.1
	[0, 4000]	0.02	0.02	0.002	0.2
	$[-4000, 8000]$	0.15	0.15	0.009	4.1
EMB	[900, 3100]	0.03	0.03	0.002	0.4
	[0, 4000]	0.05	0.06	0.003	1.9
	$[-4000, 8000]$	0.98	1.01	0.015	19.4
Mars	[900, 3100]	0.70	0.83	0.023	69.5
	[0, 4000]	1.06	1.28	0.036	107.2
	$[-4000, 8000]$	1.49	1.74	0.058	153.4

Notes. Units are in: arsec (λ , L , B) and km (R). The extension of INPOP10a is from Manche (2012, priv. comm.).

than $0.2''$ for Mercury and Venus, about $1''$ for EMB and $1.7''$ for Mars. The improvement is about a factor of 5 compared to VSOP2000. However, we see that the precision decreases with time more quickly for EMB and Mars than for Mercury and Venus. This is due to the perturbations by the asteroids which should be computed at the second order of masses to improve the accuracy of the solutions of the motion of these two planets over large time spans.

The differences between VSOP2013 and INPOP10a for the heliocentric longitudes of the telluric planets from -4000 to 8000 are illustrated in Fig. 3.

7.4.2. Major planets and Pluto

Table 12 gives, for the major planets and Pluto, the differences between VSOP2013, TOP2013, and the extension of INPOP10a for the same variables and the same time spans as in Table 11. For Jupiter and Saturn and for the time span from -4000 to 8000 , the second line of VSOP corresponds to the theory without the corrections from TOP2013. We see that the improvement due to the supplementary material from TOP2013 is about a factor of 5 for the mean and heliocentric longitudes of these two planets. From -4000 to 8000 , the precision of the heliocentric longitudes of the major planets is included between $1''$ and $5''$ for Jupiter, Uranus, and Neptune and about $11''$ for Saturn. This represents an improvement compared to VSOP2000 of a factor of 5 to 10.

Nevertheless, TOP2013 remains the more precise solution over large time spans. From -4000 to 8000 , the precision of the heliocentric longitudes of the major planets is about $0.4''$ for Jupiter, $0.7''$ for Uranus and Neptune, and $0.9''$ for Saturn. The gain in precision, compared to VSOP2013, is between 10 and 15 for Jupiter and Saturn and between 1.5 and 4 for Uranus and Neptune. The differences between VSOP2013, TOP2013, and the extension of INPOP10a for the heliocentric longitudes of the major planets from -4000 to 8000 are illustrated in Fig. 4.

For Pluto, the precision remains good from 900 to 3100 (about $0.8''$ for the heliocentric longitude) and still correct from 0 to 4000 (about $12''$). For larger time spans, the precision quickly decreases, but the existence of a libration of the

Table 12. Major planets and Pluto: maximum differences over large time spans between VSOP2013, TOP2013, and the extension of INPOP10a from -4000 to 8000 .

Planet	Time span	Theory	λ	L	B	R
Jupiter	[900, 3100]	VSOP	0.04	0.04	0.001	12
		TOP	0.01	0.01	0.001	21
	[0, 4000]	VSOP	0.22	0.24	0.004	64
		TOP	0.05	0.07	0.004	61
	$[-4000, 8000]$	VSOP	4.80	4.47	0.164	3 393
			19.90	22.21	0.505	5 858
TOP		0.40	0.45	0.051	343	
Saturn	[900, 3100]	VSOP	0.28	0.30	0.013	122
		TOP	0.03	0.04	0.002	50
	[0, 4000]	VSOP	2.07	2.24	0.098	758
		TOP	0.15	0.18	0.008	151
	$[-4000, 8000]$	VSOP	8.92	11.73	0.524	21 911
			35.58	41.97	1.543	35 637
TOP		0.74	0.89	0.081	1 299	
Uranus	[900, 3100]	VSOP	0.14	0.14	0.001	136
		TOP	0.01	0.02	0.001	102
	[0, 4000]	VSOP	0.30	0.34	0.004	433
		TOP	0.04	0.06	0.003	189
	$[-4000, 8000]$	VSOP	1.23	2.45	0.069	13 007
		TOP	0.42	0.70	0.037	4 113
Neptune	[900, 3100]	VSOP	0.06	0.07	0.002	143
		TOP	0.01	0.02	0.001	143
	[0, 4000]	VSOP	0.13	0.14	0.003	342
		TOP	0.04	0.06	0.003	311
	$[-4000, 8000]$	VSOP	1.28	1.27	0.044	7 761
		TOP	0.39	0.78	0.059	5 284
Pluto	[900, 3100]	TOP	0.36	0.80	0.222	6 593
	[0, 4000]	TOP	5.14	11.95	2.980	45 042

Notes. Variables and units are the same as in Table 11. For Jupiter and Saturn and for $[-4000, 8000]$, the second line of VSOP corresponds to the theory without the corrections from TOP2013.

longitude of Pluto, with a period about 19 900 yrs (Milani et al. 1989), means that we cannot hope to build an analytical solution of Pluto valid over time spans greater than a few thousand years.

7.5. The solutions on the WEB server of the IMCCE

7.5.1. The VSOP2013 solution²

For VSOP2013, we give on the WEB server of the IMCCE:

- The complete Poisson developments of the elliptic variables of the eight planets and Pluto.
- Developments in the form of Tchebychev polynomials for the heliocentric rectangular coordinates (positions and velocities) of the eight planets over different time spans.
- The mean elements of the elliptic variables of the eight planets.
- Subroutines for reading the developments and computing the coordinates for given values of time.

² <ftp://ftp.imcce.fr/pub/ephem/planets/vsop2013>




Fig. 3. Heliocentric longitudes of the telluric planets: differences between VSOP2013 and the extension of INPOP10a from -4000 to 8000 . Units are in arcsecond .

Fig. 4. Heliocentric longitudes of the major planets: differences between TOP2013 and the extension of INPOP10a (straight lines) and between VSOP2013 and the extension of INPOP10a (dotted lines) from -4000 to 8000 . Units are in arcseconds.

7.5.2. The TOP2013 solution³

The TOP2013 solution is much more compact than VSOP2013 and it is not necessary to use developments under the form of Tchebychev polynomials. So, we give:

- The complete Poisson developments of the elliptic variables (four major planets and Pluto) and of the heliocentric spherical and rectangular variables (four major planets).
- The mean elements of the elliptic variables of the four major planets and Pluto.
- Subroutines for reading the developments and computing the coordinates for given values of time.

7.6. Possible improvements

Some improvements of the theories can still be made. For VSOP2013, it would be possible

- to improve the accuracy of the motion of the telluric planets (specially EMB and Mars) over large time spans by computing the perturbations of the asteroids on the planets to the second order of masses;
- to improve the accuracy of the motion of Jupiter and Saturn over large time spans by developing the Poisson series up to the 20th degree with respect to time;
- to improve the accuracy of the relativistic corrections, using the techniques suggested by [Brumberg \(2012\)](#).

The TOP2013 theory could be more accurate by improving the modelling of the perturbations due to the telluric planets.

³ <ftp://ftp.imcce.fr/pub/ephem/planets/top2013>

Integrating the motion of the telluric planets in the iterative process would not be easy because the computation of very short periods in the motion of Mercury would lead to important difficulties in the harmonic analysis. A better method would be to do two kinds of iterations in the VSOP process, one with the telluric planets, another without. The difference between the two processes should give a good estimation of the perturbations of the telluric planets on the outer planets.

8. Conclusion

The new analytical theories VSOP2013 and TOP2013 built at the IMCCE are complementary and have been greatly enhanced since the previous versions. They are fitted to INPOP10a, a numerical integration of the motion of the planets that takes into account more recent observations. The VSOP2013 theory is the best analytical theory for the motion of the eight solar system planets over time spans of a few hundred years. The improvement over VSOP2000 is about a factor of 2 for Uranus, 5 for Mars and Jupiter, and greater than 10 for the other planets. Moreover, VSOP2013 remains very accurate over large time spans for the motion of the telluric planets. The TOP2013 theory was developed in a compact and easy to use form. It is the most precise theory for the motion of the four major planets over large time spans; TOP2013 also gives a solution of the motion of the Pluto-Charon barycenter with good accuracy up to time spans of a few thousand years.

Lastly, the subroutines used to build VSOP2013 and TOP2013 are ready to fit these theories to new versions of INPOP.

Acknowledgements. We dedicate this paper to the memory of Pierre Bretagnon who was the initiator of the VSOP theories and who devoted all his energy and his talent to the development of planetary analytical solutions for over thirty years. We thank the referee, Elena Pitjeva, for the helpful improvement she proposed to the original manuscript.

References

- Bretagnon, P. 1982, *A&A*, 114, 278
 Bretagnon, P., & Francou, G. 1988, *A&A*, 202, 309
 Bretagnon, P., & Simon, J.-L. 1986, Planetary programs and tables from –4000 to +2800 (Richmond: Willmann-Bell)
 Bretagnon, P., & Simon, J.-L. 1990, *A&A*, 239, 387
 Brumberg, V. A. 2012, Notes scientifiques et techniques de l’Institut de Mécanique céleste S096, 1
 Chapront, J., Bretagnon, P., & Mehl, M. 1975, *Celest. Mech.*, 11, 379
 Chapront, J., Chapront-Touzé, M., & Francou, G. 2002, *A&A*, 387, 700
 Chapront-Touze, M., & Chapront, J. 1983, *A&A*, 124, 50
 Fairhead, L., & Bretagnon, P. 1990, *A&A*, 229, 240
 Fienga, A., & Simon, J.-L. 2005, *A&A*, 429, 361
 Fienga, A., Manche, H., Laskar, J., & Gastineau, M. 2008, *A&A*, 477, 315
 Fienga, A., Laskar, J., Morley, T., et al. 2009, *A&A*, 507, 1675
 Fienga, A., Laskar, J., Kuchynka, P., et al. 2011, *Cel. Mech. Dyn. Astron.*, 111, 363
 Folkner, W. M. 2010, JPL Interoffice Memorandum, 343, R
 Folkner, W. M., Williams, J. G., & Boggs, D. H. 2009, Interplanetary Network Progress Report, 178, C1
 Irwin, A. W., & Fukushima, T. 1999, *A&A*, 348, 642
 Joutel, F. 1990, Thèse de doctorat, Observatoire de Paris
 Laskar, J. 1988, *A&A*, 198, 341
 Milani, A., Nobili, A. M., & Carpino, M. 1989, *Icarus*, 82, 200
 Moisson, X., & Bretagnon, P. 2001, *Celest. Mech. Dyn. Astron.*, 80, 205
 Pitjeva, E. V. 2005, *Solar System Research*, 39, 176
 Pitjeva, E. V. 2010, in IAU Symp. 261, eds. S. A. Klioner, P. K. Seidelmann, & M. H. Soffel, 170
 Simon, J. L. 1983, *A&A*, 120, 197
 Simon, J. L. 1986, Notes scientifiques et techniques de l’Institut de Mécanique céleste S005, 1
 Simon, J. L. 2004, Notes scientifiques et techniques de l’Institut de Mécanique céleste S081, 53
 Simon, J. L., & Bretagnon, P. 1984, *A&A*, 138, 169
 Simon, J.-L., & Joutel, F. 1988, *A&A*, 205, 328
 Simon, J. L., Joutel, F., & Bretagnon, P. 1992, *A&A*, 265, 308
 Simon, J. L., Bretagnon, P., Chapront, J., et al. 1994, *A&A*, 282, 663
 Standish, Jr., E. M. 1982, *A&A*, 114, 297
 Standish, E. 1998, JPL Interoffice Memorandum, 312, F.98
 Standish, E., Newhall, X., Williams, J., & Folkner, W. 1995, JPL Interoffice Memorandum, 314, 10