

## The 2021 report of the Lancet Countdown on health and climate change: code red for a healthy future

Article (Accepted Version)

Romanello, Marina, McGushin, Alice, Di Napoli, Claudia, Drummond, Paul, Hughes, Nick, Jamart, Louis, Kennard, Harry, Lampard, Pete, Solano Rodriguez, Baltazar, Arnell, Nigel, Ayeb-Karlsson, Sonja, Belesova, Kristine, Cai, Wenjia, Campbell-Lendrum, Diarmid, Kniveton, Dominic et al. (2021) The 2021 report of the Lancet Countdown on health and climate change: code red for a healthy future. *Lancet*, 398 (10311). pp. 1619-1662. ISSN 0140-6736

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/102413/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

### **Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

# The 2021 Report of The *Lancet* Countdown on Health and Climate Change

Marina Romanello, Alice McGushin, Claudia Di Napoli, Paul Drummond, Nick Hughes, Louis Jamart, Harry Kennard, Pete Lampard, Baltazar Solano Rodriguez, Nigel Arnell, Sonja Ayeb-Karlsson, Kristine Belesova, Wenjia Cai, Diarmid Campbell-Lendrum, Stuart Capstick, Jonathan Chambers, Lingzhi Chu, Luisa Ciampi, Carole Dalin, Niheer Dasandi, Shouro Dasgupta, Michael Davies, Paula Dominguez-Salas, Robert Dubrow, Kristie L. Ebi, Matthew Eckelman, Paul Ekins, Luis E. Escobar, Lucien Georgeson, Delia Grace, Hilary Graham, Samuel H. Gunther, Stella Hartinger, Kehan He, Clare Heaviside, Jeremy Hess, Shih-Che Hsu, Slava Jankin, Marcia P. Jimenez, Ilan Kelman, Gregor Kiesewetter, Patrick Kinney, Tord Kjellstrom, Dominic Kniveton, Jason K.W. Lee, Bruno Lemke, Yang Liu, Zhao Liu, Melissa Lott, Rachel Lowe, Jaime Martinez-Urtaza, Mark Maslin, Lucy McAllister, Celia McMichael, Zhifu Mi, James Milner, Kelton Minor, Nahid Mohajeri, Maziar Moradi-Lakeh, Karyn Morrissey, Simon Munzert, Kris A. Murray, Tara Neville, Maria Nilsson, Nick Obradovich, Maquins Odhiambo Sewe, Tadj Oreszczyn, Matthias Otto, Fereidoon Owfi, Olivia Pearman, David Pencheon, Mahnaz Rabbaniha, Elizabeth Robinson, Joacim Rocklöv, Renee N. Salas, Jan C. Semenza, Jodi Sherman, Lihua Shi, Marco Springmann, Meisam Tabatabaei, Jonathon Taylor, Joaquin Trinanes, Joy Shumake-Guillemot, Bryan Vu, Fabian Wagner, Paul Wilkinson, Matthew Winning, Marisol Yglesias, Shihui Zhang  
Peng Gong<sup>a</sup>, Hugh Montgomery<sup>a</sup>, Anthony Costello<sup>a</sup>, Ian Hamilton\*  
[Insert institutional logos for inside cover]

<sup>a</sup> Co-Chair

\* Executive Director and Corresponding Author

[Word Count: 17,822]

## Table of Contents

List of Figures, Tables and Panels .....	3
List of Figures.....	3
List of Panels.....	5
List of Abbreviations .....	5
Executive Summary.....	8
Deepening inequities in a warming world.....	8
An inequitable response leaves the most vulnerable behind and fails all of us <b>Error! Bookmark not defined.</b>	
An unprecedented opportunity to ensure a healthy future for all .....	11
Introduction .....	13
Five years of tracking progress on health and climate change .....	14
Section 1: Climate Change Impacts, Exposures, and Vulnerability.....	18
1.1 Health and Heat.....	19
Indicator 1.1.1: Vulnerability to the Extremes of Heat .....	19
Indicator 1.1.2: Exposure of Vulnerable Populations to Heatwaves.....	19
Indicator 1.1.3: Heat and Physical Activity.....	21
Indicator 1.1.4: Change in Labour Capacity .....	22
Indicator 1.1.5: Heat and Sentiment .....	23
Indicator 1.1.6: Heat-Related Mortality .....	26
1.2 Health and Extreme Weather Events .....	27
Indicator 1.2.1: Wildfires.....	27
Indicator 1.2.2: Drought .....	28
Indicator 1.2.3: Lethality of Extreme Weather Events.....	29
1.3 Climate-Sensitive Infectious Diseases .....	29
Indicator 1.3.1: Climate Suitability for Infectious Disease Transmission .....	29
Indicator 1.3.2: Vulnerability to Mosquito-Borne Diseases .....	31
1.4 Food Security and Undernutrition.....	32
Indicator 1.4.1: Terrestrial Food Security and Undernutrition .....	32
Indicator 1.4.2: Marine Food Security.....	33
1.5 Migration, Displacement, and Rising Sea Levels .....	34
Conclusion .....	36
Section 2: Adaptation, Planning, and Resilience for Health .....	37
2.1: Adaptation Planning and Assessment .....	37
Indicator 2.1.1: National Adaptation Plans for Health.....	37
Indicator 2.1.2: National Assessments of Climate Change Impacts, Vulnerability, and Adaptation for Health .....	38
Indicator 2.1.3: City-level Climate Change Risk Assessments .....	39
Indicator 2.2: Climate Information Services for Health.....	40
2.3: Adaptation Delivery and Implementation.....	40
Indicator 2.3.1: Detection, Preparedness and Response to Health Emergencies .....	40
Indicator 2.3.2: Air Conditioning: Benefits and Harms .....	41
Indicator 2.3.3: Urban Green Space .....	44
Indicator 2.4: Health Adaptation-Related Global Funding and Financial Transactions.....	46
Conclusion .....	47
Section 3: Mitigation Actions and Health Co-Benefits.....	49
Indicator 3.1: Energy System and Health .....	49
Indicator 3.2: Clean Household Energy .....	51

Indicator 3.3: Mortality from Ambient Air Pollution by Sector .....	53
Indicator 3.4: Sustainable and Healthy Road Transport.....	55
3.5: Food, Agriculture, and Health .....	55
Indicator 3.5.1: Emissions from Agricultural Production and Consumption.....	55
Indicator 3.5.2: Diet and Health Co-Benefits .....	57
Indicator 3.6: Healthcare Sector Emissions .....	58
Conclusion .....	60
Section 4: Economics and Finance .....	62
4.1 The Economic Impact of Climate Change and its Mitigation.....	62
Indicator 4.1.1: Economic Losses due to Climate-Related Extreme Events .....	62
Indicator 4.1.2: Costs of Heat-Related Mortality .....	63
Indicator 4.1.3: Loss of Earnings from Heat-Related Labour Capacity Reduction .....	64
Indicator 4.1.4: Costs of the Health Impacts of Air Pollution .....	65
4.2 The Economics of the Transition to Zero-Carbon Economies .....	67
Indicator 4.2.1: Coal and Clean Energy Investment .....	68
Indicator 4.2.2: Employment in Low-Carbon and High-Carbon Industries .....	69
Indicator 4.2.3: Funds Divested from Fossil Fuels.....	70
Indicator 4.2.4: Net Value of Fossil Fuel Subsidies and Carbon Prices.....	71
Indicator 4.2.5: Production- and Consumption-Based Attribution of CO <sub>2</sub> and PM <sub>2.5</sub> Emissions ..	72
Conclusion .....	74
Section 5: Public and Political Engagement .....	76
Indicator 5.1 Media Coverage of Health and Climate Change .....	77
Indicator 5.2: Individual Engagement in Health and Climate Change .....	78
Indicator 5.3: Coverage of Health and Climate Change in Scientific Journals.....	79
Indicator 5.4: Government Engagement in Health and Climate Change .....	81
Indicator 5.5: Corporate Sector Engagement in Health and Climate change .....	84
Conclusion .....	85
Conclusion: the 2021 Report of the <i>Lancet</i> Countdown.....	87
References .....	91

## List of Figures, Tables and Panels

### List of Figures

Figure 1. Change in person-days of heatwave exposure relative to the 1986-2005 baseline. A) in the population aged over 65 years; B) in the population aged under 1 year of age. In each case, the countries with the highest exposure averages over the past 5 years are highlighted. ....	20
Figure 2. Average potential activity hours lost per person per day by 2019 Human Development Index country group, 1980-2020. ....	21
Figure 3. Heat-related potential hours of labour lost by sector and 2019 country Human Development Index group, 1990-2000 .....	23
Figure 4. Heatwaves and sentiment. Top: Annual effect of heatwave exposure on the sentiment of online expressions from 2015-2020. Coloured intervals depict 95% CIs of the estimated average change in positive (green) and negative (orange) sentiment expressions during days with heatwaves, relative to the median daily maximum temperature baseline range for each location and year. Sentiment was extracted from Twitter posts using a dictionary-based approach across multiple	

languages. Grey bars depict the geolocated Tweet count by year of observation. Bottom: Country-level count of geolocated tweets for 2015-2020.....	25
Figure 5. Heat-related mortality among the 65-and-older population in 2019, by country .....	26
Figure 6. Annual population-weighted mean changes in days of very high and extremely high fire danger from 2001-2004 to 2017-2020 for each country/territory. Large urban areas with population density $\geq 400$ persons/km <sup>2</sup> are excluded in the calculations of population-weighted mean values. ...	27
Figure 7. Percentage of land area affected by drought events per month .....	28
Figure 8. Change in climate suitability for infectious diseases. Solid lines represent the annual change. Straight lines represent the trend since 1950 (for dengue and malaria), 1982 (for Vibrio bacteria), and 2003 (for Vibrio cholerae) .....	31
Figure 9. Change in crop growth duration relative to the 1981–2010 baseline. The grey line represents the annual global area-weighted change. The blue line represents the running mean over 11 years (5 years forward and 5 years backward). .....	33
Figure 10. Global heat-related deaths averted by household air conditioning in the 65-and-older population (red line), proportion of households with air conditioning (blue line), and carbon dioxide emissions from air conditioning (green line), 2000-2019 .....	42
Figure 11. Average urban population-weighted peak Normalized Difference Vegetation Index (NDVI) in urban centres of >500,000 inhabitants by country, for 2010 (A) and 2020 (B). For countries without an urban centre of >500,000 inhabitants, the most populated urban centre was used in the analysis.....	45
Figure 12. Per capita potential adaptation transactions in the health and health care sector (A) and health-relevant sectors (B) for financial years 2015/16 to 2019/20, by 2019 Human Development Index country group.....	47
Figure 13. The carbon intensity of the energy system for 1970-2018, by 2019 Human Development Index country group.....	51
Figure 14. Residential energy supply by 2019 Human Development Index country group for 2000 to 2019. Primary axis: per capita fuel type (coloured bars). Secondary axis: percentage of population with primary reliance on clean fuels and technology for cooking. Data taken from the WHO and IEA. 198-200 .....	53
Figure 15. Mortality attributable to ambient PM <sub>2.5</sub> in 2015, 2018 and 2019 by source and by 2019 Human Development Index country group. ....	54
Figure 16. Per capita yearly greenhouse gas emissions associated with consumption of agri-food products, by 2019 Human Development Index country group and commodity, 2000-2018.....	57
Figure 17. Deaths attributable to imbalanced diets and weight in 2018 by risk factor, sex, and 2019 Human Development Index country group. The size of each component in the stacked bar represents its individual contribution to attributable deaths. Since these contributions cannot be summed directly, the overall contribution by diet and weight components are represented by the dots as given in the key.....	58
Figure 18. National per capita healthcare greenhouse gas emissions for 2018 against 2019 country Human Development Index level. Dot size is proportional to population .....	60
Figure 19. Monetised cost of heat-related (in terms of expressed as the number of people whose average income the loss is equivalent to) by 2019 Human Development Index country group for 2000-2019 .....	64
Figure 20 Average potential loss of earnings in the low Human Development Index country group as a result of potential labour loss due to heat exposure. Losses are presented as share of GDP, by sector of employment,. The agriculture and construction (sun additional) blocks represent the losses that would have been incurred in addition to those from agriculture and construction (shade) if all of the activities in these sectors had been carried out in direct sunlight.....	65

Figure 21. Economic cost of YLLs in 2015 and 2019, relative to the annual income of the average person and total GDP, by 2019 Human Development Index country group .....	66
Figure 22. Economic value of annual investment in renewable and fossil fuel energy supply and energy efficiency, 2014-2020.....	69
Figure 23. Net carbon prices (left), net carbon revenues (centre), and net carbon revenue as a share of current national health expenditure (right), across 84 countries in 2018, arranged by 2019 Human Development Index country group: low (n=1), medium (n=7), high (n=23) and very high (n=53). Boxes show the interquartile range (IQR), horizontal lines inside the boxes show the medians, and the brackets represent the full range from minimum to maximum. ....	72
Figure 24. The flows of embodied CO <sub>2</sub> and PM <sub>2.5</sub> emissions among different Human Development Index country groups in 2019 .....	74
Figure 25. Aggregate monthly co-clicks on Wikipedia articles related to human health and climate change, 2018—2020. Blue: co-click from health-related page to climate-related page. Orange: co-click from climate-related page to health-related page. Grey: sum of all health and climate co-click activity.....	79
Figure 26. Scientific journal articles relating to health and climate change, 2007-2020, by 2019 country Human Development Index country group.....	80
Figure 27. Proportion of countries referring to climate change, health, and the intersection between the two in their UNGD statements, 1970-2020.....	82
Figure 28. Proportion of companies referring to climate change, health, and the intersection of health and climate change in their UN Global Compact Communication on Progress (GCCOP) reports, 2011-2020. ....	85

## List of Panels

Panel 1. The Indicators of the 2021 Report of the <i>Lancet</i> Countdown.....	16
Panel 2. Gender, Health and Climate Change.....	34
Panel 3. The Urban Heat Island and the Impact of Cool Roofs.....	42
Panel 4. Recovering from Covid-19: Stimulus Measures for a Sustainable Economy.....	66
Panel 5. Compatibility of Fossil Fuel Company Strategies with Well Below 2°C-Consistent Emissions Trajectories.....	70
Panel 6. The place of Health in the Enhanced NDCs.....	82

## List of Abbreviations

A&RCC – Adaptation & Resilience to Climate Change
CDP – Carbon Disclosure Project
CFU – Climate Funds Update
CO <sub>2</sub> – Carbon Dioxide
CO <sub>2</sub> e – Carbon Dioxide Equivalent
COP – Conference of the Parties
ECMWF – European Centre for Medium-Range Weather Forecasts
EE-MRIO – Environmentally Extended Multi-Region Input-Output
EJ – Exajoule
EM-DAT – Emergency Events Database
ERA – European Research Area

ETS – Emissions Trading System  
EU – European Union  
FAO – Food and Agriculture Organization of the United Nations  
GBD – Global Burden of Disease  
GDP – Gross Domestic Product  
GHG – Greenhouse Gas  
GNI – Gross National Income  
GtCO<sub>2</sub> – Gigatons of Carbon Dioxide  
GW – Gigawatt  
GWP – Gross World Product  
HIC – High Income Countries  
HDI – Human Development Index  
IEA – International Energy Agency  
IHR – International Health Regulations  
IPC – Infection Prevention and Control  
IPCC - Intergovernmental Panel on Climate Change  
IRENA - International Renewable Energy Agency  
LMICs – Low- and Middle-Income Countries  
LPG – Liquefied Petroleum Gas  
Mt – Metric Megaton  
MtCO<sub>2</sub>e – Metric Megatons of Carbon Dioxide Equivalent  
MODIS – Moderate Resolution Imaging Spectroradiometer  
MRIO – Multi-Region Input-Output  
NAP – National Adaptation Plan  
NASA – National Aeronautics and Space Administration  
NDCs - Nationally Determined Contributions  
NHS – National Health Service  
NO<sub>x</sub> – Nitrogen Oxide  
NDVI – Normalised Difference Vegetation Index  
OECD – Organization for Economic Cooperation and Development  
PM<sub>2.5</sub> – Fine Particulate Matter  
PV – Photovoltaic  
SDG – Sustainable Development Goal  
SDU – Sustainable Development Unit  
SIDS – Small Island Developing States  
SSS – Sea Surface Salinity  
SST – Sea Surface Temperature  
tCO<sub>2</sub> – Tons of Carbon Dioxide  
tCO<sub>2</sub>/TJ – Total Carbon Dioxide per Terajoule  
TJ – Terajoule  
TPES – Total Primary Energy Supply  
TWh – Terawatt Hours  
UN – United Nations  
UNFCCC – United Nations Framework Convention on Climate Change

UNGA – United Nations General Assembly  
UNGD – United Nations General Debate  
VC – Vectorial Capacity  
WHO – World Health Organization  
WMO – World Meteorological Organization



# 1 The 2020 Report of the Lancet Countdown 2 on Health and Climate Change

## 3 Executive Summary

4 The Lancet Countdown: Tracking Progress on Health and Climate Change is an international  
5 collaboration which independently monitors the health consequences of a changing climate.  
6 Publishing updated, new and improved indicators each year, it represents the consensus of  
7 leading researchers from 38 academic institutions and UN agencies. The 44 indicators of the  
8 2021 report of the Lancet Countdown expose an unabated rise in the health impacts of  
9 climate change, and the current health consequences of the delayed and uneven response of  
10 countries around the world – providing a clear imperative for accelerated action that puts  
11 the health of people and planet first.

12 This year's report coincides with the 26<sup>th</sup> UN Framework Convention on Climate Change  
13 Conference of the Parties (COP26), a moment when countries are facing pressure to realise  
14 the ambition of the Paris Agreement to keep global average temperature rise to 1.5 °C, and  
15 mobilise the finance required for all countries to deliver an effective climate response. These  
16 negotiations unfold in the context of the COVID-19 pandemic – a global health crisis which  
17 has claimed millions of lives, affected livelihoods and communities around the globe and  
18 exposed deep fissures and inequities in the world's capacity to cope with, and respond to,  
19 health emergencies. Yet, in its response to both crises, the world is faced with an  
20 unprecedented opportunity to ensure a healthy future for all.

21

## 22 Deepening inequities in a warming world

23 Record temperatures in 2020 resulted in a new high of 3.1 billion more days of heatwave  
24 exposure among people over 65 and 626 million more exposures affecting children under  
25 1 year old, as compared to a 1986-2005 baseline (indicator 1.1.2). Looking to 2021, people in  
26 these age groups, along with those facing social disadvantages, were the most affected by the  
27 record-breaking temperatures of over 40°C that affected the Pacific northwest areas of the  
28 USA and Canada in June 2021 – an event that would have been virtually impossible without  
29 human-caused climate change. Although the tally will not be known for several months,  
30 hundreds of people prematurely died from the heat. Furthermore, populations in countries  
31 with low and medium levels of UN-defined Human Development Index (HDI) have  
32 experienced the biggest increase in heat vulnerability over the past 30 years, with risks to  
33 their health further exacerbated by the lower availability of cooling mechanisms and urban  
34 green space in these countries (indicators 1.1.1, 2.3.2 and 2.3.3).

35 Agricultural workers in countries with low and medium HDI were among the worst affected  
36 by exposure to extreme temperatures, bearing almost half the 295 billion potential work  
37 hours lost due to heat in 2020 (indicator 1.1.4). These lost work hours could have devastating  
38 economic consequences to these already vulnerable workers – data in this year’s report  
39 shows that potential earnings lost were equivalent to 4-8% of national GDP in the low HDI  
40 country group (indicator 4.1.3).

41 Through these impacts and alongside rising average temperatures and altered rainfall  
42 patterns, climate change is beginning to reverse years of progress in tackling the food and  
43 water insecurity that still affects the most underserved populations around the world,  
44 undermining a cornerstone of good health. In any given month in 2020, up to 19% of the  
45 global land surface was affected by extreme drought, a value that had not exceeded 13% from  
46 1950 to 1999 (indicator 1.2.2). In parallel, warmer temperatures are affecting the yield  
47 potential of the world’s major staple crops: a 6.0% reduction for maize; 3.0% for winter  
48 wheat; 5.4% for soybean; and 1.8% for rice in 2020, relative to 1981-2010 (indicator 1.4.1);  
49 exposes the rising risk of food insecurity in a warming world.

50 Adding to these health hazards, the changing environmental conditions are also increasing  
51 the suitability for the transmission of many water-, air-, food-, and vector-borne pathogens.  
52 Although socioeconomic development, public health interventions, and advances in medicine  
53 have reduced the global burden of infectious disease transmission, climate change threatens  
54 to undermine eradication efforts.

55 The number of months with environmentally suitable conditions for the transmission of  
56 malaria (*Plasmodium falciparum*) rose by 39% from 1950-1959 to 2010-2019 in densely  
57 populated highland areas in the low HDI country group – threatening highly disadvantaged  
58 populations, previously comparably safe from this disease due to their geographical location  
59 (indicator 1.3.1). The epidemic potential for dengue, Zika and chikungunya, which currently  
60 affect primarily populations in Central and South America, the Caribbean, Africa and south  
61 Asia, increased globally by 13% for the transmission by *A. aegypti* and 7% higher for *A.*  
62 *albopictus* from the 1950s, with the biggest relative increase was seen in countries with very  
63 high HDI (indicator 1.3.1). However, it is people in the low HDI country group who are  
64 confronted with the highest vulnerability to these arboviruses (indicator 1.3.2).

65 Similar findings are observed in the environmental suitability for *Vibrio cholerae*, a pathogen  
66 estimated to cause almost 100,000 deaths annually, particularly among populations with  
67 poor access to safe water and sanitation. Between 2003 and 2019, the coastal areas suitable  
68 for *Vibrio cholerae* transmission increased significantly across all HDI country groups – but,  
69 with 98% of their coastline suitable to the transmission of *Vibrio cholerae* in 2020, it is people  
70 in the low HDI country group that face the highest environmental suitability for this disease  
71 (indicator 1.3.1).

72 The concurrent and interconnecting risks posed by extreme weather events, infectious  
73 disease transmission, and food, water and financial insecurity are overburdening the most  
74 vulnerable. Through multiple simultaneous and interacting health risks, climate change is  
75 threatening to reverse years of progress in public health and sustainable development.

76 Even with overwhelming evidence on the health impacts of climate change, countries are not  
77 delivering an adaptation response proportionate to rising risks their populations face. In  
78 2020, 63% of all countries were yet to achieve a high level of implementation of national  
79 health emergency frameworks, leaving them unprepared to respond to pandemics and  
80 climate-related health emergencies (indicator 2.3.1). Importantly, only 55% of low HDI  
81 countries had reported at least medium-level of implementation of these frameworks,  
82 compared with 89% of very high HDI countries. In addition, only 37 of 70 countries reported  
83 having a national adaptation plan for health, with insufficient human and financial resources  
84 identified as the key barrier for their implementation (indicator 2.1.1). With a world facing  
85 unavoidable temperature rise, even under the most ambitious climate change mitigation,  
86 accelerated adaptation is essential to reduce vulnerabilities and protect the health of people  
87 around the world.

88

### 89 [An inequitable response fails everyone](#)

90 Six months into 2021, the world had failed to deliver global equitable access to the COVID-19  
91 vaccine: more than 75% of all vaccine doses had been given to people in just 10 countries.  
92 Data in this report exposes similar inequities in the global climate change mitigation response.

93 To meet the Paris Agreement goals and prevent catastrophic levels of warming, global  
94 greenhouse gas emissions must halve within a decade. However, at the current pace, it would  
95 take over 150 years for the energy system to fully decarbonise (indicator 3.1), and the unequal  
96 response between countries is resulting in an uneven realisation of the health co-benefits of  
97 a low-carbon transition.

98 Partly responsible for the slow decarbonisation rate is the use of public funds to subsidise  
99 fossil fuels. Out of 84 countries reviewed, 65 were still providing an overall subsidy to fossil  
100 fuels in 2018, using funds in many cases equivalent to substantial proportions of the national  
101 health budget, and which could otherwise be redirected to deliver net benefits to health and  
102 wellbeing. Further, all the 19 countries whose carbon pricing policies did outweigh the effect  
103 of any fossil fuels subsidies came from the very high HDI group (indicator 4.2.4).

104 While countries in the very high HDI group have collectively made the greatest progress in  
105 energy system decarbonisation, they are still the main contributors to CO<sub>2</sub> emissions through  
106 their local production, accounting for 45% of the global total (indicator 4.2.5). Meanwhile,  
107 with a slower pace of decarbonisation and poorer air quality regulations, the medium and  
108 high HDI country groups produce the most PM<sub>2.5</sub> emissions and have the highest rates of air

109 pollution-related mortality – about 50% higher than the total mortality in the very high HDI  
110 group (indicator 3.3). Turning to the low HDI country group, with comparatively lower levels  
111 of industrial activity, its local production contributes to only 0.7% of global CO<sub>2</sub> emissions, and  
112 it has the lowest mortality rate from ambient air pollution. However, with only 12% of its  
113 inhabitants relying on clean fuels and technologies for cooking, the health of these  
114 populations is still at risk from dangerously high concentrations of household air pollution  
115 (indicator 3.2). Importantly, even within the most affluent countries, people in the most  
116 deprived areas overwhelmingly bear the health burden from exposure to air pollution. These  
117 findings expose the health costs of the delayed and unequal mitigation response, and  
118 underscore the millions of lives to be saved annually through a low-carbon transition that  
119 prioritises the health of all populations.

120 However, the world is not on track to realising these health gains: current global  
121 decarbonisation commitments would lead to 2.7-3.1°C of warming by the end of the century,  
122 and present direction of post-COVID-19 spending is threatening to make this situation worse,  
123 with just 18% of all funds committed for recovery by the end of 2020 expected to reduce  
124 greenhouse gas emissions. Indeed, the economic recovery from the pandemic is already  
125 predicted to lead to an unprecedented 5% increase in GHG emissions in 2021, which will bring  
126 global anthropogenic emissions back to their peak levels.

127 In addition, the current economic recession is threatening to further undermine the target of  
128 mobilising US\$100 billion per year from 2020 onwards to promote low-carbon shifts and  
129 adaptation responses in the most underserved countries - even while this quantity is now  
130 dwarfed by the trillions allocated to COVID-19 recovery. The high levels of borrowing that  
131 lower income countries had to resort to during the pandemic could further erase their ability  
132 to deliver a green recovery, and maximise the health gains to their population of a low-carbon  
133 transition.

134

135 [An unprecedented opportunity to ensure a healthy future for all](#)

136 The overshoot in emissions resulting from a carbon-intensive COVID-19 recovery would  
137 irreversibly push the world off track meeting climate commitments and the Sustainable  
138 Development Goals – and lock in humanity to an increasingly extreme and unpredictable  
139 environment. Data in this report expose the health impact and health inequities of the current  
140 1.2°C world and confirms that, on the current trajectory, climate change will become the  
141 defining narrative of human health.

142 However, by directing the trillions of dollars that will be committed to COVID-19 recovery  
143 towards the WHO's prescriptions for a healthy, green recovery, the world could meet the  
144 Paris Agreement goals, protect the natural systems that support wellbeing, and minimise  
145 inequities through reduced health impacts and maximised co-benefits of a universal low-  
146 carbon transition. Promoting equitable climate change mitigation and universal access to

147 clean energies could save millions of lives annually from reduced exposure to air pollution,  
148 healthier diets, and more active lifestyles, and contribute to reducing health inequities  
149 globally. This pivotal moment of economic stimulus represents a historical opportunity to  
150 securing the health of present and future generations.

151 There is a glimpse of this story unfolding through several promising trends in this year's data:  
152 electricity generation from renewable wind and solar energy has increased by an annual  
153 average of 17% between 2013-2018 (indicator 3.1); investment in new coal capacity  
154 decreased by 10% in 2020 (indicator 4.2.1); and the global number of electric vehicles reached  
155 7.2 million in 2019 (indicator 3.4). Additionally, the global pandemic has driven increased  
156 engagement in health and climate change across multiple domains in society, with 91 heads  
157 of state making the connection in the 2020 UN General Debate, and newly widespread  
158 engagement among very high HDI countries (indicator 5.4). Whether COVID-19 recovery  
159 supports, or reverses these trends, is yet to be seen.

160 Neither SARS-CoV-2 nor climate change respect national borders. Without widespread,  
161 accessible vaccination across all countries and societies, the virus and its new variants will  
162 continue to put the health of everyone at risk. Likewise, tackling climate change requires all  
163 countries to deliver an urgent and coordinated response, with COVID-19 recovery funds  
164 allocated to support and ensure a just transition to a low carbon future and climate change  
165 adaptation in all corners of the world. Leaders of the world have an unprecedented  
166 opportunity to deliver a future of improved health, reduced inequity, and economic and  
167 environmental sustainability. However, this will only be possible if the world acts together to  
168 ensure no one is left behind.  
169

## 170 Introduction

171 The COVID-19 pandemic has changed societies in ways previously unimaginable, with  
172 deepening and widespread concerns about global health security, inequities, and  
173 anthropogenic influences on the environment. As of the 11<sup>th</sup> of May 2021, the pandemic had  
174 resulted in almost 191 million cases and 4.1 million deaths,<sup>1,2</sup> and its multidimensional  
175 impacts on health and wellbeing, together with its disruption to work, social, and leisure  
176 activities, still continue. The overwhelming healthcare demand caused 94 of 105 countries  
177 examined to experience disruptions to the delivery of essential health services, further  
178 undermining health and wellbeing.<sup>3</sup> Adding to this, COVID-19 led to a worldwide economic  
179 recession, an estimated 90 million people were pushed below the extreme poverty threshold  
180 in 2020,<sup>4,5</sup> and pandemic-induced borrowing by the World Trade Organization's 'developing'  
181 countries amounted to US\$130 billion by July that year.<sup>6</sup>

182 But while the world's attention has been diverted towards the ongoing acute health crisis,  
183 the health effects of human-induced climate change continue to increase. Climate change  
184 contributed to the unusually high temperatures seen during 2020 in the United Kingdom and  
185 Siberia, to the record-breaking well-over 40°C heatwave that affected populations across the  
186 Pacific northwest areas of the USA and Canada in June 2021 and caused over 1000 deaths,  
187 with that number expected to increase, to an accelerated glacier retreat that is putting the  
188 city of Huaraz (Peru) under imminent flooding risk, and to Australia's devastating 2019/2020  
189 bushfire season.<sup>7-11</sup> Over a six month period in 2020, 84 disasters from floods, droughts, and  
190 storms affected 51.6 million people in countries already struggling with COVID-19,<sup>12</sup> with the  
191 escalating impacts threatening their ability to respond to health emergencies. Meanwhile,  
192 climate impacts may undermine the capacity of countries to repay their debts, further  
193 hindering their progress towards the Sustainable Development Goals (SDGs).<sup>13,14</sup> As with  
194 COVID-19, the health impacts of climate change are inequitable, with disproportionate effects  
195 on the most vulnerable in every society, including the poor, members of minority groups,  
196 women, children, older adults, people with chronic diseases and disabilities, and outdoor  
197 workers.<sup>15</sup> Such interrelationships between climate change and COVID-19 provide ongoing  
198 evidence of the interconnectedness of the world, and of the health consequences of  
199 inequities. The 2021 report of the *Lancet* Countdown depicts the synergies and interactions  
200 between these two crises.

201 The world is now 1.2°C warmer than in the pre-industrial period, the past seven years rank  
202 as the hottest seven on record, and 2020 tied with 2016 as the hottest yet.<sup>16-18</sup> Atmospheric  
203 CO<sub>2</sub> concentrations have reached a concerning milestone – now 50% higher than in the pre-  
204 industrial era.<sup>19</sup> Changes such as reduced soil moisture could limit the Earth's carbon  
205 reuptake, resulting in further CO<sub>2</sub> in the atmosphere.<sup>20</sup> Furthermore, some climate tipping  
206 points are close to or may have surpassed critical thresholds and could interact to further  
207 destabilise the Earth's climate system.<sup>21,22</sup> While the dramatic reductions in transport and  
208 industrial manufacturing during the pandemic resulted in energy-related emissions for 2020  
209 falling by 5.8%, the largest annual percentage decline since World War II, this was short-lived

210 and emissions have risen in 2021.<sup>23-25</sup> Without an adequate response, the health effects of  
211 climate change will worsen throughout the coming decades.

212 The world now turns with hope to the 2021 UN Framework Convention on Climate Change  
213 (UNFCCC) conference in Glasgow (COP26), originally scheduled for 2020. Over the past year,  
214 the world has seen more ambitious climate targets from governments and businesses, and  
215 73% of global emissions are now covered by net zero emissions targets announced by May  
216 2021. Nevertheless, these announcements are non-binding, and even with their full  
217 implementation the world would be on track to ~2.4°C (1.9-3.0°C) of warming by 2100.<sup>26</sup>

218 These climate announcements are being made against the backdrop of huge investments in  
219 economic recovery from COVID-19. Depending on their consistency with climate targets,  
220 these investments could take the world in one of two directions – either driving it towards  
221 the goals of the Paris Agreement, or locking it into increased emissions and climate change  
222 that will damage the health of current and future generations. As humanity faces a critical  
223 turning point, the indicators in this report provide the health evidence to inform a global  
224 response to the impacts of climate change, and identify the considerable health,  
225 environmental and economic benefits that would result if a ‘green recovery’ from COVID-19  
226 was prioritised.

227

## 228 [Five years of tracking progress on health and climate change](#)

229 The *Lancet* Countdown is an independent, international, and multidisciplinary collaboration  
230 that monitors the health impacts of climate change, and progress – or lack thereof – in the  
231 world’s response. It draws on the expertise of climate scientists, economists, energy and  
232 transport experts, social and political scientists, public health experts and health professionals  
233 among others, spanning 38 academic and UN institutions. Together, they report on 44  
234 indicators, organised within five domains: climate change impacts, exposures, and  
235 vulnerabilities; adaptation, planning, and resilience for health; mitigation actions and health  
236 co-benefits; economics and finance; and public and political engagement. The 2021 report of  
237 the *Lancet* Countdown is its sixth annual report, building on nine years of collaborative work.

238 The *Lancet* Countdown’s indicator domains were selected through an open, global  
239 consultation process that identified scientifically documented links between health and  
240 climate change, with indicators developed according to well-established methods, and to the  
241 availability of reliable and regularly updated data with adequate geographical and temporal  
242 scales.<sup>27</sup> Each year, the indicators have been improved upon through an open, iterative and  
243 adaptive approach, and new indicators have been introduced to provide an increasingly  
244 complete picture of the health dimensions of climate change. For the two most recent  
245 reports, prior to the formal peer review, all new indicators underwent an independent  
246 assessment process led by world experts in their respective domains, adding rigour and  
247 transparency to the collaboration’s research. Existing indicators are undergoing a similar,

248 independent quality improvement process, aimed at ensuring they continue to use the best  
249 available data and methods.

250 Three new important indicators are added to the 2021 report: the first incorporates  
251 considerations of mental wellbeing by tracking the effect of heat on expressed online  
252 sentiment; the second captures the influence of heat on safe physical activity; and the third  
253 tracks consumption-based greenhouse gas and PM<sub>2.5</sub> emissions. Most of the pre-existing  
254 indicators underwent major improvements, with strengthened methods, datasets and  
255 metrics, and expanded geographical and temporal coverage. All indicators, including their  
256 methods, data sources, caveats, and plans for future improvements, are described in detail  
257 in the appendix – an essential manual for this report. The final indicators for the 2021 report  
258 are listed in panel 1.

259 Each indicator, wherever possible and appropriate, is disaggregated into very high, high,  
260 medium, and low Human Development Index (HDI) country groups, as defined by the United  
261 Nations Development Programme (UNDP), in the latest year for which data are available  
262 (2019).<sup>28</sup> This composite index captures three key dimensions: a long and healthy life (with  
263 life expectancy as a proxy), education (captured by the mean of years of schooling for adults),  
264 and standard of living (measured by per capita gross national income).<sup>28</sup> In line with the  
265 priorities of *The Lancet's* Diversity Board, gender disparities are also considered wherever  
266 relevant. However, a stark lack of gender-disaggregated data, means that few indicators are  
267 able to capture these differences quantitatively, and often do so using sex disaggregation as  
268 a proxy for gender (see panel 2).

269 The COVID-19 pandemic will alter the trends of many of the indicators reported – some of  
270 which can be identified in this report, and others which will become apparent in the coming  
271 years. COVID-19 has also altered population demographics and mortality rates, as well as the  
272 structure and size of the labour force. These changes are not reflected in the current  
273 indicators, presenting methodological challenges in the assessment of the health impacts of  
274 climate change. How this impacts the methods and assumptions of the *Lancet Countdown's*  
275 indicators will become clearer in future reports, as more data become available.

276 The global reach of the *Lancet Countdown* is expanding. Two regional offices – in South  
277 America (based at Universidad Peruana Cayetano Heredia, Lima, Peru) and in Asia (based at  
278 Tsinghua University, Beijing, China) – were established in 2020. These regional collaborators  
279 contributed indicators to the global 2021 report and are working on nationally- and  
280 regionally-relevant health and climate change research, accompanied by local  
281 communications and policy engagement. A third regional office, based at the University of  
282 the West Indies, was established in September 2021, aiming to build on the network and  
283 evidence base of health and climate change in Small Island Developing States (SIDS). The  
284 *Lancet Countdown* is also working in collaboration with the European Environment Agency,  
285 incorporating policy-relevant data from its indicators into the European Climate and Health  
286 observatory.



287 National and regional reports were published for Australia (in partnership with the *Medical*  
288 *Journal of Australia*), China, and SIDS.<sup>29-31</sup> For the third year now, the data underpinning each  
289 of the *Lancet* Countdown's indicators have been shared through an online data visualisation  
290 platform, where they can be explored at finer spatial and temporal scales.

291 The work of this collaboration is driven by the ongoing support from *The Lancet* and the  
292 Wellcome Trust, the *Lancet* Countdown's scientific advisory group and higher-level advisory  
293 board, and, importantly, the *Lancet* Countdown's authors and collaborators. The  
294 collaboration welcomes further offers of support from new experts and new institutions,  
295 willing to build on this analysis, as the *Lancet* Countdown monitors the world's response to  
296 the health effects of climate change across this decade.  
297

Working Group	Indicator		
Climate Change Impacts, Exposure, and Vulnerability	1.1: Health and Heat	1.1.1: Vulnerability to Extremes of Heat	
		1.1.2: Exposure of Vulnerable Populations to Heatwaves	
		1.1.3: Heat and Physical Activity	
		1.1.4: Change in Labour Capacity	
		1.1.5: Heat and Sentiment	
		1.1.6: Heat-Related Mortality	
	1.2: Health and Extreme Weather Events	1.2.1: Wildfires	
		1.2.2: Drought	
		1.2.3: Lethality of Extreme Weather Events	
	1.3: Climate-Sensitive Infectious Diseases	1.3.1: Climate Suitability for Infectious Disease Transmission	
		1.3.2: Vulnerability to Mosquito-Borne Diseases	
	1.4: Food Security and Undernutrition	1.4.1: Terrestrial Food Security and Undernutrition	
		1.4.2: Marine Food Security and Undernutrition	
	1.5: Migration, Displacement and Rising Sea Levels		
	Adaptation, Planning, and Resilience for Health	2.1: Adaptation Planning and Assessment	2.1.1: National Adaptation Plans for Health
2.1.2: National Assessments of Climate Change Impacts, Vulnerability, and Adaptation for Health			
2.1.3: City-Level Climate Change Risk Assessments			
2.2: Climate Information Services for Health			
2.3: Adaptation Delivery and Implementation		2.3.1: Detection, Preparedness and Response to Health Emergencies	
		2.3.2: Air Conditioning: Benefits and Harms	
		2.3.3: Urban Green Space	
2.4: Health Adaptation-Related Global Funding and Financial Transactions			
Mitigation Actions and Health Co-Benefits	3.1: Energy System and Health		
	3.2: Clean Household Energy		
	3.3: Premature Mortality from Ambient Air Pollution by Sector		
	3.4: Sustainable and Healthy Transport		
	3.5: Food, Agriculture, and Health	3.5.1: Emissions from Agricultural Production and Consumption	
		3.5.2: Diet and Health Co-Benefits	
	3.6: Mitigation in the Healthcare Sector		
Economics and Finance	4.1: The Economic Impact of Climate Change and its Mitigation	4.1.1: Economic Losses due to Climate-Related Extreme Events	
		4.1.2: Costs of Heat-Related Mortality	
		4.1.3: Loss of Earnings from Heat-Related Labour Capacity Reduction	
		4.1.4: Costs of the Health Impacts of Air Pollution	
	4.2: The Economics of the Transition to Zero-Carbon Economies	4.2.1: Coal and Clean Energy Investment	
		4.2.2: Employment in Low-Carbon and High-Carbon Industries	
		4.2.3: Funds Divested from Fossil Fuels	
		4.2.4: Net Value of Fossil Fuel Subsidies and Carbon Prices	
		4.2.5: Production- and Consumption-Based Attribution of CO <sub>2</sub> and PM <sub>2.5</sub> Emissions	
Public and Political Engagement	5.1: Media Coverage of Health and Climate Change		
	5.2: Individual Engagement in Health and Climate Change		
	5.3: Coverage of Health and Climate Change in Scientific Journals		
	5.4: Government Engagement in Health and Climate Change		
	5.5: Corporate Sector Engagement in Health and Climate Change		

## 299 Section 1: Climate Change Impacts, Exposures, and Vulnerability

300 Climate change threatens human health and wellbeing through impacts on weather,  
301 ecosystems and human systems – increasing exposure to extreme events, changing the  
302 environmental suitability for infectious disease transmission, altering population movements,  
303 and undermining people’s livelihoods and mental health.<sup>32-36</sup> The resulting strains on health  
304 and social systems are felt disproportionately by the most disadvantaged in society, with  
305 climate change amplifying inequities.<sup>32,33</sup>

306 Section 1 of the 2021 report monitors the health impacts of climate change, with indicators  
307 tracking climate hazards, human exposure and vulnerabilities, and the resulting health  
308 outcomes. The first group of indicators addresses the direct implications of rising  
309 temperatures for health, exploring changes in the exposure and vulnerabilities of world  
310 populations to extreme heat, as well as its impacts on health and wellbeing (indicators 1.1.1–  
311 1.1.6, see panel 1). Each of these indicators takes gridded heat data as a starting point, and  
312 overlays them with relevant exposure and vulnerability data to reflect health outcomes. Two  
313 new indicators are introduced: one examines how heat is reducing the possibility to  
314 undertake outdoor exercise safely (indicator 1.1.3); the other approaches the challenge of  
315 assessing the influence of extreme heat on sentiment, using Twitter data to capture people’s  
316 online expressions (indicator 1.1.5).<sup>37</sup>

317 The second group of indicators in this section sheds light on climate-sensitive extreme events,  
318 tracking exposure to wildfire and wildfire risk (indicator 1.2.1), the incidence of droughts  
319 (indicator 1.2.2), and the lethality of extreme weather events (indicator 1.2.3). Capturing the  
320 influence of environmental changes on ecological niches for human pathogens, the section  
321 also models the changing suitability for the transmission of climate-sensitive infectious  
322 diseases, expanding the analysis from previous years to capture three new diseases of global  
323 public health relevance (Zika, chikungunya and *Vibrio cholerae*), and improving models to  
324 reflect the reproduction number for arbovirus transmission. With health outcomes of vector-  
325 borne disease transmission strongly influenced by socioeconomic factors and healthcare  
326 access, indicator 1.3.2 incorporates considerations of implemented adaptation measures to  
327 capture the changing vulnerability to arboviruses. This is followed by indicators of  
328 environmental pressure on terrestrial and marine food productivity, this year extending the  
329 analysis to assess the association between heat stress and severe food insecurity (indicators  
330 1.4.1 and 1.4.2). The final indicator in this section focuses on exposure to sea level rise and its  
331 implications for human mobility (indicator 1.5).

332

333 1.1 Health and Heat

334 Indicator 1.1.1: Vulnerability to the Extremes of Heat

335 *Headline finding: although vulnerability to heat in the low and medium Human Development*  
336 *Index country groups remains 27-38% lower than that of the very high Human Development*  
337 *Index group, it is increasing rapidly and is today 19% and 20% higher than in 1990, respectively*

338 Exposure to extreme heat poses an acute health hazard, with individuals over 65 years of  
339 age,<sup>38-40</sup> urban populations,<sup>39,40</sup> and those with underlying health conditions<sup>38,39</sup> particularly  
340 at risk. Heat disproportionately affects the marginalised or under-resourced that have limited  
341 access to cooling mechanisms and healthcare, amplifying health and social inequities.<sup>41-44</sup>

342 This indicator tracks vulnerability to extreme heat, through an index that combines the  
343 proportion of the population older than 65 years, the prevalence of relevant chronic diseases  
344 (respiratory disease, cardiovascular disease, and diabetes) in that group, and the proportion  
345 of the total population living in urban areas.

346 With aging populations, high prevalence of chronic diseases, and increasing urbanisation, the  
347 very high HDI countries exhibited the highest vulnerability to extremes of heat. Vulnerability  
348 is rising across all HDI groups, with countries of low and medium HDI experiencing the highest  
349 increases from 1990 levels (19% and 20%, respectively). The heat indicators that follow each  
350 present worsening trends, highlighting a great need to identify populations that are  
351 vulnerable to the health impacts of heat, at the national and at the local level. Further work  
352 will be done to capture other heat vulnerabilities within this indicator.


353

354 Indicator 1.1.2: Exposure of Vulnerable Populations to Heatwaves

355 *Headline finding: in 2020, compared with the 1986-2005 baseline, children under 1 and adults*  
356 *over 65 were affected by 626 million and 3.1 billion more days of heatwave exposure,*  
357 *respectively*

358 Young children and older persons are especially susceptible to the health risks of high  
359 temperatures and heatwaves.<sup>45</sup> This indicator reports the total number of days adults aged  
360 over 65 years and (for the first time) children from birth to 1 year, were exposed to life-  
361 threatening heatwave events. In an improvement from previous years, the definition of a  
362 heatwave now aligns with the World Meteorological Organization (WMO) and other scientific  
363 literature.<sup>46-48</sup> Additional details are given in the appendix (pp 5-7).

364 Results show a steady increase in the person-days of exposure for adults over 65 years, with  
 365 the last 10 years seeing an annual average of 2.9 billion additional events and 3.1 billion more  
 366 person-days of exposure (or an average of 4.1 days per person >65 years) in 2020, with  
 367 respect to the 1986-2005 baseline (Figure 1). For children under 1 year, there were an  
 368 estimated 626 million additional person-days of exposure (4.6 days per person <1 year)  
 369 affecting this vulnerable group in 2020, compared with baseline years.



370

371 *Figure 1. Change in person-days of heatwave exposure relative to the 1986-2005 baseline. A) in the*  
 372 *population aged over 65 years; B) in the population aged under 1 year of age. In each case, the*  
 373 *countries with the highest exposure averages over the past 5 years are highlighted.*


374

375 Indicator 1.1.3: Heat and Physical Activity

376 *Headline finding: the last four decades saw an increase in the number of daily hours in which*  
377 *temperatures were too high for safe outdoor exercise, with people in the medium Human*  
378 *Development Index country group experiencing an average loss of 4.4 hours of safe exercise*  
379 *per day in 2020*

380 Physical exercise provides mental health benefits, and reduces the risk of cardiovascular  
381 disease, diabetes, cancer, cognitive decline and all-cause mortality.<sup>49-53</sup> However, high  
382 temperatures can reduce the frequency and duration of physical activity, and the desire to  
383 engage in exercise,<sup>54-56</sup> and even low levels of physical activity can pose a risk to health under  
384 high temperatures.<sup>57</sup> This indicator estimates the hours of physical activity potentially lost per  
385 person due to ambient temperature, humidity, and radiant heat, by tracking the number of  
386 hours per day during which the wet bulb globe temperature (WBGT) exceeds 28°C, a  
387 threshold above which national sports medicine authorities of the USA, Australia and Japan  
388 recommend outdoor physical activities to be conducted with discretion.<sup>58,59</sup>

389 Due to rising temperatures, the loss in the number of hours available for safe physical activity  
390 per day increased in all four country HDI groups (Figure 2). The greatest rate occurred the  
391 medium HDI country group, with an average increase from 3.5 hours per person per day in  
392 1980 to 4.4 hours per person per day in 2020.



393  
394 *Figure 2. Average potential activity hours lost per person per day by 2019 Human Development Index*  
395 *country group, 1980-2020.*

396

397 Indicator 1.1.4: Change in Labour Capacity

398 *Headline finding: in 2020 the world lost 295 billion potential work hours due to extreme heat*  
399 *exposure, with 79% of all losses in countries of low Human Development Index level occurring*  
400 *in the agricultural sector*


401

402 As well as through its direct impacts on health, high temperatures can also affect people's  
403 ability to work.<sup>60</sup> This indicator estimates the potential work hours lost as a result of heat  
404 exposure, by linking WBGT with the power (metabolic rate) typically expended by a worker  
405 within the construction, manufacturing, agriculture, and all other sectors.

406 In a rising trend, 295 billion potential work hours were lost across the globe in 2020 due to  
407 heat exposure – equivalent to 88 work hours per employed person. The three most populous  
408 countries with medium HDI levels, Pakistan, Bangladesh, and India experienced the greatest  
409 losses (2.5-3 times the world average, equivalent to 216-261 hours lost per employed person  
410 in 2020). In contrast, the three most populous countries with very high HDI levels (the USA,  
411 Japan, and Russia) accounted for the smallest numbers of labour hours lost. With lockdowns  
412 around the world, COVID-19 led to the loss of millions of hours of effective labour, particularly  
413 within service, construction, and manufacturing sectors.<sup>61</sup> The changes in labour structure  
414 induced by COVID-19 are not currently accounted for by this indicator.

415 Almost half of the total potential work hours lost globally occurred in the agricultural sector  
416 of low and medium HDI countries. Occupational heat exposure disproportionately affects  
417 labourers in the agricultural sector of low HDI countries, with 79% (25.8 out of 32.6 billion  
418 hours) of these countries' losses occurring in this sector, compared with only 12% (1.1 out of  
419 9.3 billion hours) in very high HDI countries. The impacts could therefore extend to food  
420 production. While heat affects labour capacity across all genders, differences in occupation  
421 may drive gender disparity. Men make up 80% of the total employment in the construction  
422 sector, and indigenous women and women in rural areas, who are highly dependent on local  
423 natural resources for their livelihood would be particularly affected by the impacts of climate  
424 change on labour capacity.<sup>62-64</sup>

425



426

427 *Figure 3. Heat-related potential hours of labour lost by sector and 2019 country Human Development*  
 428 *Index group, 1990-2000*

429

430 Indicator 1.1.5: Heat and Sentiment


431 *Headline finding: Exposure to heatwave events significantly worsens expressed sentiment,*  
 432 *with a 155% increase in negative expressions during heatwaves in 2020 relative to the 2015-*  
 433 *2019 average*

434 Climate change-related increases in heat extremes pose diverse risks to mental health  
 435 globally, ranging from altered affective states to elevated mental health-related  
 436 hospitalisations and suicidality.<sup>34-36,65-68</sup> However, because the definition, acknowledgement,  
 437 stigmatisation, and treatment of mental health varies across different regions and cultures,<sup>37</sup>  
 438 capturing the mental health impacts of climate change still remains a challenge that the  
 439 Lancet Countdown will work to address in upcoming years.




440 This indicator, new to the 2021 report, tracks the effect of heatwaves on the sentiment of  
441 expressions from Twitter users around the world, using previously published methods for  
442 estimating climate impacts.<sup>69-71</sup> Briefly, this indicator classifies the sentiment expressed in  
443 over six billion geolocated tweets collected between 2015 and 2020, using the Linguistic  
444 Inquiry Word Count (LIWC) sentiment classification tool.<sup>72</sup> It then deploys a multivariate  
445 ordinary least squares fixed effects model to estimate the annual effect of heatwaves on  
446 expressed sentiment. In this way, it compares sentiment expression during as-good-as-  
447 random heatwave days (as defined in indicator 1.1.2) with non-heatwave days in 40,000  
448 unique localities for nearly one million individuals per day. Potential temporal and  
449 geographical confounders were adjusted for by taking into account the month, calendar  
450 date, and location of each tweet in the analysis. Further detail is provided in the appendix (pp  
451 15-18). This indicator offers a glimpse into the influence of extremes of heat on sentiment of  
452 people around the world. However, since Twitter access and social media use are not evenly  
453 distributed, higher income countries are disproportionately represented.

454 Local heatwave exposure was found to significantly reduce positive expressions and increase  
455 negative expressions (Figure 4). In 2020, the percentage point increase in negative sentiment  
456 during a heatwave day rose to 0.20 (95% CI: 0.31-0.08), 155% higher than the 2015-2019  
457 average effect. Compared to the recent 2015-2019 baseline, the magnitude of this additional  
458 increase was substantive, equivalent to three quarters of the total rise in negative sentiment  
459 observed during a benchmark flooding event (see appendix, p 18). In parallel the reduction in  
460 positive sentiment observed during 2020 was 11.9% smaller than that observed during 2015-  
461 2019.



462



463

464

465

466

467

468

469

470

Figure 4. Heatwaves and sentiment. Top: Annual effect of heatwave exposure on the sentiment of online expressions from 2015-2020. Coloured intervals depict 95% CIs of the estimated average change in positive (green) and negative (orange) sentiment expressions during days with heatwaves, relative to the median daily maximum temperature baseline range for each location and year. Sentiment was extracted from Twitter posts using a dictionary-based approach across multiple languages. Grey bars depict the geolocated Tweet count by year of observation. Bottom: Country-level count of geolocated tweets for 2015-2020.


471

472 Indicator 1.1.6: Heat-Related Mortality

473 *Headline finding: heat-related mortality in the ≥65 population reached a record high of an*  
474 *estimated 345,000 deaths in 2019. Between 2018 and 2019, all WHO regions except for*  
475 *Europe saw an increase in heat-related deaths in this vulnerable age group*

476 Exposure to extreme heat increases risk of cardiovascular, cerebrovascular and respiratory  
477 mortality, as well as all-cause mortality.<sup>73</sup> As in the 2020 report, this indicator uses the  
478 exposure-response function and minimum mortality temperature defined by Honda and  
479 collaborators<sup>74</sup> to estimate deaths attributable to extremes of heat, with work ongoing to  
480 increase the accuracy of local estimates.<sup>74,75</sup> Using life expectancy data from the Global  
481 Burden of Disease 2019 Study,<sup>76</sup> years of life lost (YLL) were also calculated to better reflect  
482 health burdens.

483 Heat-related mortality for the 65-and-older population increased throughout the period of  
484 study, reaching a record high of almost 345,000 deaths in 2019 (Figure 5) - 80.6% higher than  
485 in the 2000-2005 average. Between 2018 and 2019, India and Brazil experienced the biggest  
486 absolute increase in heat-related mortality. Although heat related mortality fell from 2018 to  
487 2019 in the WHO European region (due to fewer attributable deaths in countries such as  
488 Germany, Russia, and the UK), this region still remains the most affected, with almost 108,000  
489 deaths attributable to heat exposure in 2019.



490

491 *Figure 5. Heat-related mortality among the 65-and-older population in 2019, by country*


492 1.2 Health and Extreme Weather Events

493 Indicator 1.2.1: Wildfires

494 *Headline finding: nearly 60% of countries saw an increase in the number of days people were*  
495 *exposed to 'very high' or 'extremely high' fire danger in 2017-2020 compared to 2001-2004,*  
496 *and 72% experienced an increased human exposure to wildfires across the same period*

497 Hotter and drier conditions caused by climate change increase the risk of wildfires and the  
498 extent of their damage.<sup>77</sup> As in previous years, this indicator tracks wildfire exposure by  
499 joining satellite-observed active fire spots,<sup>78,79</sup> as well as human exposure to high and very  
500 high climatological wildfire danger by combining areas with a Fire Danger Index score of over  
501 5 and population data.<sup>80</sup> A full description of the methods can be found in the appendix (pp  
502 22-23). This indicator does not yet quantify exposure to wildfire smoke, which can affect much  
503 greater populations and have large health consequences. For example, it is estimated that  
504 smoke from the 2019/2020 Australian fires affected 80% of Australia's population and  
505 resulted in hundreds of deaths and thousands of hospital presentations.<sup>81</sup>

506 Globally, in 2017-2020, there was an additional annual average of 215,531 person-days of  
507 wildfire exposure compared to 2001-2004. Overall, 72.4% (134 out of 185) of countries  
508 experienced an increase in wildfire exposure over this time. But the increase was unequal:  
509 83% (27 out of 32) of low HDI countries experienced an increase in wildfire exposure  
510 compared with 62.5% (40 out of 64) of very high HDI countries. The largest increases were  
511 observed in The Democratic Republic of the Congo, India, and China. Over the same time  
512 period, the climatological danger of wildfire increased in 110 countries, with the largest  
513 growth occurring in Lebanon, Gambia, and Lesotho (Figure 6).



514 *Figure 6. Annual population-weighted mean changes in days of very high and extremely high fire*  
515 *danger from 2001-2004 to 2017-2020 for each country/territory. Large urban areas with population*  
516 *density  $\geq 400$  persons/km<sup>2</sup> are excluded in the calculations of population-weighted mean values.*  
517


518 Indicator 1.2.2: Drought

519 *Headline finding: in 2020, up to of 19% of the global land surface was affected by extreme*  
520 *drought in any given month*

521 Climate change is driving an increase in the frequency, intensity, and duration of drought  
522 events. This is posing threats to water security, sanitation and food productivity, and  
523 increasing the risk of wildfires and exposure to pollutants.<sup>32,82</sup>

524 This indicator tracks the land area affected by extreme drought events using the Standardised  
525 Precipitation-Evapotranspiration Index (SPEI), capturing the changes in precipitation and the  
526 effect of temperature on evaporation and moisture loss.

527 The global land surface area affected by extreme drought conditions has consistently  
528 increased over the past 30 years. The percentage of the world’s land surface experiencing  
529 extreme drought in a given month reached a maximum of 22% in the 2010-2019 decade, a  
530 value that had only ever reached 13% in 1950-1999 (Figure 7). Furthermore, the 5 years with  
531 the most area affected have all occurred since 2015, and the Horn of Africa, a region impacted  
532 by recurrent extreme droughts and food insecurity,<sup>83</sup> was one of the most affected areas in  
533 2020.



534  
535 *Figure 7. Percentage of land area affected by drought events per month*

536 Indicator 1.2.3: Lethality of Extreme Weather Events

537 *Headline finding: the last 30 years have seen statistically significant increases in the number*  
538 *of extreme weather events, yet only the low Human Development Index country group*  
539 *experienced a statistically significant increase in the number of people affected by these*  
540 *events*

541 This indicator tracks the number of occurrences of climate-sensitive weather-related  
542 disasters, and the number of people affected and killed per event. Data is taken from the  
543 Centre for Research on the Epidemiology of Disasters,<sup>84</sup> and presented as standardised  
544 anomalies across the 1990-2020 period. The number of extreme weather events has seen a  
545 consistent and statistically significant increase in all HDI country groups over the last 30 years,  
546 with the very high HDI country group experiencing the highest increase (see appendix pp 27-  
547 31). However, only the low HDI country group experienced a statistically significant increase  
548 in the number of people affected per disaster event, a situation that might reflect greater  
549 population shifts to high-risk areas or inequities in adaptive capacity and preparedness to  
550 respond to worsening climate change hazards.

551

552 1.3 Climate-Sensitive Infectious Diseases


553 Indicator 1.3.1: Climate Suitability for Infectious Disease Transmission

554 *Headline finding: in the last decade, the area of coastline suitable for Vibrio bacterial*  
555 *transmission has increased by 35% in the Baltic, 25% in the Atlantic Northeast, and 4% in the*  
556 *Pacific Northwest. The number of months suitable for malaria transmission increased by 39%*  
557 *between 1950-1959 and 2010-2019 in highland areas of the low Human Development Index*  
558 *country group*

559 Climate change is affecting the distribution of arthropod-, food-, and water-borne  
560 diseases.<sup>85,86</sup> Together with global mobility and urbanisation, climate change is a major driver  
561 of the increase in dengue cases,<sup>87</sup> which have doubled every decade since 1990.<sup>76</sup> Other  
562 important emerging or re-emerging arboviruses, transmitted by the same vectors, are likely  
563 similarly responsive to climate change.<sup>88</sup> This indicator tracks the environmental suitability  
564 for the transmission of arboviruses (dengue, chikungunya and Zika) using an improved model  
565 to capture the influence of temperature and rainfall on vectorial capacity and vector  
566 abundance, and overlaying it with human population density data to estimate the  $R_0$  (the  
567 expected number of secondary infections resulting from one infected person). The  $R_0$  for all  
568 arboviral diseases tracked has increased rapidly from the 1950s, and, in 2020, was 13% higher  
569 for the transmission by *A. aegypti* and 7% higher for *A. albopictus*, than in baseline years  
570 (1950-1954). The largest increases in epidemic potential for dengue, Zika and chikungunya  
571 were in countries with very high HDI, mainly driven by the ongoing expansion of *Aedes*  
572 mosquitoes.

573 The influence of the changing climate on the length of the transmission season for  
574 *Plasmodium falciparum* malaria is also tracked, using a threshold-based model that  
575 incorporates precipitation accumulation, average temperature, and relative humidity.<sup>36</sup>  
576 There were significant changes in the number of months suitable for transmission of malaria  
577 in highland areas ( $\geq 1,500$  m above sea level) between 2010-2019 compared to 1950-1959,  
578 with an increase of 39% in highlands within the low HDI country group, and an increase of  
579 15% in those within medium HDI group. The difference between high and medium HDI areas  
580 is even more marked at a sub-national level. This suggests that climate change might make  
581 malaria eradication efforts increasingly difficult in already disadvantaged areas.

582 Finally, this indicator monitors the environmental suitability for the transmission of *Vibrio*  
583 bacteria in coastal waters. *Vibrio* pathogens can cause gastroenteritis and life-threatening  
584 cholera, as well as severe wound infections and sepsis.<sup>37</sup> Driven by changes in sea surface  
585 temperature and sea surface salinity, the environmental area of coastline showing suitable  
586 conditions for the transmission of non-*cholerae* *Vibrio* species at any one point during the  
587 year increased by 56% (from 7.0% to 10.9% of the coastline) in northern latitudes (40-70° N)  
588 in 2020 compared to a 1980s baseline. From the 1980s to the most recent decade, the area  
589 of coastline suitable for these bacteria at any point during the year has risen from 47.5% to  
590 82.4% in the Baltic, 29.9 to 54.9% in the Atlantic Northeast, and 1.2 to 5.1% in the Pacific  
591 Northwest (Figure 8). Between 2003 and 2019, there was an increase in the proportion of  
592 coastline with suitable conditions for *Vibrio cholerae* across all HDI country groups, with the  
593 low HDI country group being having the highest suitability on average, at 98.6% of countries'  
594 coastlines in 2019. However, it was the high HDI country group that showed the biggest  
595 increase in suitable coastline area during this period, at a rate of almost an additional 1% of  
596 their coastline becoming suitable each year ( $r^2=0.78$ ,  $df=15$ ,  $p<0.01$ ).



597  
 598 *Figure 8. Change in climate suitability for infectious diseases. Solid lines represent the annual change.*  
 599 *Straight lines represent the trend since 1950 (for dengue and malaria), 1982 (for Vibrio bacteria), and*  
 600 *2003 (for Vibrio cholerae)*

601

602 **Indicator 1.3.2: Vulnerability to Mosquito-Borne Diseases**

603 *Headline finding: while vulnerabilities to arboviruses transmitted by A. albopictus and A.*  
 604 *aegypti have decreased across all countries since the year 2000, countries in the low Human*  
 605 *Development Index group remain on average the most vulnerable*

606 As demonstrated by indicator 1.3.1, climate change is making environmental conditions  
 607 increasingly favourable for the transmission of certain arboviruses. While interventions to  
 608 reduce vulnerabilities can partly counteract these threats, environmental pressures make this  
 609 increasingly challenging. This indicator combines the environmental suitability for the  
 610 transmission of dengue (as described in indicator 1.3.1) with key indicators of social  
 611 vulnerability to this disease: access to sanitation and water services, income level, and  
 612 healthcare quality.<sup>89,90</sup>



613 Due to improvements in sanitation, income and healthcare quality, vulnerability to mosquito-  
614 borne diseases is decreasing, even despite increases in their environmental suitability.  
615 However, improvements have been slower in the lower HDI country groups: while the  
616 vulnerability for transmission by *A. aegypti* has decreased by 34% in the low HDI country  
617 groups from 2000 to 2017, the reduction in the high HDI country groups has been of 61%. The  
618 vulnerability index remains inversely related to the level of HDI, with countries in the low HDI  
619 group having a vulnerability index over 360 times higher than countries in the very high HDI  
620 group in 2017 (appendix pp 45-46).

621

## 622 1.4 Food Security and Undernutrition


### 623 Indicator 1.4.1: Terrestrial Food Security and Undernutrition

624 *Headline finding: crop yield potential continues to follow a downward trend, with 6.0%*  
625 *reduction in the crop yield potential of maize; 3.0% for winter wheat; 5.4% for soybean, and*  
626 *1.8% for rice, relative to 1981-2010 levels*

627 Food insecurity is on the rise and affected 2 billion people in 2019.<sup>91</sup> Climate change threatens  
628 to exacerbate this crisis, which will disproportionately affect the most vulnerable and those  
629 already facing undernutrition. Due to socially defined gender roles and lower empowerment,  
630 food insecurity disproportionately affects rural women, reinforcing their disadvantaged  
631 position through reduced educational attainment, income and socioeconomic status.<sup>92</sup>

632 This indicator tracks the change in crop yield potential resulting from rising temperatures  
633 using the same methods as for the 2020 report. Rising temperatures shorten the time taken  
634 for crops to reach maturity, thereby leading to reduced seed yield potential.<sup>93</sup> A reduction in  
635 crop yield potential can be considered an indicator of future crop yield reductions due to  
636 higher growing season temperatures (and therefore a shortened growing season). Crop yield  
637 potential continues to follow a consistently downward trend, adding extra pressure to already  
638 strained food systems around the world. Reductions in time to maturity are observed in all  
639 staple crops tracked, amounting to 6.0% reduction for maize, 3.0% for winter wheat, 5.4% for  
640 soybean, and 1.8% for rice, relative to 1981-2010 levels (Figure 9).

641 Data from the Food Insecurity Experience Scale of the United Nations' Food and Agriculture  
642 Organization (FAO) was used to assess self-reported experience of 'severe food insecurity',  
643 defined as a situation in which an individual went at least one whole day without eating as a  
644 result of lack of resources in the prior 12 months, in 83 countries. A fixed-effects, time-varying  
645 regression showed that every 1°C of temperature increase was associated with 1.4% increase  
646 in the probability of 'severe food insecurity' (95% CI: 1.3 to 1.47; p-value: <0.001) in 2014 and  
647 1.64% (95% CI: 1.6 to 1.65; p-value: <0.001) in 2019, globally.



648  
 649 *Figure 9. Change in crop growth duration relative to the 1981–2010 baseline. The grey line*  
 650 *represents the annual global area-weighted change. The blue line represents the running mean over*  
 651 *11 years (5 years forward and 5 years backward).*

652

653 **Indicator 1.4.2: Marine Food Security**

654 *Headline finding: in 2018-2020, nearly 70% of countries showed increases in average sea*  
 655 *surface temperature in their territorial waters compared to 2003-2005, reflecting an*  
 656 *increasing threat to their marine food productivity and marine food security*

657 Per-capita fish consumption has increased steadily since the 1960s.<sup>94</sup> About 3.3 billion people  
 658 depend on marine food, with coastal populations in low and medium HDI countries, SIDS, and  
 659 indigenous people in particular, relying on it for their nutrition and livelihoods.<sup>94,95</sup> Climate  
 660 change is driving shifts in marine fish capacity and capture through increases in sea water

661 temperatures (and the associated reduced oxygenation), ocean acidification, and coral reef  
662 bleaching. As a result of this, coastal tropical countries are facing the biggest reductions in  
663 marine yield potential, while also being the most vulnerable to the associated socioeconomic  
664 impacts.<sup>95-97</sup>

665 This indicator expands its geographical scope for 2021, tracking sea surface temperature in  
666 territorial waters of 136 countries to reflect the changing threats of climate change on marine  
667 productivity, and therefore on marine food security. It is complemented by reporting the  
668 changes in marine capture based-per-capita fish consumption, using data collected by FAO  
669 (see appendix pp 50-70).

670 Average sea surface temperature increased in the territorial waters of 95 out of 136 studied  
671 countries (70%) in 2018-2020 compared to 2003-2005, posing threats to marine food  
672 productivity. Marine capture-based fish consumption was also reduced since the 1990s,  
673 coupled with an increase in the consumption of farm-based fish products of lower nutritional  
674 quality and omega-3 content.<sup>98</sup> These trends expose the threats of climate change poses on  
675 marine food security around the world.

676

## 677 1.5 Migration, Displacement, and Rising Sea Levels

678 *Headline finding: there are currently 569.6 million people settled below 5 metres above sea*  
679 *level, who could face risks from the direct and indirect hazards posed by the rising sea levels*

680 Between 1902 and 2015, the global mean sea level increased by 0.12–0.21 metres.<sup>99</sup> If  
681 unabated, sea level rise is projected to reach up to 2 metres above current levels within 80  
682 years, or even more in certain locations if considering ice sheet collapse, waves, and tidal  
683 contributions and other factors.<sup>100-103</sup> This indicator tracks the population settled in areas at  
684 risk of global mean sea level rise, based on coastal elevation and population distribution,<sup>104,105</sup>  
685 and national policies connecting climate change, human mobility, and health.

686 There are currently 146.6 million people living in coastal areas less than 1 metre above current  
687 sea levels, 27.3% of which reside in areas with low HDI levels. Further, the 569.6 million  
688 people settled below 5 metres above current sea levels could face rising risks of increased  
689 flooding, more intense storms, soil and water salinification,<sup>106</sup> and local emergence of  
690 infectious diseases,<sup>107</sup> as sea levels continue to rise; 26.6% of these people live in areas with  
691 low HDI levels. Where erosion occurs, dwellings and other infrastructure can be damaged.

692 Migration and mobility could be a response, which might be exacerbated through other  
693 impacts of climate change, like those described in other indicators in this section. This would  
694 affect livelihoods, access to essential services, and psychosocial wellbeing.<sup>108-110</sup> Up to  
695 December 31, 2020, 45 policies connecting climate change and migration were identified in  
696 37 countries (see appendix pp 72-77), all of which mentioned health or wellbeing, but

697 typically related to climate change impacts rather than to the potential health impacts of  
698 forced migration. Although they commonly accepted that mobility could be domestic and  
699 international, immobility was rarely acknowledged. National policies that recognise and  
700 respond to the health risks and health benefits of different mobility patterns will, in part,  
701 shape the overall health outcomes.<sup>111</sup>

702

---

703 *Panel 2. Gender, Health and Climate Change*

---

704 The health impacts of climate change are both underpinned and amplified by gender norms and gender  
705 inequities, with numerous examples cited throughout this report.<sup>112</sup> Gender also influences who sets the agenda  
706 and drives responses to climate change. Evidence shows that greater representation of women in parliament is  
707 associated with stronger climate change policies.<sup>113-115</sup> However, only 21% (41 out of 196) of the heads of  
708 delegation to the UNFCCC Conference of Parties in 2019 were women, and women headed just 29% of national  
709 delegations to the UNFCCC intersessional in June 2019. Moreover, of the 1,000 scholars listed by Reuters as the  
710 most influential on climate change, only 122 were women.<sup>116</sup>

711 There is an urgent need for gender-sensitive responses to the health dimensions of climate change. This must  
712 be underpinned by the collection and reporting of data that is sufficiently disaggregated, granular and  
713 intersectional to reveal local inequities – for example data disaggregated not only by gender but also by  
714 geography, age, ethnicity, class, and other markers of marginalisation and vulnerability.<sup>117-121</sup> However, in many  
715 cases a severe lack of standardised, gender-disaggregated data hampers these efforts,<sup>122-126</sup> and it is the very  
716 social structures that shape how gender is perceived and prioritised that undermine progress: cultural norms  
717 often translate into weak political and financial support, and limit the capacity of researchers to engage with  
718 gender inequities.<sup>124,127</sup> Only 6% of all scientific articles covering climate change and health in 2020 also  
719 considered gender (indicator 5.3), and despite a workstream established for this purpose, only 6 of the 44  
720 indicators in the 2021 report of the *Lancet* Countdown provide data by sex or gender.

721 Starting to reverse this, the United Nations Entity for Gender Equality and the Empowerment of Women (UN  
722 Women) is leading global efforts to increase the availability of information on gender through its “Making Every  
723 Woman and Girl Count” flagship programme. Through this programme, UN Women supports countries with the  
724 development of priority indicators to capture gender inequities – both through indicator selection and through  
725 data collection.<sup>128</sup> A model questionnaire has been developed for that purpose, and several countries, including  
726 Bangladesh, Mongolia and several Pacific Island countries, have either begun (or are currently preparing for)  
727 their rollout. With the purpose of helping countries understand the connections between the environment and  
728 gender equality, the programme also supports data reprocessing, and the integration of geospatial information  
729 with demographic and health surveys. The importance of this work is already materialising: preliminary analysis  
730 demonstrates the accentuation of gender inequities as a result of weather events, including drought episodes  
731 driving spikes in child marriage for girls in almost all Asian countries analysed.

732 Gender, as a social construction, affects everyone in society.<sup>9-13</sup> A gender-sensitive response to climate change  
733 would generate benefits for the whole of society. Ensuring gender is represented in national statistical strategies  
734 and regular data collection processes will expose the true dimensions of the challenge. This, along with more  
735 diverse leadership, will inform and drive a commensurate response.

---

## 736 Conclusion

737 In this fifth iteration of the *Lancet* Countdown indicators, section 1 of the 2021 report  
738 highlights a continuous increase in the impacts of climate change on all monitored aspects of  
739 human health, providing further evidence that climate change is having quantifiably and  
740 increasingly negative impacts on human health.

741 While its health impacts are felt across the world, climate change disproportionately affects  
742 disadvantaged populations, exacerbating their vulnerabilities. The stratification of indicators  
743 by HDI groups reveals the higher risks faced by low and medium HDI countries, particularly  
744 with regards to labour capacity and livelihoods, food security, and vector-borne disease  
745 transmission. Capturing the health impacts on disadvantaged groups, necessary for  
746 adaptation responses (described in the following section), represents a major challenge,  
747 exacerbated by the lack of disaggregated data.<sup>15</sup> With respect to gender, this is further  
748 explored in panel 2. Moreover, although this section considers the impact of heat on  
749 sentiment, the difficulties of capturing the mental health impacts of climate change still  
750 remain. Work will continue to focus on addressing this gap.

## 751 Section 2: Adaptation, Planning, and Resilience for Health

752 The past year affirmed the centrality of health and wellbeing to socioeconomic development,  
753 illustrating how health risks can compound and cascade across other sectors and nations, and  
754 dramatically highlighting the potential consequences of chronic, limited investments into  
755 climate-resilient and environmentally sustainable health systems.<sup>129,130</sup> The COVID-19  
756 pandemic also exposed stark differences in the capacity of health systems and the resilience  
757 of populations to health emergencies,<sup>4,5</sup> identifying the urgent need for health authorities to  
758 increase national and international coordination and preparedness. This should include  
759 integrated surveillance and monitoring of emerging health threats, developing and deploying  
760 early warning and response systems, and financially supporting low-resource nations and  
761 communities.<sup>131</sup> Importantly, for the public health response to be effective, it must address  
762 the needs of the most vulnerable – with the benefits of reduced inequities for the whole of  
763 society.

764 Building climate-resilient and environmentally sustainable health systems would not only  
765 help reduce the health impacts of climate change explored in the previous section, but also  
766 contribute to minimising the risk of future pandemics. This section reports eight indicators of  
767 adaptation, planning, and resilience, closely linked with the components of the WHO  
768 Operational Framework for Building Climate Resilient Health Systems: planning and  
769 assessment (indicators 2.1.1–2.1.3); information systems (indicator 2.2); delivery and  
770 implementation (indicators 2.3.1–2.3.3); and funding and spending (indicator 2.4). Each of  
771 these indicators provide insights into inequities. Data on health adaptation funding from  
772 global financing mechanisms – necessary to help low and medium HDI countries adapt to the  
773 worsening health impacts of climate change – have been reintroduced into this year’s report  
774 (indicator 2.4).

775 A remaining challenge within section 2 is the scarcity of clear metrics to monitor adaptation  
776 progress. While efforts were made to validate the indicators, self-reported data for  
777 adaptation plans, assessments, and services may be subject to reporting bias, particularly  
778 where COVID-19 resulted in redeployment of public health resources, and where surveys  
779 experienced a decline in participation.

780

### 781 2.1: Adaptation Planning and Assessment

#### 782 Indicator 2.1.1: National Adaptation Plans for Health

783 *Headline finding: in 2021, 37 countries out of 70 reported having national health and climate*  
784 *change strategies or plans in place*

785 Health systems are under pressure to respond to the acute and long-term threats from  
786 climate change, while simultaneously facing other critical public health risks. Comprehensive,

787 implemented health adaptation plans can not only improve health resilience to climate  
788 change, but also contribute to broader health systems strengthening, and catalyse effective  
789 collaboration with other health-determining sectors.

790 Data for indicators 2.1.1 and 2.1.2 are sourced from the 2021 WHO Health and Climate  
791 Change Global Survey,<sup>132</sup> that provides self-reported data on health sector response to  
792 climate change from 70 governments (described in the appendix pp 78-79). This indicator  
793 tracks the development of national health and climate change strategies and plans and  
794 barriers to implementation.

795 In the 2021 survey just over half of countries (37 out of 70) reported to have a national health  
796 and climate change strategy or plan in place, comparable to the proportion reported in 2018.  
797 Implementation remains a challenge for countries from all HDI levels with less than a quarter  
798 of these countries reaching high or very high levels of implementation. Insufficient financing  
799 was identified as a main barrier to reaching full implementation by 73% of all responding  
800 countries with one-fifth (8 out of 37) reporting to have no current sources of funding available  
801 for taking action on priorities set out in their strategies/plans. Other key barriers to  
802 implementation were insufficient human resource capacity (59%), COVID-19 related  
803 constraints (54%), and a lack of research and evidence (49%).

804 A desktop review of National Adaptation Plans (NAPs) submitted to the UNFCCC found that  
805 four of the 19 NAPs considered gender in health adaptation actions. However, although NAPs  
806 may mention the principles of gender equality, they often fail to demonstrate mainstreaming  
807 gender in a way that challenges gender norms, power, and structures. The recommendations  
808 in the *WHO Guidance for Mainstreaming Gender in Health and Climate Change Programmes*  
809 provide countries with guidance for achieving gender mainstreaming, including through  
810 national health and climate change plans.<sup>133,134</sup>

811

812 [Indicator 2.1.2: National Assessments of Climate Change Impacts, Vulnerability, and](#)  
813 [Adaptation for Health](#)

814 *Headline finding: 36 out of 70 countries in 2021 reported having conducted a climate change*  
815 *and health vulnerability and adaptation assessment*

816 Evidence-based policy development and planning require a comprehensive evaluation of the  
817 climate change-associated health risks faced by populations and health systems. This  
818 indicator monitors the number of countries who report having conducted a climate change  
819 and health vulnerability and adaptation assessment. These assessments are critical, because  
820 they not only allow countries to establish and re-evaluate health risks, but also consider the  
821 vulnerabilities contributing to health outcomes.

822 While 36 out of 70 countries disclosed they had conducted a climate change and health  
823 vulnerability and adaptation assessment, only about one-third of these reported that the  
824 findings influenced the allocation of human and financial resources. However, 56% (20 out of  
825 36) reported that the findings informed the development of their national health and climate  
826 change strategy/plan, suggesting evidence-based policy setting. Over two-thirds of countries  
827 specifically considered vulnerable population groups in their assessments, including children,  
828 women, the elderly, workers, rural/urban populations, those living in poverty and, to a lesser  
829 extent, indigenous groups, migrant or displaced populations. However, the  
830 comprehensiveness of these assessments varied widely.

831 As explored in section 1, health vulnerabilities to climate change are unevenly distributed and  
832 can exacerbate existing health inequities. As health vulnerability and adaptation assessments  
833 inform national health and climate change plans and programmes, it is essential that data  
834 gathered for these assessments are disaggregated according to social determinants of health.  
835 This will enable public health interventions to actively identify and support the most  
836 vulnerable communities, and proactively reduce sub-national health inequities relating to  
837 climate change.

838

### 839 [Indicator 2.1.3: City-level Climate Change Risk Assessments](#)

840 *Headline finding: in 2020, 546 of 670 cities reported having completed or being in the process*  
841 *of undertaking climate change risk assessments. Heat-related illness was the most common*  
842 *climate-related health concern, identified by 169 out of 308 cities*

843 The COVID-19 pandemic revealed the persistent health inequities and vulnerabilities of cities  
844 and urban sub-populations.<sup>135,136</sup> Home to over half the world's population (a proportion  
845 projected to increase to 70% by 2050), cities play a crucial role in leading local health  
846 adaptation to climate change.<sup>137</sup> Using data from the CDP's 2020 survey of global cities, this  
847 indicator captures the number of cities that report having completed a climate change risk or  
848 vulnerability assessment; and the climate-related health impacts and vulnerabilities that  
849 cities identified.

850 In 2020, 546 of 670 cities (81%) reported they had completed or were currently undertaking  
851 climate change risk assessments. For those cities who responded in both 2019 and 2020, an  
852 additional 45 (9%) reported having completed a climate change risk assessment in 2020.  
853 Importantly, however, 94% of responding cities belonged to countries with high or very high  
854 HDI, meaning that cities and countries of low and medium levels of HDI were  
855 underrepresented in the data. 308 cities identified that climate change poses a threat to one  
856 or more health areas. The most prominent perceived health concern pertained to heat-  
857 related illness, with 169 (55%) cities reporting this concern. The most vulnerable groups  
858 identified were the elderly (reported by 213 [69%] cities), children and youth (180 [58%]), and



859 low-income households (170 [55%]), while 94 cities (31%) identified women as vulnerable to  
860 climate-related health impacts.

861

## 862 Indicator 2.2: Climate Information Services for Health

863 *Headline finding: in 2020, national meteorological and hydrological services of 86 countries*  
864 *reported providing climate information to the health sector; only five out of the 86 indicated*  
865 *these climate services guide health sector policy and investment plans*

866 Health adaptation to climate change relies on accurate meteorological data and forecasts for  
867 the integrated surveillance and monitoring of emerging health threats, the development and  
868 deployment of early warning and response systems, and the implementation of adaptation  
869 interventions. This indicator monitors the extent to which national health and meteorological  
870 services provide climate information services to the health sector, using data reported to the  
871 World Meteorological Organization (WMO).

872 In 2020, 86 national meteorological and hydrological services reported providing climate  
873 services to the health sector. In very high HDI countries, 50% of those providing services to  
874 the health sector reported that they were co-designing or providing tailored climate  
875 information services or products, in contrast with 36% of low HDI countries.

876

## 877 2.3: Adaptation Delivery and Implementation

### 878 Indicator 2.3.1: Detection, Preparedness and Response to Health Emergencies

879 *Headline finding: 124 out of 166 countries reported medium-to-high implementation of a*  
880 *national health emergency framework in 2020; an increase of 14% compared to 2019*

881 The International Health Regulations (IHR) are legally-binding instruments defining countries'  
882 rights and obligations in handling public health events and emergencies that could cross  
883 national borders.<sup>11</sup> Under the IHR, State Parties are required to provide self-evaluations of  
884 emergency response preparedness against 13 core capacities published in the State Party  
885 Annual Report (SPAR). Limitations of the IHR in ensuring an effective response to the COVID-  
886 19 pandemic were identified and continue to be evaluated, and reviews currently underway  
887 are discussed in the appendix (pp 89-90). Notwithstanding, countries with higher SPAR scores  
888 had lower incidence and mortality per 100,000 population within 30 days from first COVID-  
889 19 diagnosis, stressing the relevance of the IHR.<sup>138</sup>

890 This indicator tracks the degree to which countries have implemented a national health  
891 emergency framework under IHR core capacity eight, which includes emergency

892 preparedness and response planning, emergency management structures, and mobilisation  
893 of resource. This assesses whether countries are prepared and operationally ready to respond  
894 to all public health events, including climate-related emergencies. In 2020, 166 (85%) of State  
895 Parties to the IHR completed the relevant section of the SPAR relating to capacity eight, and  
896 75% of countries reported medium-to-high degrees of implementation of a national health  
897 emergency framework – a 14% increase compared to 2019. Importantly, however, only 37%  
898 of countries reported high implementation, indicated by a capacity score of 75% or greater.  
899 The level of implementation varied greatly by HDI, with 89% of very high HDI countries  
900 reporting medium-to-high implementation, compared to 55% of low HDI countries.

901 In preparing for future health crises, it is essential that global institutions improve emergency  
902 response preparedness, using lessons learned during the pandemic. The ongoing review of  
903 the IHR is an important step in this direction to ensure that the IHR is effective when faced  
904 with health emergencies associated with climate change.

905

#### 906 [Indicator 2.3.2: Air Conditioning: Benefits and Harms](#)


907 *Headline finding: use of air conditioning, a widespread technology for indoor cooling in some*  
908 *regions of the world, averted an estimated 195,000 heat-related deaths among people ≥65*  
909 *years of age in 2019. However, it also contributed to greenhouse gas emissions, air pollution,*  
910 *peak electricity demand, and urban heat islands*

911 Indoor cooling represents an effective strategy for preventing heat-related mortality.<sup>139</sup> In  
912 this year's report, this indicator combines the prevented fraction<sup>140</sup> and heat-related death  
913 estimates from indicator 1.1.6, to track the number of heat-related deaths averted by air  
914 conditioning in the 65-and-older population (methods described in appendix pp 91-101).

915 Applying country- and region-specific prevented fractions to the data from indicator 1.1.6  
916 revealed that, in the absence of air conditioning, an estimated 195,400 heat-related deaths  
917 would have occurred globally in the 65-and-older population, in addition to the 345,000 heat-  
918 related deaths that are estimated to have occurred in 2019. In this age group, air conditioning  
919 averted an estimated 69,500 deaths in China (where 72,000 deaths attributable to heat  
920 exposure are estimated to have occurred; 65% of households had air conditioning), 47,800 in  
921 the USA (where 20,500 deaths are estimated to have occurred; 92% of households had air  
922 conditioning), 30,400 in Japan (where 12,400 deaths are estimated to have occurred; 93% of  
923 households had air conditioning), but only 2,400 in India (where 46,600 deaths are estimated  
924 to have occurred; 6% of households had air conditioning). These figures demonstrate the  
925 power of indoor cooling to prevent mortality, as well as the inequities in access to indoor  
926 cooling across countries.

927 Current air conditioning technology is unsustainable, and leads to adverse health outcomes  
 928 from increased air pollution, urban heat, and greenhouse gas emissions (see panel 3).<sup>141</sup> In  
 929 2019, the number of premature deaths from PM<sub>2.5</sub> exposure attributable to fossil-fuel  
 930 powered electricity used for air conditioning is estimated (using the same approach as in  
 931 indicator 3.3) to have been 21,000 globally. Between 2000 and 2019, the global proportion of  
 932 households with air conditioning rose 57% and CO<sub>2</sub> emissions from air conditioning use rose  
 933 61% (Figure 10).

934 Sustainable indoor cooling approaches are urgently needed, including strong, enforced codes  
 935 that mandate energy-efficient buildings,<sup>141</sup> a return to traditional tropical and sub-tropical  
 936 building designs in these regions and elsewhere,<sup>141</sup> use of fans in climate zones where they  
 937 provide effective cooling,<sup>142</sup> stringent minimum energy performance standards for air  
 938 conditioners,<sup>141</sup> cool roofs (see panel 3), and increased urban green space (Indicator 2.2.3).



939  
 940 *Figure 10. Global heat-related deaths averted by household air conditioning in the 65-and-older*  
 941 *population (red line), proportion of households with air conditioning (blue line), and carbon dioxide*  
 942 *emissions from air conditioning (green line), 2000-2019*

943

---

944 *Panel 3. The Urban Heat Island and the Impact of Cool Roofs*

---

945 As a result of human activity and the urban fabric, cities tend to be hotter than surrounding rural or suburban  
946 areas, a phenomenon known as the urban heat island (UHI) effect.

947 With increasing temperatures and urbanisation, the demand for cooling mechanisms is on the rise. While  
948 offering protection from life-threatening extreme heat exposure, the use of air conditioning contributes to  
949 climate change through its energy consumption and its leakage of hydrofluorocarbons that act as powerful  
950 greenhouse gases; contributes to intensifying the urban heat island through its waste heat emissions; and  
951 contributes to increasing peak electricity demand and urban air pollution (see indicator 2.3.2).<sup>143-145</sup>  
952 Furthermore, its high costs are amplifying the energy poverty gap.<sup>145,146</sup> The development of sustainable and  
953 affordable cooling alternatives is therefore crucial to protect the health of urban populations, while keeping the  
954 world on track to meeting the Paris Agreement goals.

955 This case study explores the use of ‘cool’ (reflective) roofs as sustainable cooling mechanisms, ranging from  
956 specially designed roofing materials, to affordable alternatives such as light-coloured paint. Focusing on  
957 Birmingham and the West Midlands region of the UK, urban air temperatures were simulated at 1 km x 1 km  
958 horizontal resolution by combining detailed land use data with a building energy parameterisation scheme in a  
959 regional climate model (WRF).<sup>147</sup> To estimate the impact of the UHI, temperatures are compared with those  
960 from a simulated counterfactual scenario, with urban surfaces replaced by rural types.

961 The UHI intensity was found to be around 3°C during summer, and up to 9°C during heatwaves in this region.  
962 The resulting overheating was estimated to contribute to approximately 40% of mortality associated with UHI  
963 over a summer season, and up to 50% during heatwaves.<sup>148-150</sup> Spatial analysis further revealed that the most  
964 underserved population groups were particularly exposed to urban heat.<sup>150</sup>

965 Simulations introducing reflective surfaces found that cool roofs could reduce maximum daytime air  
966 temperatures by 0.5°C on average, and up to 3°C during heatwaves. This has the potential of reducing heat-  
967 related mortality due to the UHI by 18% over a summer season, and 23% during a heatwave.<sup>151</sup> Considering this  
968 assessment was done in a country with relatively cool climate, the impact of cool roofs might be even greater if  
969 applied in warmer parts of the world. Moreover, while the UHI can reduce cold-related mortality by around 15%  
970 in the winter, cool roofs were shown to have negligible effects in winter months, suggesting they would not  
971 contribute to increased mortality in the winter.<sup>152,153</sup>

972 Because roofs may affect other factors such as precipitation, their use must be assessed on a case-by-case  
973 basis.<sup>154</sup> However, with a net annual benefit on temperature-related mortality, adoption of cool roofs in the face  
974 of a warming world could provide a low-carbon cooling alternative, with health benefits to the whole urban  
975 population.

---

976 Indicator 2.3.3: Urban Green Space

977 *Headline finding: globally in 2020, 27% of urban centres were classified as being moderately*  
978 *green or above, an increase from 14% in 2010. This level of greenness varied between 17% of*  
979 *urban centres in the low Human Development Index country group and 39% of urban centres*  
980 *in the very high country group*


981 There is increasing evidence that access to urban green spaces provides benefits to human  
982 physical and mental health. This includes reducing exposures to air pollution, relieving stress,  
983 and increasing social interaction and physical activity, with overall improved general health  
984 outcomes and lower mortality risk.<sup>155,156</sup> Green space also helps climate change mitigation  
985 and adaptation by sequestering carbon and delivering local cooling benefits. However, urban  
986 green spaces must be carefully designed and managed to conserve biodiversity, and ensure  
987 they do not provide habitats and breeding sites for vectors of human diseases, or contribute  
988 to increased gender and other social inequities.<sup>157-163</sup>

989 Indicator 2.3.3 provides an estimate of the magnitude of green vegetation in urban centres,  
990 using the satellite-based Normalized Difference Vegetation Index (NDVI), with higher values  
991 indicating higher greenness levels. In the 2021 report, the sample size was increased to  
992 include 1,029 urban centres across 139 countries. These encompass all urban centres of over  
993 500,000 inhabitants, as well as the most populated one in those countries that had no urban  
994 centre above this threshold. Full details are in the appendix (pp 102-106).


995 Averaged across all urban centres sampled, population-weighted peak NDVI increased 23%  
996 from 2010 to 2020 (mean NDVI 0.26 to 0.32), with 27% of urban centres being classified as  
997 moderately green or above (an NDVI  $\geq 0.40$ ) in 2020 (Figure 11). The level of greenness varies  
998 greatly by HDI level. In the very high HDI country group, 39% of urban centres have at least  
999 moderate levels of greenness (mean NDVI 0.34) in 2020, compared to 17% (mean NDVI 0.27),  
1000 36% (mean NDVI 0.33), and 15% (mean NDVI 0.30) in low, medium and high HDI country  
1001 groups, respectively. This highlights the inequities in the availability of green spaces across  
1002 urban centres.

1003 With their potential to simultaneously improve health outcomes, reduce health inequities,  
1004 and facilitate climate mitigation and adaptation, urban green space design must involve  
1005 interdisciplinary experts to ensure the health and environmental benefits are maximised.<sup>164</sup>  
1006 More broadly, with health at the centre of planning in areas such as housing, transport,  
1007 energy, and water and sanitation, urban centres can be places that are safe, comfortable, and  
1008 enjoyed by everyone.<sup>165</sup>

A



B



1009 *Figure 11. Average urban population-weighted peak Normalized Difference Vegetation Index (NDVI)*  
1010 *in urban centres of >500,000 inhabitants by country, for 2010 (A) and 2020 (B). For countries without*  
1011 *an urban centre of >500,000 inhabitants, the most populated urban centre was used in the analysis.*

1012

1013 **Indicator 2.4: Health Adaptation-Related Global Funding and Financial Transactions**


1014 *Headline finding: globally, adaptation funding directed at health systems represents a small*  
1015 *portion of total climate change adaptation funding (0.3%), and only 5.6% of all transactions*  
1016 *with adaptation potential were relevant to health in 2019/20*

1017 This indicator monitors two elements of spending that could provide adaptation for health:  
1018 1) the global funding approved for health-related adaptation projects through multilateral  
1019 funds, and 2) global financial transactions with the potential to deliver adaptation in the  
1020 health and care sector, as well as in other sectors that are relevant to the determinants of  
1021 health. The former draws on data from the Climate Funds Update Data Dashboard, while the  
1022 latter uses the Adaptation and Resilience to Climate Change (A&RCC) dataset produced by  
1023 kMatrix.<sup>166,167</sup> These complementary elements provide an evaluation of proactive adaptation  
1024 funding potentially related to health, and of the global size of all economic transaction that  
1025 can offer climate change adaptation potential for health.

1026 Between 2018 and 2020, \$5.1 billion of multilateral climate change adaptation funding was  
1027 approved globally. Only \$711 million (13.9%) was related to health. This consisted of \$14.0  
1028 million (0.3%) of approved funding directed specifically at health systems, and \$697 million  
1029 (13.6%) with potential secondary benefits for health identified.

1030 Meanwhile, the value of all financial transactions with the potential to deliver adaptation for  
1031 health (adaptation-relevant transactions within the dataset-defined “health and healthcare”  
1032 sectors) increased by 14.0% in 2019/20 compared with 2018/19, reaching 5.6% of total  
1033 adaptation spending. Spending in other sectors that could be relevant to health (including in  
1034 the waste and water management, built environment, or agricultural sectors, for example) is  
1035 estimated to have increased by 7.6%, representing 28.6% of total transactions. Grouped by  
1036 HDI, \$234 million (1%) of spending was within low HDI countries (Figure 12). This compares  
1037 to \$1.8 billion (8%) in medium HDI countries, \$5.7 billion (27%) in high HDI countries, and  
1038 \$13.3 billion (64%) in very high HDI countries. For spending in health-relevant economic  
1039 sectors, a similar narrative emerges: \$1.2 billion (1%) of spending occurred in low HDI  
1040 countries, compared with 62% in countries with very high HDI. As the data covers financial  
1041 years, the data up to 31<sup>st</sup> of March 2020 presented in this indicator are unlikely to reflect the  
1042 anticipated economic impact of the COVID-19 pandemic on adaptation spending.

1043 These findings highlight a growing global market for health-relevant adaptation transactions,  
1044 but this has yet to translate into sufficient targeted health adaptation funding. As world  
1045 economies recover from COVID-19, sufficient resources must be redirected towards health  
1046 adaptation to build resilience to the increasing health threats of climate change.



1048

1049 *Figure 12. Per capita potential adaptation transactions in the health and health care sector (A) and*  
 1050 *health-relevant sectors (B) for financial years 2015/16 to 2019/20, by 2019 Human Development*  
 1051 *Index country group*

1052

## 1053 Conclusion

1054 The indicators in this section paint a complex landscape of adaptation, planning, and  
 1055 resilience for health in the past 12 months, where the small global improvements to  
 1056 adaptation planning and assessment (indicators 2.1.1, 2.1.2, and 2.1.3) and intersectoral  
 1057 collaboration (indicator 2.2) are overshadowed by slow progress in implementation  
 1058 (indicators 2.3.2, 2.3.3) and insufficient investment (indicator 2.4). A key theme across all the  
 1059 indicators is inequities, and while these indicators largely track inequities between countries,  
 1060 within-country inequities are significant for moving towards resilience and sustainability.

1061 While the world economy and health systems are on the road to recovery from a significant  
 1062 acute global health crisis, climate change poses a much greater health threat in the coming  
 1063 decades. It is crucial that organisations and institutions capitalise on the insights generated  
 1064 from the pandemic to improve adaptability and resilience. Research is needed to identify  
 1065 current and future vulnerabilities; project risks from climate change at scales relevant for  
 1066 decision-making under different climate and development scenarios; and identify and  
 1067 evaluate adaptation options to prepare for and protect health in a changing climate.  
 1068 Adaptation plans should be reviewed and updated to consider medium and long-term risks  
 1069 of climate change for health, and to further build resilience. Greater collaboration and



1070 coordination are necessary across public and private sectors and global institutions, along  
1071 with increasing investments in adaptation.  
1072

## 1073 Section 3: Mitigation Actions and Health Co-Benefits

1074 Global atmospheric CO<sub>2</sub> levels passed 415 ppm in January 2021 – continuing an unbroken  
1075 upward trend – and for the first time, the concentrations for much of 2020 are expected to  
1076 be 50% higher than the 1750-1800 average.<sup>169</sup> Total emissions of all greenhouse gases in 2019  
1077 were 59.1 GtCO<sub>2</sub>e (±5.9) including those generated by land-use changes. To limit warming to  
1078 1.5°C, annual global emissions must be reduced to 25 GtCO<sub>2</sub>e by 2030.<sup>168</sup>

1079 COVID-19 and associated lockdowns across the globe have had profound impacts on the  
1080 global economy – most significantly in the surface and air transportation, and industrial  
1081 sectors.<sup>169</sup> Emissions from very high HDI countries, which account for 48% of the global total,  
1082 were around 10% lower than 2019 levels.<sup>169</sup> However, without targeted intervention,  
1083 emissions will rebound as the world recovers from the pandemic. Indeed, the 5.8% drop in  
1084 energy-related CO<sub>2</sub> emissions seen in 2020 is forecast to be matched with an unprecedented  
1085 4.8% rise in 2021.<sup>25</sup>

1086 The necessity of steering the economic recovery to a lower-emissions pathway has been well  
1087 publicised but has yet to be well-integrated into recovery plans (see panel 4).<sup>170</sup> Nevertheless,  
1088 the COVID-19 recovery presents the challenge and simultaneous opportunity to encourage  
1089 action that yields benefits to health.

1090 Tracking this global challenge, section 3 covers the relationships between climate change  
1091 mitigation actions and health. It provides an overview of the global energy system (indicator  
1092 3.1) alongside associated global exposure to ambient PM<sub>2.5</sub> air pollution and its health impacts  
1093 (indicator 3.3). Energy use in the home is also reported, with new detail on fuels used and  
1094 estimates of indoor air pollution concentrations (indicator 3.2). Following this, individual  
1095 sectors are examined: transport (indicator 3.4); food and agriculture (indicators 3.5.1 and  
1096 3.5.2); and the global healthcare sector (indicator 3.6). Where possible, the ways in which  
1097 relationships between health and climate change mitigation both influence and are  
1098 influenced by societal inequities are explored.

1099

### 1100 Indicator 3.1: Energy System and Health

1101 *Headline finding: from 2014 to 2018, despite strong growth in renewables in very high Human*  
1102 *Development Index countries, the carbon intensity of the global energy system has seen an*  
1103 *annual average decline of just 0.6% - a rate incompatible with meeting the ambitions of the*  
1104 *Paris Agreement*

1105 Fossil fuel combustion within the energy system is the largest single source of greenhouse gas  
1106 emissions, with a global share of 65%.<sup>168</sup> The rapid shift from coal to renewable energy use is  
1107 crucial, not only to mitigate these emissions, but also to prevent deaths due to ambient air


1108 pollution (indicator 3.3) and eliminate other harmful pollutants related to coal mining and  
1109 combustion.<sup>171</sup> Drawing data from the IEA, this indicator tracks three components: the carbon  
1110 intensity of the global energy system; coal phase-out; and zero-emission electricity. Full  
1111 details are described in the appendix (pp 110-115).

1112 The carbon-intensity of the global energy system fell slightly for the fifth year in a row, to 56.0  
1113 tCO<sub>2</sub>e/TJ (excluding land use emissions) in 2018. However, progress remains very limited, with  
1114 an annual rate of decline of just 0.6% from 2014 to 2018. At this rate it would take over 150  
1115 years to fully decarbonise the energy system – far from the 2040 deadline required to keep  
1116 temperature rise to 1.5°C.<sup>172</sup> Progress has been made in the very high HDI country group since  
1117 1970 and carbon intensity in the high HDI country group could be at a possible peak. However,  
1118 driven by the need to develop, the low and medium HDI country groups have shown sustained  
1119 growth in emissions per unit of energy over the period (Figure 13).

1120 China continues to dominate global coal consumption – although it represents 18.1% of the  
1121 world's population, it accounted for 53% of global coal use in 2019. While global coal use for  
1122 all activities fell 1.2% in 2019, including a fall of 13.4% in the USA and 21% in Europe, China's  
1123 usage grew by 1.1%.

1124 For the five years until 2018, electricity generation from renewable wind and solar increased  
1125 by an annual average of 17%, with its global share of electricity generation reaching 7.2% in  
1126 2018. While energy demand for coal, gas, oil and nuclear fell in 2020, renewables demand  
1127 grew by a small amount (0.9%).<sup>173</sup>

1128 Concerningly, global coal demand is expected to rise by 4.5% in 2021, although at the same  
1129 time demand for renewables is set to expand by over 8%.<sup>25</sup> A redirection of efforts towards  
1130 the decarbonisation of the energy system (see panel 4), could put the world on track to meet  
1131 the 1.5°C temperature goal and prevent deaths associated with climate change and air  
1132 pollution.



1133

1134

1135 *Figure 13. The carbon intensity of the energy system for 1970-2018, by 2019 Human Development*  
 1136 *Index country group*

1137

1138 **Indicator 3.2: Clean Household Energy**

1139 *Headline finding: in 2019, only 5% of rural households in countries in the low Human*  
 1140 *Development Index country group relied primarily on clean fuels and technologies for cooking*  
 1141 *(up from just 2% in 2000) – putting them at risk of morbidity and mortality due to exposure to*  
 1142 *household air pollution*

1143 Around 10% of the world’s population, three quarters of whom live in sub-Saharan Africa,  
 1144 lack access to electricity for any service provision, and 2.6 billion people continue to lack  
 1145 access to clean fuel for cooking.<sup>173,174</sup> COVID-19 poses further impediment to achieving the  
 1146 energy access goal (SDG7), with 2020 seeing a 2% rise in lack of access to electricity in sub-  
 1147 Saharan Africa,<sup>175</sup> driving low-income communities in places such as Nairobi, Kenya to  
 1148 increase their usage of wood and kerosene.<sup>176</sup> Energy poverty remains a concern even in high  
 1149 and very high HDI countries – around 7% of people in the EU struggle to afford sufficient heat  
 1150 for their homes,<sup>177</sup> putting them at risk of cold-related adverse health outcomes.<sup>178</sup> Here, and


1151 around the world (as highlighted in panel 3), energy poverty related to excess heat is also an  
1152 important issue.<sup>179</sup>

1153 This indicator tracks energy usage in the home, using data from both the IEA and the  
1154 WHO.<sup>174,180-182</sup> The WHO household energy database compiles data from national surveys,  
1155 collected up to 2017 and projected to 2019, presenting information on fuels and technologies  
1156 used for cooking, heating, and lighting. Using these data, this indicator also presents an  
1157 estimation for household air pollution concentration for 29 countries. A full description of the  
1158 methods, data, and caveats is given in the appendix (pp 116-119).

1159 In the low HDI country group, domestic energy use is dominated by bio-fuels. Primary reliance  
1160 on clean fuels and technologies for cooking in households in the low HDI country group is  
1161 estimated at only 12% in 2019. The share is even lower in rural households of this HDI group,  
1162 with only 5% relying on clean fuels and technologies – a marginal increase from 2% in 2000  
1163 (Figure 14). In homes in the medium and high HDI country groups, the share of solid biofuel  
1164 use has fallen more rapidly, and clean cooking fuel and technology use has risen substantially  
1165 – although in rural areas it remains at 54% for the high HDI group and 39% for the medium  
1166 HDI group.

1167 These patterns of energy use, as well as the infiltration of air from outside, have implications  
1168 on household air pollution concentrations. In rural households in several low and medium  
1169 HDI countries the average PM<sub>2.5</sub> concentration in the main indoor cooking area is estimated  
1170 to be above 500µg m<sup>-3</sup>. In Ethiopia it is over 1200 µg m<sup>-3</sup> – 120 times the WHO threshold of  
1171 10 µg m<sup>-3</sup>.<sup>183</sup> Exposure to these harmful air pollutants in the home results in an estimated 2.31  
1172 million deaths per year.<sup>184</sup>

1173 While gender-differentiated impacts might change across different geographies and  
1174 cultures,<sup>185</sup> exposure to household air pollution is estimated to be around 40% higher for  
1175 women than for men.<sup>186</sup> In many places women are also at higher risk of musculoskeletal  
1176 injuries and violence that result from their domestic role in collecting and using fuels for  
1177 cooking and heating, which further poses risks to their physical and mental wellbeing.<sup>187-190</sup>  
1178 Thus, progress towards meeting the SDG7 would improve health and reduce gender  
1179 inequities.



1181 *Figure 14. Residential energy supply by 2019 Human Development Index country group for 2000 to*  
 1182 *2019. Primary axis: per capita fuel type (coloured bars). Secondary axis: percentage of population*  
 1183 *with primary reliance on clean fuels and technology for cooking. Data taken from the WHO and IEA.*  
 1184 *180-182*  
 1185

1186

1187 **Indicator 3.3: Mortality from Ambient Air Pollution by Sector**


1188 *Headline finding: 3.3 million deaths were attributable to ambient PM<sub>2.5</sub> pollution from human*  
 1189 *sources in 2019 – a third of which were directly related to fossil fuel combustion. The medium*  
 1190 *and high Human Development Index country groups suffered the highest mortality rates*

1191 Awareness of the health impacts of air pollution has increased over the past years, with  
 1192 legislation shifts such as the proposed revision of the EU Ambient Air Quality Directives<sup>191</sup> and  
 1193 a landmark ruling on the death of nine-year-old Ella Adoo-Kissi-Debrah in 2020 in the UK  
 1194 thought to be the first time air pollution was listed as a cause of death in a death certificate.<sup>192</sup>  
 1195 This indicator estimates ambient PM<sub>2.5</sub> exposure and the resulting attributable deaths from

1196 different economic sectors. For the 2021 report, the methods have been updated to use the  
 1197 integrated exposure-response functions (MR-BRTs) used by the 2019 GBD study.<sup>193</sup>

1198 In total, 4.0 million deaths were estimated to be attributable to exposure to ambient PM<sub>2.5</sub> in  
 1199 2019 – 3.3 million of which were from anthropogenic sources and 1.1 million were directly  
 1200 related to fossil fuel combustion. Deaths due to coal combustion have decreased from  
 1201 620,000 in 2015 to 507,000 in 2019, largely due to strict air pollution control measures in  
 1202 China, including the reduction of coal for residential heating.

1203 Ambient concentrations of PM<sub>2.5</sub> differ strongly across world regions and between urban and  
 1204 rural areas. As a result of higher industrial activity, poorer emissions controls, and the  
 1205 continuing use of solid fuels in the domestic sector, countries in medium and high HDI groups  
 1206 face the highest rates of air pollution-related mortality (60 deaths per 100,000 inhabitants,  
 1207 and 65 deaths per 100,000 inhabitants, respectively) (Figure 15). Deaths are lower in both the  
 1208 low and very high HDI country groups (at 34 deaths per 100,000 inhabitants, and 40 deaths  
 1209 per 100,000 inhabitants, respectively). This is due to lower industrial activity and younger  
 1210 populations in low HDI countries; and cleaner electricity generation, industrial production,  
 1211 and end-of-pipe emission controls in very high HDI countries.



1212  
 1213 *Figure 15. Mortality attributable to ambient PM<sub>2.5</sub> in 2015, 2018 and 2019 by source and by 2019*  
 1214 *Human Development Index country group.*

1215

## 1216 Indicator 3.4: Sustainable and Healthy Road Transport

1217 *Headline finding: electricity use in transport rose by 15% from 2017 to 2018 and the global*  
1218 *electric vehicle fleet topped 7.2 million cars in 2019. However, emissions from road transport*  
1219 *also continued to increase*

1220 With road transport accounting for nearly 18% of global CO<sub>2</sub> emissions in 2019, the shift to  
1221 electric vehicles is an important mitigation measure.<sup>194</sup> Beyond this, the promotion of walking  
1222 and cycling (active travel) could not only cut emissions, but also provide enormous health  
1223 dividends through the increase of physical activity.<sup>195</sup> The mode share of cycling varies greatly  
1224 between and within countries of different levels of HDI – ranging from 0.3% and 0.6% of all  
1225 trips in São Paulo and Cape Town, to 1.1-1.9% in USA and Australian cities, to 4.8% in Delhi,  
1226 to 14.1-28.7% in cities in Germany, Japan and the Netherlands – with a higher mode share  
1227 being associated with more equal gender representation in cycling.<sup>196</sup> Unless active travel  
1228 infrastructure is rolled out with consideration of sociocultural inequities, the benefits may not  
1229 be equally manifested across all groups.<sup>197-201</sup>

1230 This indicator uses data from the IEA to monitor fuels used for transport and electric vehicles,  
1231 with full details provided in the appendix (pp 122-123).<sup>202-204</sup> The global number of electric  
1232 vehicles (EVs) rose from 5.1 million in 2018 to 7.2 million in 2019. However, EVs still only  
1233 represent 1% of global car stock, and road transport emissions also increased in 2019, as  
1234 demand for larger vehicles grew in the USA, Europe, and Asia, in tandem with increasing  
1235 demand for transport in low and medium HDI countries. Overall, total direct use of fossil fuels  
1236 for road transport increased by 0.7% whereas the use of electricity in transport rose by 15%  
1237 from 2017 to 2018, although it remains just 0.27% of total road transport energy use.

1238 With respect to the same period in 2019, the COVID-19 pandemic led to a nearly 50%  
1239 decrease in global road transport demand by the end of March 2020.<sup>205,206</sup> However, while  
1240 the use of fossil fuels for road travel has largely rebounded, many public transport networks  
1241 now face critical decreases in ridership.<sup>207</sup> City governments around the world implemented  
1242 measures to promote active travel during their lockdowns, many of which are intended to be  
1243 permanent.<sup>205,206</sup> As cities emerge from the COVID-19 crisis, implementing policies to  
1244 reinforce positive shifts in travel modality presents a triple opportunity to promote physical  
1245 activity, reduce urban air pollution, and mitigate climate change.<sup>208</sup>

1246

## 1247 3.5: Food, Agriculture, and Health

### 1248 Indicator 3.5.1: Emissions from Agricultural Production and Consumption

1249 *Headline finding: mostly driven by high levels of red meat consumption, per capita emissions*  
1250 *from food consumption are considerably greater in of the very high Human Development*




1251 *Index country group – 61% higher than in those in the low Human Development Index group*  
1252 *in 2018*

1253 Food systems, including agricultural production, are responsible for 21-37% of all greenhouse  
1254 gas emissions, while also holding high carbon sequestration potential.<sup>209</sup> This makes them  
1255 key to limiting global warming to 1.5°C. This indicator tracks emissions from agricultural  
1256 production and consumption of food products, combining modelling and FAO data.<sup>210</sup>

1257 Despite moderate improvements in efficiency, total agricultural production emissions  
1258 continued to grow, reaching 5.6 GtCO<sub>2</sub>e in 2018 (1.5% higher than in 2017). Of this total, cattle  
1259 products (mainly meat and milk) contributed 52% of global agricultural production emissions.

1260 Data reveal stark differences in per-capita consumption-based agricultural emissions across  
1261 countries in different levels of HDI: per capita emissions in the very high HDI country group  
1262 are 39% above those of the high HDI group, and 45% higher than those of the low HDI group  
1263 (Figure 16). This is despite a high emission-intensity for beef products in the low HDI group  
1264 (around three times higher than in the very high HDI group), which is mitigated by a much  
1265 lower per capita consumption of beef. Importantly, 68% of the total consumption-based  
1266 agricultural emissions in the very high HDI country group are attributable to cattle products,  
1267 mainly beef production, which is slightly down from 71% in 2000.

1268 Progress towards zero hunger (SDG2) will likely be associated with increases in consumption-  
1269 based agricultural emissions in low and medium HDI countries. In order to meet emission  
1270 reduction goals, consumption of red meat should be safely reduced in relevant population  
1271 groups, especially in very high HDI countries.<sup>211</sup> This would also deliver substantial health co-  
1272 benefits, as indicator 3.5.2 shows. Further scope to reduce emissions from the food  
1273 production system comes from waste reduction, deforestation curtailment and yield  
1274 improvement.<sup>212</sup>



1275

1276

1277 *Figure 16. Per capita yearly greenhouse gas emissions associated with consumption of agri-food*  
 1278 *products, by 2019 Human Development Index country group and commodity, 2000-2018.*

1279

1280 **Indicator 3.5.2: Diet and Health Co-Benefits**

1281 *Headline finding: between 2017 and 2018, estimated deaths due to excess red meat*  
 1282 *consumption rose by 1.8% to 842,000*


1283 With current production efficiency interventions failing to curb or reduce agricultural  
 1284 greenhouse gas emissions, dietary shifts – greatly reducing red meat and increasing plant-  
 1285 based foods – are necessary, particularly in the very high and high HDI countries.<sup>195</sup> For the  
 1286 low and medium HDI countries, sustainable farming and agricultural practices will help keep  
 1287 agricultural emissions low while efforts are made to meet the nutritional requirements of  
 1288 populations.<sup>213</sup> Monitoring this dietary transition, this indicator models deaths attributable  
 1289 to dietary risk factors, using updated data on food consumption and mortality rates by sex,  
 1290 age and country.<sup>214,215</sup>

1291 In 2018, 9.6 million deaths were attributable to imbalanced diets (both dietary composition  
 1292 and caloric intake). Although dietary risks and baseline mortality rates declined, there was an  
 1293 overall increase compared with 2017 (see appendix, pp 130-138). Diets in the high and very

1294 high HDI country groups contain 4 to 7 times more red meat than in the low and medium HDI  
 1295 groups. Together with greater non-communicable disease-related mortality rates, this  
 1296 translates to a rate of red meat-related mortality almost nine times greater in the very high  
 1297 HDI country group (19 deaths per 100,000) compared with the low HDI group (2 deaths per  
 1298 100,000).

1299 Diets and the associated health impacts differ across sexes. In general, male diets tend to be  
 1300 less healthy than those of females, containing fewer fruits (-6% on average globally),  
 1301 vegetables (-1%), and legumes (-10%), and more red meat (+4%).<sup>216-219</sup> The differences in  
 1302 risks resulted in an estimated 455,000 (10%) more men dying from preventable, diet-related  
 1303 diseases than women – a pattern reflected across each of the HDI country groupings (Figure  
 1304 17).

1305



1306  
 1307 *Figure 17. Deaths attributable to imbalanced diets and weight in 2018 by risk factor, sex, and 2019*  
 1308 *Human Development Index country group. The size of each component in the stacked bar represents*  
 1309 *its individual contribution to attributable deaths. Since these contributions cannot be summed*  
 1310 *directly, the overall contribution by diet and weight components are represented by the dots as given*  
 1311 *in the key.*

1312

1313 **Indicator 3.6: Healthcare Sector Emissions**


1314 *Headline finding: in 2018, emissions from the healthcare sector increased slightly to 4.9% of*  
 1315 *global greenhouse gas emissions. Healthcare emissions are positively associated with Human*  
 1316 *Development Index levels, largely through health spending, but minimal association is seen*  
 1317 *after 400 kg CO<sub>2</sub>e per capita*

1318 The healthcare sector is central to improving human development. In providing services,  
1319 healthcare systems mobilise a vast array of products and use energy in various forms, all of  
1320 which result in emissions of greenhouse gases and other pollutants that can be calculated  
1321 throughout global supply chains. With this contribution to greenhouse gas emissions and  
1322 their important leadership role in improving patient care in the face of climate change,<sup>220</sup>  
1323 healthcare institutions are beginning to seriously commit to reducing emissions.<sup>221</sup>

1324 In this indicator, both direct and indirect emissions from the global healthcare sector are  
1325 modelled using environmentally extended multi-region input-output (EE-MRIO) models,  
1326 combined with annual WHO data on national healthcare expenditure, with a full description  
1327 in the appendix (pp 139-140).

1328 In 2018, the global healthcare sector contributed approximately 4.9% of global GHG  
1329 emissions, a rise of 5.2% from 2017. Expansion of healthcare services in China was responsible  
1330 for more than half of this global increase. Although its national healthcare emissions are now  
1331 35% greater than those of the USA, on a per-capita basis, China ranks 21<sup>st</sup> among all major  
1332 economies assessed.

1333 Per-capita comparisons do not account for differences in healthcare access and quality,  
1334 specifically measured through health outcomes, such as life expectancy, which is one of the  
1335 components of the HDI. Plotting per capita healthcare emissions against HDI (Figure 18)  
1336 reveals that emissions are positively associated with HDI, an association strongest for lower  
1337 emissions levels. A wide range of HDI levels are associated with per capita healthcare  
1338 emissions of 500-600 kgCO<sub>2</sub>e, reflecting both differences in health system efficacy and other  
1339 development indicators, but also in emissions intensities. Above these levels, additional  
1340 emissions are not associated with improved HDI.



1341  
 1342 *Figure 18. National per capita healthcare greenhouse gas emissions for 2018 against 2019 country*  
 1343 *Human Development Index level. Dot size is proportional to population*

1344  
 1345 **Conclusion**

1346 Prior to the pandemic, the rapid rate of growth in renewable electricity generation was  
 1347 insufficient to counteract the sluggish decline in coal use. The result of this was that the  
 1348 carbon intensity of the global energy system remained virtually unchanged. At the same time,  
 1349 there has been very little progress in increasing the use of clean household energy. These  
 1350 delays are costing millions of lives each year, from both household and ambient air pollution.  
 1351 Food-related agricultural emissions continue to rise and so too do deaths attributable to  
 1352 dietary risk factors.

1353 Across this section, many inequities can be highlighted. Low HDI countries have the highest  
 1354 use of dirty fuels in the home, putting them at greater risk of morbidity and mortality from  
 1355 exposure to household air pollution. Countries of medium and high levels of HDI have the  
 1356 highest carbon intensity of energy and the greatest burden of deaths due to ambient air  
 1357 pollution, as a result of higher industrial activity and inadequate emissions controls. People  
 1358 in very high HDI countries have the most carbon intensive diets, and, with high levels of red  
 1359 meat consumption, they also have the most to gain from a shift towards more plant-based  
 1360 foods.

1361 Although the impacts of the COVID-19 pandemic are not yet fully captured, there was a  
1362 temporary, but significant drop in emissions due to lockdowns, and the associated reductions  
1363 in economic activities and international travel. However, emissions are already rebounding.  
1364 The challenge moving forward will be to adopt measures that provide near-term economic  
1365 relief, whilst building towards long-term emission reductions and protecting future health –  
1366 a challenge further explored in the following section.

## 1367 Section 4: Economics and Finance

1368 Avoiding the worst of the climate change impacts described in section 1 will require both  
1369 sustained adaptation efforts (section 2), as well as a rapid transformation of the world's  
1370 economies to cut greenhouse gas emissions (section 3). Section 4 examines the economic and  
1371 financial implications of this transition.

1372 First, this section explores the economic impact of climate change and its mitigation  
1373 (indicators 4.1.1 to 4.1.4). The indicators use a range of methods to estimate some of the  
1374 costs that climate change may already be imposing on society through its impacts on human  
1375 health. Then, the economics of the transition to zero-carbon economies (indicators 4.2.1 to  
1376 4.2.5), which are fundamental to the improvement of human health and wellbeing are  
1377 investigated. The indicators consider whether investments and jobs are beginning to move  
1378 away from fossil fuels, and if the appropriate economic signals are encouraging this. A new  
1379 indicator for this year's report (indicator 4.2.5) explores the effect of global trade on  
1380 greenhouse gas and PM<sub>2.5</sub> emissions associated with economic activities, highlighting that  
1381 harms may occur in countries different from the demands that drive them.

1382 Achieving the required investments in the low-carbon transition requires clear and  
1383 committed action from both governments and private sector actors and could result in both  
1384 health and economic benefits. Aiming for a green global recovery from COVID-19 over  
1385 'business as usual' economic growth will ensure that the economy recovers through the  
1386 generation of new jobs in low-carbon industries, as well as accelerate progress towards the  
1387 Paris Agreement goals and the SDGs – yielding health gains through the prevention of further  
1388 climate change and through the co-benefits of climate change mitigation.<sup>222</sup> International  
1389 economic cooperation will be essential to ensure global emission targets are met, and to  
1390 prevent the widening of inequity gaps.<sup>13</sup> This section also therefore reflects on the extent to  
1391 which post-COVID-19 recovery spending has prioritised green investment (panel 4), and the  
1392 alignment of fossil fuel companies' strategies with the requirements of the transition (panel  
1393 5).

1394

### 1395 4.1 The Economic Impact of Climate Change and its Mitigation

#### 1396 Indicator 4.1.1: Economic Losses due to Climate-Related Extreme Events

1397 *Headline finding: when normalised by GDP, economic losses from climate-related extreme*  
1398 *events in 2020 were collectively three times greater in with the medium Human Development*  
1399 *Index country group compared with the very high Human Development Index group*

1400 The loss of physical infrastructure and resulting economic losses due to climate-related  
1401 extreme events can further exacerbate the health impacts described in section 1. This  
1402 indicator tracks the total annual economic losses (insured and uninsured) that result from  
1403 climate-related extreme events, using data provided by Swiss Re, with methods described in  
1404 the appendix (pp 141-143).<sup>223,224</sup>

1405 In 2020 there were 242 recorded climate-related extreme events, with absolute economic  
1406 losses totalling US\$178 billion. Although two-thirds of these losses occurred in very high HDI  
1407 economies, when normalised by GDP, losses the medium HDI country groups are around  
1408 three times greater. Importantly, while two-thirds of losses in the very high HDI country group  
1409 are insured, almost 93% of losses were uninsured in the high HDI group. This number rises to  
1410 97% and 100% of measurable losses in the medium and low HDI country groups, respectively  
1411 – creating a bigger economic burden for these disadvantaged countries, as uninsured losses  
1412 are either not replaced, or are replaced through out-of-pocket expenses, reinforcing  
1413 inequities.

1414


1415 [Indicator 4.1.2: Costs of Heat-Related Mortality](#)

1416 *Headline finding: the monetised value of global heat-related mortality increased by 6.7% from*  
1417 *0.27% of gross world product in 2018, to 0.28% in 2019. Europe continued to be the worst*  
1418 *affected region, facing costs equivalent to the average income of 6.1 million of its citizens*

1419 The increase in morbidity and mortality due to extremes of heat represents a high cost to all  
1420 of society. This indicator uses data on years of life lost due to extremes of heat from indicator  
1421 1.1.6 to provide a measure of the costs of global deaths attributable to heat.<sup>74</sup> Improved in  
1422 the 2021 report, it combines a value of statistical life-year (VSLY) with years of life lost (YLLs),  
1423 to monetise the loss caused by premature mortality. The valuation of life across varying HDI  
1424 levels presents a methodological and ethical challenge, which this indicator addresses by  
1425 presenting costs as the proportion of GDP and the equivalent annual average income.

1426 The monetised value of global heat-related mortality in the 65-and-over population increased  
1427 by 6.7%, from 0.27% of gross world product in 2018 to 0.28% in 2019 (Figure 19). Reflecting  
1428 the distribution of impacts found in indicator 1.1.6, the costs of heat-related mortality for the  
1429 low, medium, high, and very high HDI country groups, were found to be equivalent to the  
1430 average income of 0.94, 4.80, 8.20, and 7.52 million of their citizens, respectively. As in  
1431 indicator 1.1.6, the WHO's European region was the worst affected in 2019, with costs equal  
1432 to the average income of 6.1 million of its citizens and 0.66% of regional GDP. However, the  
1433 costs were lower than the year before, due to fewer estimated heat-related deaths in 2019  
1434 compared to 2018 in this region (indicator 1.1.6). On the other hand, costs increased in other  
1435 regions, especially the WHO's South-East Asia region.





1436  
 1437 *Figure 19. Monetised cost of heat-related (in terms of expressed as the number of people whose*  
 1438 *average income the loss is equivalent to) by 2019 Human Development Index country group for 2000-*  
 1439 *2019*

1440


1441 **Indicator 4.1.3: Loss of Earnings from Heat-Related Labour Capacity Reduction**

1442 *Headline finding: working in conditions of extreme heat is a health risk. Such conditions could*  
 1443 *reduce the capacity for paid labour, with an impact on workers' earnings equivalent to 4-8%*  
 1444 *of GDP in the low Human Development Index country group in 2020*

1445 As reflected in indicator 1.1.4, higher temperatures, driven by climate change, are affecting  
 1446 people's ability to work. This indicator considers the loss of earnings that could result from  
 1447 such reduced capacity, Such earnings losses could further compound the health impacts  
 1448 through effects on the socioeconomic determinants of good health.<sup>225</sup> It combines the  
 1449 outputs of indicator 1.1.4 with data on average earnings by country and sector held in the  
 1450 International Labour Organization (ILO) databases, with methods and additional analysis  
 1451 described in the appendix (pp 146-151).<sup>226</sup> In this year's report, the number of countries  
 1452 covered in this indicator has been increased from 25 to 183.

1453 Indicators 1.1.6 and 4.1.2 found Europe to be the region most affected by heat-related  
 1454 mortality in populations aged 65 and over. In contrast, this indicator focusses on working age  
 1455 populations and, in alignment with the outputs of indicator 1.1.4, finds that greater loss of  
 1456 earnings due to labour capacity loss occur in low and medium HDI countries. Countries with  
 1457 lower HDI levels tend to experience greater proportional losses of earnings, emphasising the  
 1458 impact of climate change on deepening inequities. In the low HDI country group, potential

1459 income losses in 2020 were equivalent to 4-8% of GDP, depending on the degree of shade or  
 1460 sun exposure during agricultural and construction work (Figure 20). The ranges for the  
 1461 medium, high, and very high HDI country groups in 2020 were 2-4%, 1-2% and 0.3-0.5% of  
 1462 GDP, respectively. The impacts will mainly affect men in sectors such as construction, where  
 1463 they represent more than 90% of the workforce globally, and in manufacturing and  
 1464 agriculture where, where they represent more than 60% of the workforce.<sup>63</sup> However, the  
 1465 data does not account for informal or unpaid domestic and agricultural work, in which women  
 1466 are often overrepresented.<sup>227-229</sup> The indirect economic impacts from reduced labour capacity  
 1467 extend well beyond the loss of earnings. For example, modelling both direct and indirect  
 1468 impacts, the heat-related economic cost of labour loss in 2020 was estimated at 1.36% of  
 1469 China's GDP and 6.75% of GDP in Hainan Province.<sup>30</sup>



1470 *Figure 20 Average potential loss of earnings in the low Human Development Index country group as a*  
 1471 *result of potential labour loss due to heat exposure. Losses are presented as share of GDP, by sector*  
 1472 *of employment,. The agriculture and construction (sun additional) blocks represent the losses that*  
 1473 *would have been incurred in addition to those from agriculture and construction (shade) if all of the*  
 1474 *activities in these sectors had been carried out in direct sunlight.*  
 1475

1476


1477 **Indicator 4.1.4: Costs of the Health Impacts of Air Pollution**

1478 *Headline finding: equivalent to the annual income of 71.1 million and 99.1 million people, the*  
 1479 *greatest economic costs of mortality due to air pollution fall on countries in the medium and*  
 1480 *high Human Development Index country groups. Costs relative to GDP decreased between*  
 1481 *2015 and 2019 globally, with the exception of costs in South-East Asia*

1482 As described in indicator 3.3, global mortality due to ambient PM<sub>2.5</sub> pollution has increased.  
 1483 This indicator captures the cost of this mortality by placing an economic value on the YLLs  
 1484 that result from exposure to anthropogenic ambient PM<sub>2.5</sub>. This indicator has been expanded  
 1485 for the 2021 report, from a European-only focus to global coverage, and with a revised  
 1486 definition of YLLs. The methods, data and further analysis are described in full in the appendix  
 1487 (pp 152-154).

1488 Figure 21 presents the economic value of YLLs in 2015 and 2019 by country HDI groups,  
 1489 relative to both total GDP and the annual income of the average person in these categories.  
 1490 The greatest relative costs fall on the medium and high HDI country groups, equivalent to the  
 1491 annual income of 74.6 million and 99.1 million people, respectively. Costs relative to average  
 1492 income increased between 2015 and 2019 in the low and medium HDI country groups.  
 1493 However, with rates of growth of GDP outpacing those of population, costs relative to total  
 1494 GDP have decreased in all HDI groups.

1495



1496

1497 Figure 21. Economic cost of YLLs in 2015 and 2019, relative to the annual income of the average  
 1498 person and total GDP, by 2019 Human Development Index country group

1499 4.2 The Economics of the Transition to Zero-Carbon Economies

1500

---

1501 *Panel 4. Recovering from Covid-19: Stimulus Measures for a Sustainable Economy*

1502 The COVID-19 pandemic, and measures to tackle it, triggered a global recession of a depth only exceeded in the  
1503 last 150 years by two World Wars and the Great Depression of the 1930s.<sup>230</sup> Governments with the fiscal capacity  
1504 have responded with massive spending packages; by the end of 2020, the world's 50 largest economies had  
1505 committed USD 14.6 trillion in fiscal measures (many times higher than the value of global stimulus measures  
1506 following the 2008-09 financial crisis). Although 87% of this was designed to prevent an even deeper health and  
1507 economic crisis (USD 12.7 trillion), rather than encourage recovery (USD 1.9 trillion),<sup>231</sup> as time goes by and  
1508 further measures are announced, promoting recovery will come to the fore.

1509 How these measures are designed and targeted will determine whether this spending entrenches existing  
1510 technical, economic, and social structures and systems, or promotes those that are more sustainable, healthy,  
1511 and equitable. Evidence from stimulus measures introduced following the 2008-09 financial crisis shows that  
1512 'green' stimulus measures often have advantages over 'brown' or 'colourless' measures.<sup>222</sup>

1513 So far, the signs are not encouraging. Of the USD 1.9 trillion directed toward recovery by the end of 2020, just  
1514 18% is expected to reduce greenhouse gas emissions (or 2.5% of the value of all fiscal measures), while the  
1515 overall impact on air pollution, and particularly on natural capital – through the expansion of road transport and  
1516 defence services in particular – is likely to be negative. Positive measures are highly concentrated in just a few  
1517 nations, particularly in Europe,<sup>231</sup> although measures announced so far in 2021 indicate some movement  
1518 towards greater consideration of sustainability in other countries.<sup>232,233</sup> However, despite global CO<sub>2</sub> emissions  
1519 dropping by a record 6% in 2020 overall, they have rebounded quickly, with global CO<sub>2</sub> emissions in December  
1520 2020 around 2% higher than in December 2019.<sup>234</sup> The urgency with which the trillions of dollars for stimulus  
1521 measures yet to be announced must be oriented toward a green and healthy recovery is therefore great.

1522 In May 2020, the WHO published six prescriptions for a healthy and green recovery: (1) Protect and preserve  
1523 the source of human health: Nature; (2) Invest in essential services, from water and sanitation to clean energy  
1524 and healthcare facilities; (3) Ensure a quick, healthy energy transition; (4) Promote healthy, sustainable food  
1525 systems; (5) Build healthy, liveable cities; and (6) Stop using taxpayers money to fund pollution (particularly  
1526 through fossil fuel subsidies).<sup>235</sup> If governments are serious about their commitments under the Paris Agreement  
1527 and SDGs, they must take note of these priorities, plan ahead, and learn from both their own previous  
1528 experience and from that generated elsewhere, to implement them using well-designed and context-  
1529 appropriate policy. Where necessary, multilateral institutions, processes and instruments should be galvanised  
1530 in support of a global recovery that is both sustainable and equitable.<sup>231</sup>

---

1531


1532 Indicator 4.2.1: Coal and Clean Energy Investment

1533 *Headline finding: global investment in energy supply and energy efficiency reduced 13%*  
1534 *between 2019 and 2020. Investment in renewable energy and energy efficiency increased by*  
1535 *3%, but investment in new coal capacity reduced by 13%*

1536 Coal combustion has been responsible for over 30% of the global average temperature  
1537 increase above pre-industrial levels and for 491,000 deaths from PM<sub>2.5</sub> exposure in 2019  
1538 (indicator 3.3).<sup>236</sup> Therefore, coal phase-out is essential for both mitigating climate change  
1539 and for reducing premature mortality due to air pollution. At the same time, it is necessary to  
1540 invest in renewables, energy efficiency, and the electricity grid in order to reduce the carbon  
1541 intensity of energy supply, as described in indicator 3.1. Taking data from the IEA, this  
1542 indicator tracks global investment in energy supply and energy efficiency, and highlights  
1543 ongoing capital spending in new coal-fired power generation, globally and for key countries  
1544 and regions. The data, presented as an index, represents ongoing capital spending.

1545 Between 2019 and 2020 investment in global energy supply and energy efficiency reduced  
1546 from nearly \$2 trillion to around \$1.7 trillion, almost entirely due to declining investment in  
1547 fossil fuels, following reduced demand as a result of the pandemic (investment in coal power  
1548 capacity declined by 13%). In parallel, investment in renewables and energy efficiency  
1549 increased by 3%, with their share of total investment in global energy supply increasing from  
1550 33% to 39%. However, for a pathway consistent with 1.5°C of warming this century, annual  
1551 investments in clean energy must at least triple over the 2020s.<sup>237</sup>

1552



1553

1554 *Figure 22. Economic value of annual investment in renewable and fossil fuel energy supply and*  
1555 *energy efficiency, 2014-2020*

1556

1557 [Indicator 4.2.2: Employment in Low-Carbon and High-Carbon Industries](#)

1558 *Headline finding: direct employment in fossil fuel extraction declined by 14% from 13.1 million*  
1559 *in 2019 to 12.7 million in 2020*

1560 Evidence suggests that employees in some fossil fuel extraction industries, particularly coal  
1561 mining, and their local communities, suffer a greater incidence of cardiovascular and  
1562 cerebrovascular disease, respiratory disease and cancers.<sup>238</sup> Investments in renewable  
1563 energies and energy efficiency are estimated to create almost three times more jobs per unit  
1564 of spend than those in fossil fuel industries.<sup>239</sup> Along with strong labour and environmental  
1565 standards, investment and employment in renewables present an opportunity to improve  
1566 health and livelihoods. This indicator tracks global direct employment in fossil fuel extraction  
1567 industries and direct and indirect (supply chain) employment in renewable energy, with a full  
1568 description available in the appendix (pp 158-159).

1569 Around 11.5 million people globally were employed directly or indirectly by the renewable  
1570 energy industry in 2019, representing an increase of 4.2% from 2018. At the time of writing  
1571 data for 2020 was unavailable, although due to the pandemic, the extent to which such data  
1572 will be indicative of a long-term trend is currently unclear. Fossil fuel extraction industries  
1573 continue to employ more people globally than all renewable energy industries combined,

1574 although the number of jobs in 2019 are slightly lower than in 2018, at 12.7 million compared  
1575 with 13.1 million.

1576 While men are still overrepresented in the energy sector, the field of renewable energy  
1577 employs a considerably higher share of women (32%) than the oil and gas industry (22%).<sup>240</sup>  
1578 With adequate policies in place, the transition to a low carbon economy therefore represents  
1579 an additional opportunity to reduce gender inequities and empower women.

1580 With trillions of dollars earmarked for COVID-19 recovery, investments in the renewable fuel  
1581 industry could offer a triple gain in terms of better health through safer jobs and improved  
1582 livelihoods, climate change mitigation, and more employment opportunities.

1583

1584 [Indicator 4.2.3: Funds Divested from Fossil Fuels](#)

1585 *Headline finding: the global value of funds committing to fossil fuel divestment between 2008*  
1586 *and 2020 is US\$14.52 trillion, with health institutions accounting for US\$42 billion*

1587 By reducing financial interests in the fossil fuel industry, divestment both reduces the ‘social  
1588 licence to operate’ of fossil fuel companies, and hedges against investors’ risk of losses due  
1589 to ‘stranded assets’ in an increasingly decarbonising world (panel 5).<sup>241,242</sup> Investors can also  
1590 effect change through shareholder action, exemplified recently by activist hedge fund Engine  
1591 No 1 taking seats on ExxonMobil’s board.<sup>243</sup> Concerned with the immediate and long-term  
1592 damages of continued fossil fuel use, health institutions have the imperative to lead the way  
1593 in divesting, to ensure they ‘first, do no harm’. This indicator tracks the total global value of  
1594 funds divested from fossil fuels, and the value of funds divested by health institutions, using  
1595 data provided by 350.org.<sup>244</sup>

1596 From 2008 until the end of 2020, 1,398 organisations, with assets worth at least US\$14.52  
1597 trillion, have committed to divestment. Of these, only 25 are health institutions, with assets  
1598 totalling US\$42 billion. The value of new funds committed to divesting in 2020 was US\$2.5  
1599 trillion, with health institutions accounting for US\$47 million of these.

1600

1601

1602 *Panel 5. Compatibility of fossil fuel company strategies with well below 2°C-consistent emissions trajectories*

---

1603 Globally, carbon dioxide (CO<sub>2</sub>) from the combustion of fossil fuels represents 65% of total greenhouse gas  
1604 emissions.<sup>168</sup> In the 2015 Paris Agreement, countries agreed to reduce their emissions to keep global warming  
1605 to ‘well below 2°C’ with respect to pre-industrial levels. The carbon budget for a 66% probability of limiting  
1606 global warming to 1.5°C has been estimated at 420 GtCO<sub>2</sub>.<sup>14</sup> However, the potential CO<sub>2</sub> emissions from reserves  
1607 held by the 200 largest public fossil fuel companies is at least 1,541 GtCO<sub>2</sub>,<sup>245</sup> whilst the carbon contained in  
1608 global resources of fossil fuels is estimated at about 11,000 GtCO<sub>2</sub>,<sup>246</sup> well beyond the maximum that can be  
1609 used if the world is to meet the Paris Agreement goals. A third of oil reserves, half of gas reserves and over 80  
1610 per cent of coal reserves worldwide should remain unused to keep global warming below 2°C,<sup>246</sup> representing  
1611 stranded assets and unburnable carbon.<sup>233,247</sup> Future energy system scenarios with strict carbon constraints, low  
1612 fossil fuel demand, high capital costs projects and carbon-intensive reserves increase the risk of stranding  
1613 assets,<sup>248</sup> with considerable financial consequences for their owners and industry stakeholders.<sup>249</sup>

1614 Although the fossil fuel industry has begun to acknowledge that the energy system is transitioning away from  
1615 unabated oil, gas and coal, countries’ fossil fuel production plans to 2030 could exceed levels consistent with  
1616 limiting warming to 2°C by 50%, and by 120% in relation to 1.5°C.<sup>250</sup> Companies are following diverging business  
1617 strategies,<sup>251</sup> with most of them falling short of what is required to mitigate transition risks. While an increasing  
1618 number of oil and gas companies are announcing net-zero commitments, for these to be consistent with climate  
1619 ambitions they must be framed on the basis of their total emissions rather than on their emission intensities,  
1620 cover scope 1, 2 and 3 emissions, and account for activities based on a company’s full equity share.<sup>62,252</sup> Those  
1621 companies who better understand systemic risks, stress-test potential scenarios, and develop business  
1622 strategies with interim targets and investments that align adequately with well below 2°C targets (and preferably  
1623 1.5°C) are likely to become more resilient over the coming years, as climate-risk scrutiny from investors and  
1624 financial regulators increases.

---

1625

1626 [Indicator 4.2.4: Net Value of Fossil Fuel Subsidies and Carbon Prices](#)

1627 *Headline finding: 77% of the 84 countries reviewed had a net-negative carbon price in 2018.*  
1628 *The resulting net loss of revenue was in many cases equivalent to substantial proportions of*  
1629 *the national health budget*

1630 Placing a carbon price on fossil fuel use helps to reflect more accurately its negative  
1631 externalities, including its impact on health, and to encourage the transition away from fossil  
1632 fuels. However, not all countries set carbon prices, and where they are imposed, they can be  
1633 undermined by subsidies provided for fossil fuels.


1634 This indicator compares carbon prices and fossil fuel subsidies to calculate ‘net’ economy-  
1635 wide average carbon prices and revenues. It covers 84 countries, which are responsible for  
1636 around 92% of global CO<sub>2</sub> emissions. The indicator is based on data from the IEA,<sup>253</sup> OECD,<sup>254</sup>



1637 the World Bank,<sup>255</sup> and the WHO, with methods and further analysis in the appendix (pp 162-  
1638 165).<sup>256</sup>

1639 In 2018, 65 out of the 84 countries analysed (77%) had net-negative carbon prices, reflecting  
1640 an overall subsidising of fossil fuels. The median value of the subsidy in these countries was  
1641 US\$1 billion, with some countries providing net subsidies to fossil fuels in the tens of billions  
1642 of dollars each year. 42 countries had a carbon pricing mechanism in place, but only 19  
1643 succeeded in discouraging fossil fuels with net-positive carbon prices – all of which were very  
1644 high HDI countries. Nonetheless, most very high HDI countries still had net-negative carbon  
1645 prices (Figure 23). These net subsidies are equivalent to substantial proportions of national  
1646 health spending in many countries.

1647 With low-income populations vulnerable to energy costs, removing subsidies can be a  
1648 challenge, but redirecting spending from fossil subsidy to healthcare and health-related  
1649 services would most likely deliver net benefits to their wellbeing.<sup>257</sup> Furthermore,  
1650 international financing mechanisms to support low-income countries in their transition to  
1651 sustainable energy sources are essential to safeguard all dimensions of human health.<sup>258</sup>



1652 *Figure 23. Net carbon prices (left), net carbon revenues (centre), and net carbon revenue as a share*  
1653 *of current national health expenditure (right), across 84 countries in 2018, arranged by 2019 Human*  
1654 *Development Index country group: low (n=1), medium (n=7), high (n=23) and very high (n=53). Boxes*  
1655 *show the interquartile range (IQR), horizontal lines inside the boxes show the medians, and the*  
1656 *brackets represent the full range from minimum to maximum.*  
1657

1658

1659 [Indicator 4.2.5: Production- and Consumption-Based Attribution of CO<sub>2</sub> and PM<sub>2.5</sub> Emissions](#)

1660 *Headline finding: in 2019, 18% of CO<sub>2</sub> and 17% of PM<sub>2.5</sub> global emissions were embodied in*  
1661 *trades between countries of different Human Development Index levels*


1662 The production of goods and services often drives both greenhouse gas and PM<sub>2.5</sub> emissions,  
1663 thus contributing to impacts on health and wellbeing. Emissions from local production  
1664 ('production-based emissions') occur within the geographical territories of nations through  
1665 the local production of goods and services. An alternative way of accounting for the burden

1666 of pollution is to assign the emissions to the country which is the final consumer of the  
1667 products that are made – known as ‘consumption-based emissions’. A comparison of  
1668 production- and consumption-based emissions gives a better understanding of how  
1669 emissions are embodied in global trade, which is essential to enable better international  
1670 policy formulation that protects human health in all geographies.

1671 This indicator captures the pollution burden from a country’s local production, as well as that  
1672 driven by a nation’s domestic final consumption, including the burden embedded in its  
1673 imports. It uses an EE-MRIO model and the EXIOBASE database, to estimate CO<sub>2</sub>  
1674 emissions,<sup>259,260</sup> and the GAINS model to produce a PM<sub>2.5</sub> emission inventory.<sup>261</sup> More details  
1675 on the methodology, and further analysis, can be found in the appendix (pp 166-172).

1676 In 2019, 18% of CO<sub>2</sub> (of 35.6 Gt world total) and 17% of PM<sub>2.5</sub> (of 37.4 Mt world total) global  
1677 emissions were embodied in trades among countries of different HDI levels. The largest  
1678 contributors to global consumption-based CO<sub>2</sub> and PM<sub>2.5</sub> emissions were China (28 % and  
1679 18%), the USA (17% and 5%), the EU (10% and 6%), and India (7 % and 16%). The USA did the  
1680 most ‘outsourcing’ of emissions, with 21% CO<sub>2</sub> (of 5.9 Gt total) and 49% PM<sub>2.5</sub> (of 1.7 Mt total)  
1681 emissions resulting from the production of goods it consumed, actually occurring in other  
1682 countries. In contrast, 16% of CO<sub>2</sub> (of 10.8 Gt total) and 13% of PM<sub>2.5</sub> (of 6.8 Mt total)  
1683 emissions that occurred in China resulted from the local production of goods that were  
1684 ultimately exported to consumers in other countries.

1685 The very high HDI country group contributed the most production-based (45%) and  
1686 consumption-based (49%) CO<sub>2</sub> emissions in 2019. However, the high HDI country group was  
1687 the biggest contributor to both production-based (38%) and consumption-based (35%) PM<sub>2.5</sub>  
1688 emissions (Figure 24), with the very high HDI country group the lowest emitter of PM<sub>2.5</sub>, partly  
1689 as a result of stricter local air pollution regulations. Importantly, the very high HDI country  
1690 group was the only group with higher consumption-based emissions than production-based  
1691 emissions, i.e. a net ‘outsourcing’ in terms of their consumption-related emissions.



1692  
 1693 *Figure 24. The flows of embodied CO<sub>2</sub> and PM<sub>2.5</sub> emissions among different Human Development Index*  
 1694 *country groups in 2019*

1695  
 1696 **Conclusion**

1697 The impacts of climate change on health are already having significant economic  
 1698 consequences and fall in different ways across countries of all levels of HDI. The economic  
 1699 losses of climate-related extreme events are three times higher in medium HDI countries than  
 1700 in very high HDI countries. However, the monetised value of global heat-related deaths is  
 1701 highest in Europe, and the greatest costs of premature mortality due to air pollution fall in  
 1702 countries with medium and high HDI levels. South-East Asia was the only region with  
 1703 increasing air pollution mortality costs between 2015 and 2019, relative to GDP. Extreme heat  
 1704 can create economic impacts by reducing labour capacity. In this case, those employed in low-  
 1705 wage, outdoor work in low HDI countries are likely to be most affected.

1706 Because of the potentially large and unequally distributed impacts of climate change on  
1707 human health, incomes and wellbeing, substantial and sustained investment in the low  
1708 carbon transition is required. Overall, global investments in coal power continue to decline,  
1709 although with worrying counter-trends in certain countries. Investments in renewables and  
1710 energy efficiency continue to grow, as do divestments from fossil fuel assets, however a  
1711 considerable increase in the pace of change is required.

1712 Both governments and the private sector have crucial roles to play in bringing about the  
1713 required transition. Governments across all HDI groups must address fossil fuels subsidies in  
1714 countries. Although withdrawing energy subsidies is challenging when it affects people on  
1715 low incomes, other forms of government spending, including on health services, can provide  
1716 better and more targeted support to decrease inequities and maximise wellbeing. The global  
1717 trade system means that almost a fifth of CO<sub>2</sub> and PM<sub>2.5</sub> emissions occur in the production of  
1718 goods that are subsequently traded between countries of different HDI levels. This underlines  
1719 the importance of inclusive global agreements that facilitate cooperation on policies for the  
1720 reduction of both production and consumption emissions.

1721 As governments begin to invest in recovery from COVID-19, there is a crucial window of  
1722 opportunity to reduce fossil fuel subsidies, invest more in clean energy, and support a green  
1723 recovery. Policies and regulations must be developed that subject fossil fuel companies to  
1724 greater scrutiny and ensure their alignment with a world well below 2°C.

## 1725 Section 5: Public and Political Engagement

1726 As the preceding sections make clear, climate change is damaging people's health and  
1727 widening the fault lines of inequality, with the human costs amplified by COVID-19.<sup>28,262,263</sup>  
1728 Those least responsible for climate change are most exposed to impacts that are 'hitting  
1729 harder and sooner' than climate assessments indicated even a decade ago.<sup>264</sup> Action at the  
1730 speed and scale needed to meet the ambitions of the Paris Agreement requires public and  
1731 political engagement, particularly in industrialised countries where 'the major part of  
1732 emissions originate'.<sup>265</sup> This section tracks engagement in health and climate change in the  
1733 media as well as by individuals, scientists, governments and the corporate sector.

1734 The mainstream media is a major platform for public engagement. It remains the most widely-  
1735 used source of information,<sup>266</sup> shaping public perceptions<sup>267-269</sup> and influencing the social  
1736 media agenda.<sup>270</sup> Indicator 5.1 tracks coverage of health and climate change in 67 newspapers  
1737 from 37 countries, including the *People's Daily (Renmin Ribao)*, China's longest-running  
1738 national newspaper and the official outlet of government.<sup>271,272</sup> The indicator also includes a  
1739 content analysis of coverage in India and the USA, focusing on 'prestige' newspapers with  
1740 influence on the countries' political and economic elites.<sup>273-275</sup>

1741 Individual engagement (indicator 5.2) is tracked through individuals' searches on Wikipedia,  
1742 the online information source with wider reach and coverage than traditional  
1743 encyclopaedias.<sup>276-278</sup> The third indicator (Indicator 5.3) tracks engagement in peer-reviewed  
1744 journals, the primary source of scientific evidence for the media, government, and the  
1745 public.<sup>279</sup>

1746 Government engagement (indicator 5.4) is tracked by statements made by national leaders  
1747 at the UN General Assembly, the policy-making body of the UN. The annual meeting opens  
1748 with the General Debate where heads of government, or their high-ranking representatives,  
1749 address the global community on issues they consider important.<sup>280,281</sup> Indicator 5.4 also  
1750 considers engagement with health in the enhanced NDCs, submitted in compliance with the  
1751 2015 Paris Agreement.<sup>282-284</sup> Panel 6 compares health engagement in the initial and enhanced  
1752 set of NDCs held on the UNFCCC NDC registry on 1 April 2021.

1753 Action by the corporate sector will be decisive in moving societies away from dependence on  
1754 fossil fuels.<sup>285-287</sup> Indicator 5.5 tracks engagement in health and climate change by companies  
1755 within the UN Global Compact, the world's biggest corporate sustainability initiative.<sup>288,289</sup>  
1756 Companies commit to shared principles of sustainable behaviour and submit annual reports  
1757 on progress.

1758 With increasing acknowledgement of the need to recognise and investigate gender inequities  
1759 in the representation, communication, and governance of climate change,<sup>290-293</sup> engagement  
1760 with gender is incorporated where appropriate. Engagement with health, climate change and

1761 COVID-19, and analyses by WHO region and HDI country group are also included. Details of  
1762 data sources and methods for all indicators are provided in the appendix, along with  
1763 additional analyses.

1764

## 1765 [Indicator 5.1 Media Coverage of Health and Climate Change](#)

1766 *Headline finding: in 2020, the upward trend in coverage of health and climate change*  
1767 *continued, but failed to match the increase seen in 2019. In 2020, most of the coverage of*  
1768 *health and climate change referred to COVID-19*

1769 Newspapers provide an important forum for public engagement. They shape public  
1770 understanding of climate change, both through their influence on their readers and on the  
1771 wider political agenda.<sup>268,294</sup> This indicator tracks coverage of health and climate change from  
1772 2007, the year before the WHO World Health Assembly made a multilateral commitment to  
1773 protect people’s health from climate change.<sup>295</sup> The indicator includes 66 newspapers  
1774 spanning 36 countries and four languages, together with an additional analysis of China’s  
1775 People’s Daily. The indicator also examines the content of 2020 coverage in newspapers in  
1776 India and the USA. Methods and further analysis are provided in the appendix (pp 172-195)

1777 Across the 36 countries, the upward trend in newspaper coverage of health and climate  
1778 change continued, reaching 11,371 articles in 2020. However, the rate of increase was lower  
1779 than that of 2019 – 6% from 2019 to 2020, compared with 96% from 2018 to 2019. As in 2019,  
1780 coverage was greatest in the WHO America and Europe regions and lowest in the African  
1781 region.

1782 Engagement with gender and with COVID-19 was examined in English language newspapers  
1783 across 23 countries. While the proportion of articles referring to gender increased between  
1784 2007 and 2020 (from 97 (2%) of 6,044 articles to 573 (6%) of 10,092), gender remains marginal  
1785 to the representation of health and climate change in the mainstream press. In 2020, over  
1786 60% (6,238) of the 11,371 articles referring to health and climate change also referred to  
1787 COVID-19; in April and May 2020, it was over 80%.

1788 In China’s *People’s Daily*, the limited coverage of health and climate change noted in earlier  
1789 *Lancet Countdown* reports was again evident in 2020. Of the 1,106 articles discussing climate  
1790 change, 2% were related to human health. Across the 2008-2020 period, no articles related  
1791 to health and climate change engaged with gender issues. In 2020, no articles discussed the  
1792 relationships between climate change and COVID-19, or how they together influenced health.

1793 Analysis of the content of coverage of health and climate change focuses on India (medium  
1794 HDI) and the USA (very high HDI). The selected newspapers, the *Times of India* and *Hindustan*  
1795 *Times* along with the *New York Times* and *Washington Post*, form part of the ‘prestige’ press,

1796 seen to exercise influence on political and economic elites and on the wider policy  
1797 agenda.<sup>275,296</sup>

1798 One set of themes related to the health impacts of climate-related hazards, including  
1799 heatwaves and wildfires. For example, the *New York Times* (18 June), noted that “people with  
1800 health issues, older people and young children are especially susceptible to the effects of  
1801 extreme heat [and...] it’s a threat that grows as climate change continues”.<sup>297</sup> Another set of  
1802 themes related to the spread of infectious disease, including COVID-19. For example, the  
1803 *Hindustan Times* (25 February) reported that “climate change may revert back successes of  
1804 controlling infectious diseases” with “consensus among scientists that there has been a rise  
1805 in zoonotic diseases - Nipah, Ebola, Zika, Corona viruses - in recent decades ... driven by  
1806 biodiversity loss and climate change”.<sup>298</sup> As this last comment indicates, climate change and  
1807 environmental change are often linked together; scientific reports (including the *Lancet*  
1808 Countdown reports) are cited as evidence that “we are close to running out of time —  
1809 approaching a point of no return for human health, which depends on planetary health” (*New*  
1810 *York Times*, 28 April).<sup>299</sup>

1811

## 1812 [Indicator 5.2: Individual Engagement in Health and Climate Change](#)


1813 *Headline finding: individual information-seeking about health and climate change decreased*  
1814 *overall by 15% from 2019 to 2020; spikes in engagement in mid-2020 were almost exclusively*  
1815 *due to interest in pandemic-related content*

1816 Individual engagement in climate change and health is tracked through the digital footprint  
1817 of users of the online encyclopaedia, Wikipedia. Wikipedia has outpaced traditional  
1818 encyclopaedias in terms of reach, coverage, and comprehensiveness and is one of the most-  
1819 visited websites worldwide.<sup>276,300,301</sup> The analysis is based on the English-language Wikipedia  
1820 which represents around 50% of global traffic to all Wikipedia language editions.<sup>302,303</sup>

1821 The indicator focuses on ‘clickstream’ activity, where an individual clicks between an article  
1822 on health and climate change (or vice versa). Because clickstream activity captures only pairs  
1823 of sequential visits, for the 2021 report, the set of articles was extended to include a wider  
1824 range of health and climate change articles. In 2020, as in previous years, individuals seldom  
1825 moved between health and climate change; instead, co-click activity was predominantly  
1826 within the set of articles on health or climate change.

1827 Figure 25 tracks co-click activity from 2018 to 2020, looking separately at the volume  
1828 generated by clicks on a climate-related link in a health-related page, vice versa, and the sum  
1829 of both. Overall numbers are very low, confirming that engagement in either climate change  
1830 or health rarely triggers engagement in the other topic. Further, the volume of health-climate  
1831 co-views fell in 2020 by 15%, reversing the upward trend evident in 2019. When co-clicks to

1832 an article relating to COVID-19 are excluded, the downward trend in 2020 becomes even  
 1833 more pronounced. The spike in co-clicks in mid-2020 was almost exclusively due to interest  
 1834 in pandemic-related content, which then sparked interest in climate change, whereas the rise  
 1835 over September/October was generated by an initial interest in climate change.



1836

1837 *Figure 25. Aggregate monthly co-clicks on Wikipedia articles related to human health and climate*  
 1838 *change, 2018—2020. Blue: co-click from health-related page to climate-related page. Orange:*  
 1839 *co-click from climate-related page to health-related page. Grey: sum of all health and climate co-click*  
 1840 *activity.*

1841

1842 [Indicator 5.3: Coverage of Health and Climate Change in Scientific Journals](#)

1843 *Headline finding: original research on health and climate change increased eleven-fold*  
 1844 *between 2007 and 2020, driven primarily by scientists in countries of the highest Human*  
 1845 *Development Index levels. Gender remained marginal to research on health and climate*  
 1846 *change across the period. In 2020, 7% of health and climate change articles referred to COVID-*  
 1847 *19*

1848 Scientific evidence is a key resource for the media, individuals and governments, and is  
 1849 playing a critical role shaping public and political engagement in health and climate  
 1850 change.<sup>278,304</sup> The indicator is based on searches in OVID Medline and OVID Embase, using  
 1851 references to health and climate change in article titles and abstracts, with methods and  
 1852 further analyses provided in the appendix (pp 218-231).


1853 The upward trend in scientific engagement in health and climate change noted in previous  
 1854 *Lancet Countdown* reports has been maintained, with the number of articles on health and  
 1855 climate change increasing by 28% between 2019 and 2020, to reach its highest recorded level



1856 of 858 articles. The trend is driven by the rapid increase in original research (primary studies  
1857 and systematic reviews), which increased by 32% between 2019 and 2020. Research-related  
1858 articles (e.g. evidence reviews, editorials, letters) also increased, but at a lower rate.

1859 Increasing scientific engagement in health and climate change is driven by very high HDI  
1860 countries (Figure 26); 76% of the total output in 2020 was led by researchers in this group. In  
1861 contrast, scientists in low HDI countries were lead authors of just 1% of journal articles.

1862



1863

1864 *Figure 26. Scientific journal articles relating to health and climate change, 2007-2020, by 2019*  
1865 *country Human Development Index country group*

1866

1867 In 2007/08, under 2% of health and climate change articles engaged with gender in some way;  
1868 in 2020, the proportion was 6%. Similarly, 2020, only 7% of the articles on health and climate  
1869 change addressed COVID-19, suggesting this rise in scientific research in health and climate  
1870 change is independent of the concurrent global health crisis. Articles engaging with gender  
1871 and with COVID-19 were predominantly led by scientists in the very high HDI countries.

1872


1873 Indicator 5.4: Government Engagement in Health and Climate Change

1874 *Headline finding: in 2020, 47% of government leaders engaged with the health dimensions of*  
1875 *climate change in their statements at the UN General Debate, more than double the*  
1876 *proportion in 2019. The increase was linked to engagement with the COVID-19 pandemic*

1877 Government leadership, backed by strong near-term policies, is required if the increase in  
1878 global temperature is to be halted.<sup>168</sup> This indicator examines government engagement with  
1879 health and climate change in the UN General Debate (UNGD). Engagement with health in  
1880 commitments to emissions reduction made by governments under the 2015 Paris Agreement  
1881 is also considered in panel 6.

1882 The UNGD opens each new session of the UN General Assembly. It provides all UN member  
1883 states with an opportunity to address the global community on priorities for action. Among  
1884 many global challenges, including economic recession and social conflict, the indicator  
1885 captures whether government leaders draw attention to health and climate change. Analysis  
1886 is based on the application of a key word search in the United Nations General Debate corpus  
1887 using natural language processing,<sup>305,306</sup> with 8,288 statements analysed across 1970-2020.

1888 Figure 27 tracks the proportion of countries referring to health and climate change in their  
1889 UNGD statements between 1970 and 2020. In 2020, the proportion of countries engaging  
1890 with the health dimensions of climate change reached the highest recorded level, increasing  
1891 from 22% (43 countries) in 2019 to 47% (91 countries) in 2020. Additionally, and for the first  
1892 time in the UNGD, every member state referred to health in their 2020 address – a reflection  
1893 of the ongoing global pandemic.



1894  
1895  
1896

*Figure 27. Proportion of countries referring to climate change, health, and the intersection between the two in their UNGA statements, 1970-2020*

1897  
1898  
1899  
1900  
1901  
1902

Increased engagement in health and climate change is linked to discussion of the COVID-19 pandemic, represented by government leaders as both a threat and an opportunity. The pandemic highlights “the vulnerabilities of our societies [to]...global disasters...lurking just around the corner... [like] climate change” (Austria). It also presents an opportunity to tackle the climate crisis: “our recovery from this pandemic must mark a transition to a decarbonized, climate-resilient economic system” (Fiji).

1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911

Engagement in health and climate change continues to be led by countries in the low HDI group and, in particular, by the SIDS.<sup>283,284</sup> For the SIDS, COVID-19 has amplified the risks of climate change: “our unique circumstances and consequent vulnerabilities have left us exposed to the ravages of the twin crises of the pandemic and climate change” (St Lucia). In 2020, 75% of the SIDS discussed health and climate change in the 2020 UNGA. However, 2020 also saw greater engagement among higher-income countries. A key issue is whether this pandemic-related increase in engagement among richer countries will be maintained in future years.

1912

=====

1913

*Panel 6: The place of health in the enhanced NDCs*

1914 The 2015 Paris Agreement is the only global framework for reducing greenhouse gas emissions to protect  
1915 people’s health.<sup>307</sup> Countries committed to emissions reductions via Nationally Determined Contributions  
1916 (NDCs), to be enhanced every five years. In 2015/16, 185 countries, including an EU submission for 27 countries,  
1917 submitted initial NDCs. By July 2021, 87 countries, including an EU submission for 26 countries, had submitted  
1918 enhanced or new NDCs.<sup>308</sup>

1919 Compared with their initial NDCs, the proportion of countries referring to health increased, from 56% (49) to  
1920 91% (79). However, health engagement remained low. Overall, in both initial and enhanced NDCs, under 3% of  
1921 the text related to health; in the enhanced NDCs, this represented an average of 240 of 10466 words. Of the  
1922 references to health, 30% (249 references) noted health impacts, challenges or risks; for example, “the Kenyan  
1923 economy is dependent on climate-sensitive sectors, such as rain-fed agriculture, water, energy, tourism, wildlife,  
1924 and health, whose vulnerability is increased by climate change” (Kenya, updated submission). A further 25%  
1925 (210) related to health sector adaptation; for example, climate change “threatens the ability of health  
1926 institutions and organizations to maintain and improve health services into the future” (Marshall Islands, second  
1927 submission).

1928 The enhanced NDCs demonstrate an increased engagement with gender, health, and climate change with 9  
1929 (10%) NDCs making a meaningful connection compared with just 2 (2%) in their initial contributions. The  
1930 majority of these are references to the specific impact of climate change on women; for example, “further strain  
1931 on the workload of women and climate change related stress during pregnancy could contribute to low birth  
1932 weight, leading to increases in risks of undernutrition and non-communicable diseases” (Cambodia, updated  
1933 submission).

1934 In summary, while health engagement remains low, there is greater recognition that climate change takes a  
1935 disproportionate toll on women.

---

---

1936 Indicator 5.5: Corporate Sector Engagement in Health and Climate change

1937 *Headline finding: in 2020, engagement in health and climate change increased to its highest*  
1938 *level among companies in the UN Global Compact. Over a third (38%) of companies referred*  
1939 *to the health dimensions of climate change in their 2020 progress reports*

1940 The indicator tracks engagement in health and climate change among companies signed up  
1941 to the UN Global Compact, established to promote corporate social and environmental  
1942 responsibility,<sup>288</sup> although its effectiveness has been critiqued, with the suggestion that  
1943 membership could be a form of ‘greenwashing’ and ‘bluewashing’ for some companies.<sup>309</sup>  
1944 The Compact represents over 12,000 companies from 160 countries, with each submitting an  
1945 annual Communication on Progress (GCCOP) against a set of social and environmental  
1946 principles.

1947 The indicator is based on the application of a key word search in the text corpus of 17,984  
1948 GCCOP reports submitted in English between 2011 and 2020.<sup>288</sup> In the 2019 and 2020 *Lancet*  
1949 *Countdown* reports, the focus was on the healthcare sector. This report considers corporate  
1950 engagement across all sectors.

1951 Figure 28 tracks engagement in health and climate change in annual GCCOP reports from  
1952 2011 to 2020. As it indicates, the large majority of reports refer to health (84% of 2029 reports  
1953 in 2020) and climate change (75%) as separate topics. In contrast, only a minority made  
1954 reference to the health dimensions of climate change (38% in 2020). However, it represents  
1955 a large increase from 2014, the low point of engagement, when only 21% of corporations  
1956 made reference to the intersection between climate change and health. Three sectors stand  
1957 out for their high levels of engagement in health and climate change: food and drug retailers,  
1958 oil and gas producers, and alternative energy. In 2020, over 70% of corporations in these  
1959 sectors made reference to health and climate change; in the healthcare sector, the proportion  
1960 was only 37%.

1961 Additional analyses examined references to gender in the GCCOP reports engaging with  
1962 health and climate change. Only a minority additionally referred to gender. However, the  
1963 proportion increased from 2014 to 2019, with a particularly sharp rise (to 19%) in the 2019  
1964 report. In 2020, gender engagement fell to 13% (see appendix pp 249-264).




Figure 28. Proportion of companies referring to climate change, health, and the intersection of health and climate change in their UN Global Compact Communication on Progress (GCCOP) reports, 2011-2020.

1965  
1966  
1967  
1968

1969

1970 **Conclusion**

1971 Public and political engagement is essential if the ambitions of the Paris Agreement are to be  
1972 realised.<sup>168</sup> Section 5 has focused on five areas of engagement: the media, the public, the  
1973 scientific community, national government and the corporate sector. Three conclusions can  
1974 be drawn.

1975 Firstly, health and climate change are increasingly addressed together. The trend is  
1976 particularly pronounced for indicators relating to the media, science, government and the  
1977 corporate sector. In all these areas, engagement with health and climate change reached its  
1978 highest recorded level in 2020. Gender is rarely integrated into engagement within the  
1979 health-climate change nexus, although there is increased recognition in countries' enhanced  
1980 NDCs.

1981 Secondly, the COVID-19 pandemic appears to be a major driver of engagement in 2020. For  
1982 example, over half of newspaper coverage of health and climate change was linked to COVID-  
1983 19 and individual engagement in health and climate change was largely sustained by searches  
1984 for articles related to COVID-19. Government engagement in the health dimensions of climate

1985 change was similarly underpinned by engagement in the pandemic. It remains to be seen if  
1986 the heightened engagement in health and climate change will be maintained if and when the  
1987 pandemic-related crises are contained.

1988 Thirdly, social inequities remain deeply etched into public and political engagement. In the  
1989 media and in science, coverage of health and climate change engagement is greatest in the  
1990 very high HDI countries, the group exerting the greatest pressure on the planet but relatively  
1991 protected from the health impacts of climate change. Meanwhile, medium and low HDI  
1992 countries have much smaller carbon and environmental footprints – yet, are shouldering the  
1993 immediate burden of climate change, and are far less represented in the scientific literature.  
1994 As in previous years, the SIDS are leading global engagement with the health impacts of  
1995 climate change at the UN General Debate. What is required is for their leadership to be  
1996 matched by a decisive break with ‘business as usual’ by countries and communities  
1997 contributing most to climate change.

## 1998 Conclusion: the 2021 Report of the *Lancet* Countdown

1999 The 2021 report of the *Lancet* Countdown finds a world overwhelmed by an ongoing global  
2000 health crisis, while it has made little progress to protect its population from the  
2001 simultaneously aggravated health impacts of climate change. The inequities of these impacts  
2002 and the response, including those of gender, are brought into sharp focus within each of the  
2003 indicators presented. This exposes the urgent need for collection of standardised data to  
2004 capture inequities and vulnerabilities (panel 2).

2005 Climate-sensitive infectious diseases are of increasing global concern and the environmental  
2006 suitability for the transmission of all infectious diseases tracked is rising (indicator 1.3.1). For  
2007 non-cholerae *Vibrio* bacteria, the environmental suitability for transmission in northern  
2008 latitudes increased by 56% since the 1980s. The number of months suitable for malaria  
2009 transmission has increased by 39% in highland areas of the low HDI country group and, over  
2010 the past 5 years, the environmental suitability for the transmission of emerging arboviruses  
2011 – dengue, chikungunya and Zika – was between 7% and 13% higher than in the 1950s.

2012 The high temperatures in 2020, a year that tied with 2016 as the hottest year on record,  
2013 resulted in extreme heat-related health impacts, affecting the emotional and physical  
2014 wellbeing of populations around the world (indicators 1.1.1-1.1.6). These higher  
2015 temperatures and altered weather patterns are also leading to more frequent extreme  
2016 weather events and increased wildfire exposure (indicators 1.2.1, 1.2.2 and 1.2.3), and are  
2017 putting years of progress on food and water security at risk in many parts of the world. The  
2018 five years with the greatest area of the world's surface affected by droughts have all occurred  
2019 since 2015 (indicator 1.2.2), the yield potential of all major staple crops continues to fall as a  
2020 result of the rising temperatures (indicator 1.4.1), and 79% of all potential work hours lost to  
2021 extreme heat in low HDI countries occurred in the agricultural sector in 2020 (indicator 1.1.4).

2022 However, measures to curb emissions have been grossly inadequate. Emissions are declining  
2023 too slowly or heading in the wrong direction in the highest emitting sectors (indicators 3.1,  
2024 3.4 and 3.5.1). This delay in progress is contributing to millions of deaths each year due to  
2025 exposure to indoor and ambient PM<sub>2.5</sub> pollution, and due to high-carbon, unhealthy diets  
2026 (indicators 3.2, 3.3 and 3.5.2). Importantly, these impacts manifest differently between HDI  
2027 country groups and genders, underscoring profound inequities.

2028 Despite years of scientific reporting on climate change impacts, efforts to build resilience have  
2029 been slow and unequal, with countries of low levels of Human Development Index the least  
2030 prepared to respond to the changing health profile of climate change, and funding remaining  
2031 a consistent challenge (indicators 2.1.1, 2.3.1, and 2.4). At the same time, 65 out of 84  
2032 countries reviewed continue to provide subsidies for fossil fuels that outweigh any revenue



2033 received from carbon pricing instruments. The resulting ‘net carbon subsidy’ is in many cases  
2034 equivalent to substantial proportions of countries’ national health budgets (indicator 4.2.4).

2035 Governments with the fiscal capacity have responded to the COVID-19 pandemic with  
2036 massive spending packages, to cushion the impacts of the crisis and start to bring about  
2037 economic recovery. But as the world approaches COP26, the response to climate change, and  
2038 commensurate investment, remains inadequate. The opportunity for the green recovery is in  
2039 danger of being missed. A fossil-fuel driven recovery, whilst potentially meeting narrow and  
2040 near-term economic targets, could push the world irrevocably off course for the ambitions of  
2041 the Paris Agreement, with enormous costs to human health.

2042 With government leaders more engaged with the health dimensions of climate change than  
2043 ever before (indicator 5.4), countries across the globe must pursue low carbon economic  
2044 recovery pathways, implementing policies that reduce inequities and improve human health.  
2045 The *Lancet* Countdown indicators show the evidence to support the urgency and opportunity  
2046 of this transition, and that none of us is safe until everyone is safe.

2047

## 2048 Contributors

2049 The Lancet Countdown and the work for this paper was conducted by five working groups,  
2050 which were responsible for the design, drafting, and review of their individual indicators and  
2051 sections. All authors contributed to the overall paper structure and concepts, and provided  
2052 input and expertise to the relevant sections.

2053 ER, CDN, NA, SA-K, JC, LC, SD, LEE, SHG, IK, TK, DK, BL, JKWL, YL, ZL, RL, JM-U, CM, KMi, MM-  
2054 L, KAM, NO, MO, FO, MRa, JCS, LS, MT, JTr, BV, and MY contributed to Working Group 1. KLE,  
2055 MN, LJ, DC-L, RD, LG, DG, CH, JH, MPJ, PLK, MM, KMo, TN, MOS, JR, and JS-G contributed to  
2056 Working Group 2. TO, IH, HK, KB, CD, MD, PD-S, ME, SH, S-CH, GK, ML, NM, JM, DP, JS, MS,  
2057 JTa, PW, and MW contributed to Working Group 3. PE, PD, NH, BSR, WC, KH, ZM, FW, and SZ  
2058 contributed to Working Group 4. HG, PL, WC, SC, ND, SJ, LM, SM, and OP contributed to  
2059 Working Group 5. AC, HM, PG, IH, MRo, AM and RS provided coordination, strategic direction,  
2060 and editorial support.

## 2061 Declaration of interest

2062 We declare no competing interests.

## 2063 Acknowledgements

2064 We thank the Wellcome Trust, in particular Madeleine Thomson and Lukasz Aleksandrowicz,  
2065 for financial and strategic support, without which this research collaboration would not be  
2066 possible.

2067 The Lancet Countdown's work is supported by an unrestricted grant from the Wellcome Trust  
2068 (grant number 209734/Z/17/Z). 10 of the authors (IH, MRo, AM, CDN, LJ, HK, PD, NH, BSR, PL)  
2069 were compensated for their time while working on the drafting and development of the  
2070 Lancet Countdown's report.

2071 The work of CH is supported by a NERC fellowship (grant number NE/R01440X/1) and funding  
2072 from the Wellcome Trust HEROIC project (grant number 216035/Z/19/Z). The work of CD was  
2073 supported by the UK Natural Environment Research Council Independent Research  
2074 Fellowship (grant number NE/N01524X/1) and contributes to the Sustainable and Healthy  
2075 Food Systems (SHEFS) project supported by the Wellcome Trust (grant number  
2076 205200/Z/16/Z). The work of MD was supported by the Wellcome Trust's Complex Urban  
2077 Systems for Sustainability and Health (CUSSH) project (grant number 209387/Z/17/Z). The  
2078 work of TO was supported by the Engineering and Physical Sciences Research Council Centre  
2079 for Research in Energy Demand Solutions (grant number EP/R035288/1). The work of YL, LS,  
2080 and BV was supported by the NASA Applied Sciences Program, managed by J. Haynes (grant  
2081 number 80NSSC21K0507). RL was supported by a Royal Society Dorothy Hodgkin Fellowship.  
2082 The work of JR and MOS was supported by the Swedish Research Council Formas (grant  
2083 number 2018–01754 and 2017–01300). The work of MW was supported by the UK Energy  
2084 Research Centre research programme which is funded by the UK Research and Innovation  
2085 Energy Programme (grant number EP/S029575/1). The work of SHG and JKWL was supported  
2086 by the National Research Foundation, Prime Minister's Office, Singapore under its Campus  
2087 for Research Excellence and Technological Enterprise (CREATE) programme, and a research  
2088 grant from the NUS Initiative to Improve Health in Asia (NIHA) coordinated by the Global Asia  
2089 Institute of the National University of Singapore and supported by the Glaxo Smith Kline-  
2090 Economic Development Board (Singapore) Trust Fund. Any opinions, findings and conclusions  
2091 or recommendations expressed in this material are those of the authors and do not reflect  
2092 the views of the National University of Singapore, Singapore or the National Research  
2093 Foundation, Singapore.

2094 While carrying out its work, The Lancet Countdown received invaluable technical advice and  
2095 input from several individuals, including Heather Adair-Rohani, Miguel Gomez-Escolar Viejo  
2096 and Jessica Lewis (World Health Organization, Geneva, Switzerland), Ginette Azcona, Antra  
2097 Bhatt, and Sara Duerto Valero (UN Women), Simon Bennett, Chiara Delmastro, Ryszard  
2098 Pospiech, and Michael Waldron (International Energy Agency, Paris, France), Peter James and  
2099 Catherine Ngo (Harvard University, Boston, MA, USA), Sebastian Ramirez Ruiz (Hertie School,  
2100 Berlin, Germany), Nicholas Goh (National University of Singapore, Singapore), Kaixin Huang  
2101 (Northeastern University, Boston, MA, USA), Kai Chen and Amy Darefsky (Yale University, New

2102 Haven, CT, USA). Policy and communications advice was given by Emma-Louise Frost, Frances  
2103 MacGuire, and Kim van Daalen (the Lancet Countdown).  
2104

## 2105 References

- 2106 1. Johns Hopkins Center for Systems Science and Engineering. COVID-19 Dashboard. 2021.  
2107 <https://www.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b48e9ecf6>  
2108 (accessed 7 April 2021).
- 2109 2. Office for National Statistics. Comparisons of all-cause mortality between European countries  
2110 and regions: 2020. 2021.  
2111 [https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/article](https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/article/s/comparisonsofallcausemortalitybetweeneuropeancountriesandregions/2020)  
2112 [s/comparisonsofallcausemortalitybetweeneuropeancountriesandregions/2020](https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/article/s/comparisonsofallcausemortalitybetweeneuropeancountriesandregions/2020) (accessed 9 April  
2113 2021).
- 2114 3. WHO. Pulse survey on continuity of essential health services during the COVID-19 pandemic:  
2115 interim report. Geneva, Switzerland: World Health Organization, 2020.
- 2116 4. International Monetary Fund. World Economic Outlook Update January 2021. Washington,  
2117 DC, USA: International Monetary Fund, 2021.
- 2118 5. World Bank. Poverty and Shared Prosperity 2020: Reversals of Fortune. Washington, DC, USA:  
2119 World Bank, 2020.
- 2120 6. Stiglitz J, Rashid H. Averting Catastrophic Debt Crises in Developing Countries. Extraordinary  
2121 challenges call for extraordinary measures: Centre for Economic Policy and Research, 2020.
- 2122 7. McCarthy M, Christidis N, Stott P, Kaye N. Met Office: A review of the UK's climate in 2020.  
2123 2021. <https://www.carbonbrief.org/met-office-a-review-of-the-uks-climate-in-2020> (accessed 8 April  
2124 2021).
- 2125 8. Stuart-Smith RF, Roe GH, Li S, Allen MR. Increased outburst flood hazard from Lake Palcacocha  
2126 due to human-induced glacier retreat. *Nature Geoscience* 2021; **14**(2): 85-90.
- 2127 9. van Oldenborgh GJ, Krieken F, Lewis S, et al. Attribution of the Australian bushfire risk to  
2128 anthropogenic climate change. *Nat Hazards Earth Syst Sci Discuss* 2020; **2020**: 1-46.
- 2129 10. Philip SY, Kew SF, van Oldenborgh GJ, et al. Rapid attribution analysis of the extraordinary  
2130 heatwave on the Pacific Coast of the US and Canada June 2021, 2021.
- 2131 11. Ciavarella A, Cotterill D, Stott P, et al. Siberian heatwave of 2020 almost impossible without  
2132 climate change. 2020. [https://www.worldweatherattribution.org/siberian-heatwave-of-2020-almost-](https://www.worldweatherattribution.org/siberian-heatwave-of-2020-almost-impossible-without-climate-change/)  
2133 [impossible-without-climate-change/](https://www.worldweatherattribution.org/siberian-heatwave-of-2020-almost-impossible-without-climate-change/) (accessed 8 April 2021).
- 2134 12. Walton D, van Aalst M. Climate-related extreme weather events and COVID-19. A first look at  
2135 the number of people affected by intersecting disasters. Geneva, Switzerland: International  
2136 Federation of Red Cross and Red Crescent Societies, 2020.
- 2137 13. Dibley A, Wetzler T, Hepburn C. National COVID debts: climate change imperils countries'  
2138 ability to repay. *Nature* 2021; **592**(7853): 184-7.
- 2139 14. IPCC. Global warming of 1.5°C. An IPCC Special Report on the impacts of global warming of  
2140 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context  
2141 of strengthening the global response to the threat of climate change. Geneva, Switzerland: World  
2142 Meteorological Organization, 2018.
- 2143 15. Levy BS, Patz JA. Climate Change, Human Rights, and Social Justice. *Ann Glob Health* 2015;  
2144 **81**(3): 310-22.
- 2145 16. GISTEMP Team. GISS Surface Temperature Analysis (GISTEMP), version 4. 2021.  
2146 <https://data.giss.nasa.gov/gistemp/> (accessed 7 April 2021).
- 2147 17. Lenssen NJL, Schmidt GA, Hansen JE, et al. Improvements in the GISTEMP Uncertainty Model.  
2148 *Journal of Geophysical Research: Atmospheres* 2019; **124**(12): 6307-26.
- 2149 18. NOAA. More Near-Record Warm Years Are Likely On Horizon. 2021.  
2150 <https://www.ncei.noaa.gov/news/projected-ranks> (accessed May 11, 2021).

- 2151 19. Met Office. Mauna Loa carbon dioxide forecast for 2021. 2021.  
 2152 [https://www.metoffice.gov.uk/research/climate/seasonal-to-decadal/long-range/forecasts/co2-](https://www.metoffice.gov.uk/research/climate/seasonal-to-decadal/long-range/forecasts/co2-forecast-for-2021)  
 2153 [forecast-for-2021](https://www.metoffice.gov.uk/research/climate/seasonal-to-decadal/long-range/forecasts/co2-forecast-for-2021) (accessed 7 April 2021).
- 2154 20. Green JK, Seneviratne SI, Berg AM, et al. Large influence of soil moisture on long-term  
 2155 terrestrial carbon uptake. *Nature* 2019; **565**(7740): 476-9.
- 2156 21. Lenton TM, Rockström J, Gaffney O, et al. Climate tipping points—too risky to bet against.  
 2157 Nature Publishing Group; 2019.
- 2158 22. Wunderling N, Donges JF, Kurths J, Winkelmann R. Interacting tipping elements increase risk  
 2159 of climate domino effects under global warming. *Earth Syst Dynam* 2021; **12**(2): 601-19.
- 2160 23. Le Quéré C, Jackson RB, Jones MW, et al. Temporary reduction in daily global CO2 emissions  
 2161 during the COVID-19 forced confinement. *Nature Climate Change* 2020.
- 2162 24. Friedlingstein P, O'Sullivan M, Jones MW, et al. Global carbon budget 2020. *Earth System*  
 2163 *Science Data* 2020; **12**(4): 3269-340.
- 2164 25. IEA. Global Energy Review 2021. Paris, France: IEA, 2021.
- 2165 26. Climate Action Tracker. Global Update: Climate Summit Momentum. 2021.  
 2166 <https://climateactiontracker.org/publications/global-update-climate-summit-momentum/> (accessed  
 2167 7 May 2021).
- 2168 27. Watts N, Adger WN, Ayeb-Karlsson S, et al. The Lancet Countdown: tracking progress on  
 2169 health and climate change. *The Lancet* 2017; **389**(10074): 1151-64.
- 2170 28. UNDP. Human Development Report 2020 - The Next Frontier: Human Development and the  
 2171 Anthropocene. 2020. <https://report.hdr.undp.org/> (accessed March 21 2021).
- 2172 29. Beggs PJ, Zhang Y, McGushin A. The 2021 report of the MJA-Lancet Countdown on health and  
 2173 climate change: Australia increasingly out on a limb. *Medical Journal of Australia* 2021 **In Press**.
- 2174 30. Cai W, Zhang C, Zhang S. The 2021 China Report of the Lancet Countdown on Health and  
 2175 Climate Change: Seizing the window of opportunity. *The Lancet Public Health* 2021; **In Press**.
- 2176 31. Allen C, West R, Beagley J, McGushin A. Climate Change and Health in Small Island Developing  
 2177 States. London, UK The Lancet Countdown: Tracking Progress on Health and Climate Change,  
 2178 2021.
- 2179 32. IPCC. Climate Change 2014. Impacts, Adaptation, and Vulnerability. Working Group II  
 2180 Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.  
 2181 Cambridge and New York, 2014.
- 2182 33. Watts N, Amann M, Arnell N, et al. The 2020 report of The *Lancet* Countdown on health and  
 2183 climate change: responding to converging crises. *The Lancet* 2021; **397**(10269): 129-70.
- 2184 34. Hayward G, Ayeb-Karlsson S. 'Seeing with Empty Eyes': a systems approach to understand  
 2185 climate change and mental health in Bangladesh. *Climatic Change* 2021; **165**(1): 29.
- 2186 35. Kelman I, Ayeb-Karlsson S, Rose-Clarke K, et al. A review of mental health and wellbeing under  
 2187 climate change in small island developing states (SIDS). *Environmental Research Letters* 2021; **16**(3):  
 2188 033007.
- 2189 36. Position Statement PS03/21: Our planet's climate and ecological emergency: Royal College of  
 2190 Psychiatrists, 2021.
- 2191 37. Gopalkrishnan N. Cultural diversity and mental health: Considerations for policy and practice.  
 2192 *Frontiers in public health* 2018; **6**: 179.
- 2193 38. Li M, Gu S, Bi P, Yang J, Liu Q. Heat Waves and Morbidity: Current Knowledge and Further  
 2194 Direction-A Comprehensive Literature Review. *International Journal of Environmental Research and*  
 2195 *Public Health* 2015; **12**(5): 5256-83.
- 2196 39. Kovats RS, Hajat S. Heat Stress and Public Health: A Critical Review. *Annu Rev Public Health*  
 2197 2008; **29**(1): 41-55.

- 2198 40. Basu R. Relation between Elevated Ambient Temperature and Mortality: A Review of the  
2199 Epidemiologic Evidence. *Epidemiologic Reviews* 2002; **24**(2): 190-202.
- 2200 41. Schmeltz M, Petkova E, Gamble J. Economic Burden of Hospitalizations for Heat-Related  
2201 Illnesses in the United States, 2001–2010. *International Journal of Environmental Research and Public  
2202 Health* 2016; **13**(9): 894.
- 2203 42. Hansen A, Bi L, Saniotis A, Nitschke M. Vulnerability to extreme heat and climate change: is  
2204 ethnicity a factor? *Global Health Action* 2013; **6**(1): 21364.
- 2205 43. Chambers J. Global and cross-country analysis of exposure of vulnerable populations to  
2206 heatwaves from 1980 to 2018. *Climatic Change* 2020; **163**(1): 539-58.
- 2207 44. Bassil KL, Cole DC. Effectiveness of public health interventions in reducing morbidity and  
2208 mortality during heat episodes: a structured review. *International Journal of Environmental Research  
2209 and Public Health* 2010; **7**(3): 991-1001.
- 2210 45. WHO. Heat and Health. 2018. [https://www.who.int/news-room/fact-sheets/detail/climate-  
2211 change-heat-and-health](https://www.who.int/news-room/fact-sheets/detail/climate-change-heat-and-health) (accessed 17 April 2021).
- 2212 46. WMO Catalogue of Major Meteorological Hazards. Available at  
2213 <http://puslitbang.bmkg.go.id/litbang/wmo-catalogue-of-major-meteorological-hazards/>. 2021.
- 2214 47. de Perez EC, van Aalst M, Bischiniotis K, et al. Global predictability of temperature extremes.  
2215 *Environmental Research Letters* 2018; **13**(5).
- 2216 48. Di Napoli C, Pappenberger F, Cloke HL. Verification of Heat Stress Thresholds for a Health-  
2217 Based Heat-Wave Definition. *Journal of Applied Meteorology and Climatology* 2019; **58**(6): 1177-94.
- 2218 49. Arem H, Moore SC, Patel A, et al. Leisure Time Physical Activity and Mortality: A Detailed  
2219 Pooled Analysis of the Dose-Response Relationship. *JAMA Internal Medicine* 2015; **175**(6): 959-67.
- 2220 50. Warburton DER, Nicol CW, Bredin SSD. Health benefits of physical activity: the evidence.  
2221 *Canadian Medical Association Journal* 2006; **174**(6): 801-9.
- 2222 51. Nuzum H, Stickel A, Corona M, Zeller M, Melrose RJ, Wilkins SS. Potential Benefits of Physical  
2223 Activity in MCI and Dementia. *Behavioural Neurology* 2020; **2020**: 7807856.
- 2224 52. Peluso MAM, Andrade LHS. Physical activity and mental health: the association between  
2225 exercise and mood. *Clinics* 2005; **60**: 61-70.
- 2226 53. Zhang X, Li X, Sun Z, et al. Physical activity and COVID-19: an observational and Mendelian  
2227 randomisation study. *Journal of Global Health* 2020; **10**(2).
- 2228 54. An R, Shen J, Li Y, Bandaru S. Projecting the Influence of Global Warming on Physical Activity  
2229 Patterns: a Systematic Review. *Current Obesity Reports* 2020; **9**(4): 550-61.
- 2230 55. Heaney AK, Carrión D, Burkart K, Lesk C, Jack D. Climate Change and Physical Activity:  
2231 Estimated Impacts of Ambient Temperatures on Bikeshare Usage in New York City. *Environmental  
2232 Health Perspectives* 2019; **127**(3): 037002.
- 2233 56. Nazarian N, Liu S, Kohler M, et al. Project Coolbit: can your watch predict heat stress and  
2234 thermal comfort sensation? *Environmental Research Letters* 2021; **16**(3): 034031.
- 2235 57. Andrews O, Le Quéré C, Kjellstrom T, Lemke B, Haines A. Implications for workability and  
2236 survivability in populations exposed to extreme heat under climate change: a modelling study. *The  
2237 Lancet Planetary Health* 2018; **2**(12): e540-e7.
- 2238 58. Armstrong LE, Casa DJ, Millard-Stafford M, Moran DS, Pyne SW, Roberts WO. Exertional Heat  
2239 Illness during Training and Competition. *Medicine & Science in Sports & Exercise* 2007; **39**(3): 556-72.
- 2240 59. Casa DJ, DeMartini JK, Bergeron MF, et al. National Athletic Trainers' Association Position  
2241 Statement: Exertional Heat Illnesses. *Journal of Athletic Training* 2015; **50**(9): 986-1000.
- 2242 60. Flouris AD, Dinas PC, Ioannou LG, et al. Workers' health and productivity under occupational  
2243 heat strain: a systematic review and meta-analysis. *The Lancet Planetary Health* 2018; **2**(12): e521-  
2244 e31.

- 2245 61. International Labour Organization. ILO Monitor: COVID-19 and the world of work. Seventh  
2246 edition. Geneva, Switzerland: International Labour Organization, 2021.
- 2247 62. ILO. Working on a warmer planet: The impact of heat stress on labour productivity and decent  
2248 work. Geneva, Switzerland: International Labour Organization, 2019.
- 2249 63. International Labour Organization. Employment by sex and economic activity — ILO modelled  
2250 estimates. 2020.
- 2251 64. International Labour Organization. Indigenous peoples and climate change: from victims to  
2252 change agents through decent work Geneva: ILO, 2017.
- 2253 65. Obradovich N, Migliorini R, Paulus MP, Rahwan I. Empirical evidence of mental health risks  
2254 posed by climate change. *Proc Natl Acad Sci U S A* 2018; **115**(43): 10953-8.
- 2255 66. Mullins JT, White C. Temperature and mental health: Evidence from the spectrum of mental  
2256 health outcomes. *J Health Econ* 2019; **68**: 102240.
- 2257 67. Burke M, González F, Baylis P, et al. Higher temperatures increase suicide rates in the United  
2258 States and Mexico. *Nature Climate Change* 2018; **8**(8): 723-9.
- 2259 68. Carleton TA. Crop-damaging temperatures increase suicide rates in India. *Proceedings of the  
2260 National Academy of Sciences* 2017; **114**(33): 8746-51.
- 2261 69. Baylis P. Temperature and temperament: Evidence from Twitter. *Journal of Public Economics*  
2262 2020; **184**: 104161.
- 2263 70. Baylis P, Obradovich N, Kryvasheyev Y, et al. Weather impacts expressed sentiment. *PLoS One*  
2264 2018; **13**(4): e0195750.
- 2265 71. Carleton TA, Hsiang SM. Social and economic impacts of climate. *Science* 2016; **353**(6304):  
2266 aad9837.
- 2267 72. Pennebaker JW, Boyd RL, Jordan K, Blackburn K. The development and psychometric  
2268 properties of LIWC2015, 2015.
- 2269 73. Song X, Wang S, Hu Y, et al. Impact of ambient temperature on morbidity and mortality: An  
2270 overview of reviews. *Science of The Total Environment* 2017; **586**: 241-54.
- 2271 74. Honda Y, Kondo M, McGregor G, et al. Heat-related mortality risk model for climate change  
2272 impact projection. *Environ Health Prev Med* 2014; **19**(1): 56-63.
- 2273 75. Guo Y, Gasparrini A, Armstrong BG, et al. Temperature Variability and Mortality: A Multi-  
2274 Country Study. *Environ Health Perspect* 2016; **124**(10): 1554-9.
- 2275 76. Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204 countries  
2276 and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The  
2277 Lancet* 2020; **396**(10258): 1204-22.
- 2278 77. Abatzoglou JT, Williams AP, Barbero R. Global Emergence of Anthropogenic Climate Change  
2279 in Fire Weather Indices. *Geophysical Research Letters* 2019; **46**(1): 326-36.
- 2280 78. NASA EarthData. Active Fire Data. Available at [https://earthdata.nasa.gov/earth-observation-  
data/near-real-time/firms/active-fire-data](https://earthdata.nasa.gov/earth-observation-<br/>2281 data/near-real-time/firms/active-fire-data). 2021.
- 2282 79. NASA Socioeconomic Data and Applications Center (SEDAC) Gridded Population of the World  
2283 (GPWv4). Available at <https://beta.sedac.ciesin.columbia.edu/data/collection/gpw-v4>. 2021.
- 2284 80. Vitolo C, Di Giuseppe F, Barnard C, et al. ERA5-based global meteorological wildfire danger  
2285 maps. *Sci Data* 2020; **7**(1): 216.
- 2286 81. Royal Commission into National Natural Disaster Arrangements. Royal Commission into  
2287 National Natural Disaster Arrangements - Report, 2020.
- 2288 82. Stanke C, Kerac M, Prudhomme C, Medlock J, Murray V. Health effects of drought: a  
2289 systematic review of the evidence. *PLoS currents* 2013; **5**.
- 2290 83. Thomas MA, Lin T. Illustrative Analysis of Probabilistic Sea Level Rise Hazard. *Journal of  
2291 Climate* 2020; **33**(4): 1523-34.

- 2292 84. Centre for Research on the Epidemiology of Disasters. EM-DAT The International Disaster  
2293 Database. Available at <https://emdat.be/>. 2021.
- 2294 85. Caminade C, McIntyre KM, Jones AE. Impact of recent and future climate change on vector-  
2295 borne diseases. *Ann N Y Acad Sci* 2019; **1436**(1): 157-73.
- 2296 86. Semenza JC, Herbst S, Rechenburg A, et al. Climate Change Impact Assessment of Food- and  
2297 Waterborne Diseases. *Crit Rev Environ Sci Technol* 2012; **42**(8): 857-90.
- 2298 87. Iwamura T, Guzman-Holst A, Murray KA. Accelerating invasion potential of disease vector  
2299 *Aedes aegypti* under climate change. *Nature Communications* 2020; **11**(1): 2130.
- 2300 88. Ryan SJ, Carlson CJ, Mordecai EA, Johnson LR. Global expansion and redistribution of *Aedes*-  
2301 borne virus transmission risk with climate change. *PLOS Neglected Tropical Diseases* 2019; **13**(3):  
2302 e0007213.
- 2303 89. Hagenlocher M, Delmelle E, Casas I, Kienberger S. Assessing socioeconomic vulnerability to  
2304 dengue fever in Cali, Colombia: statistical vs expert-based modeling. *International Journal of Health*  
2305 *Geographics* 2013; **12**(1): 36.
- 2306 90. Vincenti-Gonzalez MF, Grillet M-E, Velasco-Salas ZI, et al. Spatial Analysis of Dengue  
2307 Seroprevalence and Modeling of Transmission Risk Factors in a Dengue Hyperendemic City of  
2308 Venezuela. *PLOS Neglected Tropical Diseases* 2017; **11**(1): e0005317.
- 2309 91. FAO. The state of food security and nutrition in the world. Rome, Italy: Food and Agriculture  
2310 Organization of the United Nations, 2020.
- 2311 92. Botreau H, Cohen MJ. Gender inequality and food insecurity: A dozen years after the food  
2312 price crisis, rural women still bear the brunt of poverty and hunger. *Advances in Food Security and*  
2313 *Sustainability* 2020; **5**: 53-117.
- 2314 93. Craufurd PQ, Wheeler TR. Climate change and the flowering time of annual crops. *Journal of*  
2315 *Experimental Botany* 2009; **60**(9): 2529-39.
- 2316 94. FAO. The State of World Fisheries and Aquaculture 2020. Sustainability in action. Rome: Food  
2317 and Agriculture Organization, 2020.
- 2318 95. Lynn K, Daigle J, Hoffman J, et al. The impacts of climate change on tribal traditional foods.  
2319 *Climatic Change* 2013; **120**(3): 545-56.
- 2320 96. Allison EH, Perry AL, Badjeck MC, et al. Vulnerability of national economies to the impacts of  
2321 climate change on fisheries. *Fish Fish* 2009; **10**(2): 173-96.
- 2322 97. Barange M, Bahri T, Beveridge MC, Cochrane KL, Funge-Smith S, Poulain F. Impacts of climate  
2323 change on fisheries and aquaculture: synthesis of current knowledge, adaptation and mitigation  
2324 options: FAO; 2018.
- 2325 98. FAO. New Food Balance Sheets. Available at <http://www.fao.org/faostat/en/#data/FBS>. 2021.
- 2326 99. Pörtner H, Roberts D, Masson-Delmotte V, et al. IPCC Special Report on the Ocean and  
2327 Cryosphere in a Changing Climate. *IPCC Intergovernmental Panel on Climate Change: Geneva,*  
2328 *Switzerland* 2019.
- 2329 100. Melet A, Meyssignac B, Almar R, Le Cozannet G. Under-estimated wave contribution to coastal  
2330 sea-level rise. *Nature Climate Change* 2018; **8**(3): 234-9.
- 2331 101. Kirezci E, Young IR, Ranasinghe R, et al. Projections of global-scale extreme sea levels and  
2332 resulting episodic coastal flooding over the 21st Century. *Sci Rep* 2020; **10**(1): 11629.
- 2333 102. Le Bars D, Drijfhout S, de Vries H. A high-end sea level rise probabilistic projection including  
2334 rapid Antarctic ice sheet mass loss. *Environmental Research Letters* 2017; **12**(4): 044013.
- 2335 103. Bakker AMR, Wong TE, Ruckert KL, Keller K. Sea-level projections representing the deeply  
2336 uncertain contribution of the West Antarctic ice sheet. *Sci Rep* 2017; **7**(1): 3880.
- 2337 104. Rose AN, McKee JJ, Sims KM, Bright EA, Reith AE, Urban ML. LandScan 2019. 2019 ed. Oak  
2338 Ridge, TN: Oak Ridge National Laboratory; 2020.



- 2339 105. Kulp SA, Strauss BH. CoastalDEM: a global coastal digital elevation model improved from SRTM  
2340 using a neural network. *Remote sensing of environment* 2018; **206**: 231-9.
- 2341 106. Vineis P, Chan Q, Khan A. Climate change impacts on water salinity and health. *Journal of*  
2342 *Epidemiology and Global Health* 2011; **1**(1): 5-10.
- 2343 107. Dvorak AC, Solo-Gabriele HM, Galletti A, et al. Possible impacts of sea level rise on disease  
2344 transmission and potential adaptation strategies, a review. *Journal of Environmental Management*  
2345 2018; **217**: 951-68.
- 2346 108. Schütte S, Gemenne F, Zaman M, Flahault A, Depoux A. Connecting planetary health, climate  
2347 change, and migration. *The Lancet Planetary Health* 2018; **2**(2): e58-e9.
- 2348 109. Dannenberg AL, Frumkin H, Hess JJ, Ebi KL. Managed retreat as a strategy for climate change  
2349 adaptation in small communities: public health implications. *Climatic change* 2019; **153**(1-2): 1-14.
- 2350 110. Ayeb-Karlsson S, Kniveton D, Cannon T. Trapped in the prison of the mind: Notions of climate-  
2351 induced (im)mobility decision-making and wellbeing from an urban informal settlement in  
2352 Bangladesh. *Palgr Commun* 2020; **6**(1).
- 2353 111. Schwerdtle PN, McMichael C, Mank I, Sauerborn R, Danquah I, Bowen KJ. Health and migration  
2354 in the context of a changing climate: A systematic literature assessment. *Environmental Research*  
2355 *Letters* 2020; **15**(10): 103006.
- 2356 112. Glazebrook T, Noll S, Opoku E. Gender Matters: Climate Change, Gender Bias, and Women's  
2357 Farming in the Global South and North. *Agriculture (Switzerland)* 2020; **10**(267): 1-25.
- 2358 113. Chingarande D, Huyer S, Lanzarini S, et al. Background paper on mainstreaming gender into  
2359 National Adaptation Planning and implementation in Sub-Saharan Africa. Wageningen, the  
2360 Netherland, 2020.
- 2361 114. Delaney P, Shrader E. Gender and Post-Disaster Reconstruction: The Case of Hurricane Mitch  
2362 in Honduras and Nicaragua, 2000.
- 2363 115. Roehr U. Gender, climate change and adaptation. Introduction to the gender dimensions.  
2364 August, 2007.
- 2365 116. Schipper ELF, Ensor J, Mukherji A, et al. Equity in climate scholarship: a manifesto for action.  
2366 *Climate and Development* 2021: 1-4.
- 2367 117. Ampaire EL, Acosta M, Huyer S, et al. Gender in climate change, agriculture, and natural  
2368 resource policies: insights from East Africa. *Climatic Change* 2020; **158**: 43-60.
- 2369 118. Butler J. Gender Trouble: Feminism and the Subversion of Identity. New York; 1990.
- 2370 119. Colebrook C. Gender. Hampshire, UK; 2004.
- 2371 120. Cornwall A, Harrison E, Whitehead A. Gender Myths and Feminist Fables: The Struggle for  
2372 Interpretive Power in Gender and Development. *Development and Change* 2007; **38**(1): 1-20.
- 2373 121. Kabeer N. Gender equality and women's empowerment: A critical analysis of the Third  
2374 Millennium Development Goal. *Gender and Development* 2005; **13**(1): 13- 24.
- 2375 122. Azcona G, Valero SD. Making women and girls visible: Data for gender equality, 2018.
- 2376 123. Equal M. SDG Gender Index. 2021.
- 2377 124. Sanga D. Addressing Gender Issues through the Production and Use of Gender-Sensitive  
2378 Information. *The African Statistical Journal* 2008; **7**: 116-39.
- 2379 125. World Economic F. Global Gender Gap Report; 2020.
- 2380 126. Center on Gender Equity and Health U, Data2X, Foundation BMG, et al. Strengthening gender  
2381 measures and data in the COVID-19 era: an urgent need for change, 2021.
- 2382 127. U. N. Women. Making every women and girl count: Supporting the monitoring and  
2383 implementation of the SDGs through better production and use of gender statistics. May, 2016.
- 2384 128. ESCAP. Work of the secretariat and partners on mainstreaming gender in environment  
2385 statistics. Bangkok: Economic and Social Commission for Asia and the Pacific 2020.

- 2386 129. Ibn-Mohammed T, Mustapha KB, Godsell J, et al. A critical analysis of the impacts of COVID-  
2387 19 on the global economy and ecosystems and opportunities for circular economy strategies.  
2388 *Resources, Conservation and Recycling* 2021; **164**: 105169.
- 2389 130. Nicola M, Alsafi Z, Sohrabi C, et al. The socio-economic implications of the coronavirus  
2390 pandemic (COVID-19): A review. *Int J Surg* 2020; **78**: 185-93.
- 2391 131. Ebi K, Boyer C, Bowen K, Frumkin H, Hess J. Monitoring and Evaluation Indicators for Climate  
2392 Change-Related Health Impacts, Risks, Adaptation, and Resilience. *International Journal of*  
2393 *Environmental Research and Public Health* 2018; **15**(9): 1943.
- 2394 132. WHO. WHO Health and Climate Change Survey Report: Tracking Global Progress. Geneva,  
2395 Switzerland: World Health Organization, 2019.
- 2396 133. WHO. Gender, Climate Change and Health, 2014.
- 2397 134. WHO. Mainstreaming gender in health adaptation to climate change programmes. Geneva,  
2398 Switzerland, 2012.
- 2399 135. Moloney A. How COVID-19 is exposing 'hidden poverty' across unequal cities. World Economic  
2400 Forum. 2020 2020/11.
- 2401 136. Sharifi A, Khavarian-Garmsir AR. The COVID-19 pandemic: Impacts on cities and major lessons  
2402 for urban planning, design, and management. *Science of The Total Environment* 2020; **749**: 142391.
- 2403 137. WBG. Urban Development. 2020.  
2404 <https://www.worldbank.org/en/topic/urbandevelopment/overview> (accessed 28 April 2020).
- 2405 138. Wong MCS, Huang J, Wong SH, Yuen-Chun Teoh J. The potential effectiveness of the WHO  
2406 International Health Regulations capacity requirements on control of the COVID-19 pandemic: a cross-  
2407 sectional study of 114 countries. *J R Soc Med* 2021; **114**(3): 121-31.
- 2408 139. Bouchama A, Dehbi M, Mohamed G, Matthies F, Shoukri M, Menne B. Prognostic factors in  
2409 heat wave related deaths: a meta-analysis. *Archives of internal medicine* 2007; **167**(20): 2170-6.
- 2410 140. Miettinen OS. Proportion of disease caused or prevented by a given exposure, trait or  
2411 intervention. *American journal of epidemiology* 1974; **99**(5): 325-32.
- 2412 141. Birol F. The future of cooling: opportunities for energy-efficient air conditioning. Paris, France:  
2413 International Energy Agency, 2018.
- 2414 142. Hospers L, Smallcombe JW, Morris NB, Capon A, Jay O. Electric fans: A potential stay-at-home  
2415 cooling strategy during the COVID-19 pandemic this summer? *Science of the Total Environment* 2020;  
2416 **747**: 141180.
- 2417 143. Salamanca F, Georgescu M, Mahalov A, Moustouli M, Wang M. Anthropogenic heating of the  
2418 urban environment due to air conditioning. *Journal of Geophysical Research: Atmospheres* 2014;  
2419 **119**(10): 5949-65.
- 2420 144. Waite M, Cohen E, Torbey H, Piccirilli M, Tian Y, Modi V. Global trends in urban electricity  
2421 demands for cooling and heating. *Energy* 2017; (127): 786-802.
- 2422 145. Randazzo T, De Cian E, Mistry MN. Air conditioning and electricity expenditure: The role of  
2423 climate in temperate countries. *Economic Modelling* 2020; **90**: 273-87.
- 2424 146. Mastrucci A, Byers E, Pachauri S, Rao ND. Improving the SDG energy poverty targets:  
2425 Residential cooling needs in the Global South. *Energy and Buildings* 2019; **186**: 405-15.
- 2426 147. Chen F, Kusaka H, Bornstein R, et al. The integrated WRF/urban modelling system:  
2427 development, evaluation, and applications to urban environmental problems. *International Journal of*  
2428 *Climatology* 2011; **31**(2): 273-88.
- 2429 148. Heaviside C, Cai XM, Vardoulakis S. The effects of horizontal advection on the urban heat  
2430 island in Birmingham and the West Midlands, United Kingdom during a heatwave. *Q J Roy Meteor Soc*  
2431 2015; **141**(689): 1429-41.
- 2432 149. Heaviside C, Vardoulakis S, Cai XM. Attribution of mortality to the urban heat island during  
2433 heatwaves in the West Midlands, UK. *Environmental Health: A Global Access Science Source* 2016; **15**.

2434 150. Macintyre HL, Heaviside C, Taylor J, et al. Assessing urban population vulnerability and  
2435 environmental risks across an urban area during heatwaves – Implications for health protection.  
2436 *Science of the Total Environment* 2018; **610-611**: 678-90.

2437 151. Macintyre H, Heaviside C. Potential benefits of cool roofs in reducing heat-related mortality  
2438 during heatwaves in a European city. *Environment international* 2019; **127**: 430-41.

2439 152. Macintyre HL, Heaviside C, Cai X, Phalkey R. The winter urban heat island: Impacts on cold-  
2440 related mortality in a highly urbanized European region for present and future climate. *Environment*  
2441 *International* 2021; **154**: 106530.

2442 153. Macintyre HL, Heaviside C, Cai X, Phalkey R. Comparing temperature-related mortality  
2443 impacts of cool roofs in winter and summer in a highly urbanized European region for present and  
2444 future climate. *Environment International* 2021; **154**: 106606.

2445 154. He C, Zhao J, Zhang Y, et al. Cool Roof and Green Roof Adoption in a Metropolitan Area:  
2446 Climate Impacts during Summer and Winter. *Environmental Science & Technology* 2020; **54**(17):  
2447 10831-9.

2448 155. Kardan O, Gozdyra P, Misic B, et al. Neighborhood greenspace and health in a large urban  
2449 center. *Sci Rep* 2015; **5**(1): 11610.

2450 156. Gascon M, Triguero-Mas M, Martínez D, et al. Residential green spaces and mortality: A  
2451 systematic review. *Environment International* 2016; **86**: 60-7.

2452 157. Abelt K, McLafferty S. Green Streets : Urban Green and Birth Outcomes. *Int J Environ Res Public*  
2453 *Health* 2017; **14**(771).

2454 158. Aronson MF, Lepczyk CA, Evans KL, et al. Biodiversity in the city: key challenges for urban  
2455 green space management. *Frontiers in Ecology and the Environment* 2017; **15**(4): 189-96.

2456 159. Ode Sang Å, Knez I, Gunnarsson B, Hedblom M. Urban Forestry & Urban Greening The effects  
2457 of naturalness, gender, and age on how urban green space is perceived and used. *Urban Forestry &*  
2458 *Urban Greening* 2016; **18**: 268-76.

2459 160. Rahman KMA, Zhang D. Analyzing the Level of Accessibility of Public Urban Green Spaces to  
2460 Different Socially Vulnerable Groups of People. *Sustainability* 2018; **10**(3917).

2461 161. Richardson EA, Mitchell R. Gender differences in relationships between urban green space  
2462 and health in the United Kingdom. *Soc Sci Med* 2010; **71**(3): 568-75.

2463 162. Schipperijn J, Ekholm O, Stigsdotter UK, et al. Landscape and Urban Planning Factors  
2464 influencing the use of green space : Results from a Danish national representative survey. *Landscape*  
2465 *and Urban Planning* 2010; **95**: 130-7.

2466 163. Mitchell R, Popham F. Effect of exposure to natural environment on health inequalities: an  
2467 observational population study. *Lancet* 2008; **372**(9650): 1655-60.

2468 164. Geary RS, Wheeler B, Lovell R, Jepson R, Hunter R, Rodgers S. A call to action: Improving urban  
2469 green spaces to reduce health inequalities exacerbated by COVID-19. *Preventive Medicine* 2021; **145**:  
2470 106425.

2471 165. UN-Habitat, World Health Organization. Integrating health in urban and territorial planning.  
2472 Geneva, Switzerland: UN-Habitat and World Health Organization, 2020.

2473 166. International Monetary F. World Economic Outlook Database. 2018.

2474 167. kMatrix Ltd. Adaptation and Resilience to Climate Change dataset. Rutland, UK; 2021.

2475 168. UNEP, UNEP DTU Partnership. Emissions Gap Report 2020. 2020.  
2476 <https://www.unep.org/emissions-gap-report-2020>.

2477 169. Le Quéré C, Peters GP, Friedlingstein P, et al. Fossil CO2 emissions in the post-COVID-19 era.  
2478 *Nature Climate Change* 2021; **11**(3): 197-9.

2479 170. UNEP. Are We Building Back Better? Evidence from 2020 and Pathways for Inclusive Green  
2480 Recovery Spending. Nairobi, Kenya: UN Environment Programme, 2021.

2481 171. Hendryx M, Zullig KJ, Luo J. Impacts of Coal Use on Health. *Annu Rev Public Health* 2020; **41**(1):  
2482 397-415.

2483 172. Matthews HD, Tokarska KB, Nicholls ZRJ, et al. Opportunities and challenges in using  
2484 remaining carbon budgets to guide climate policy. *Nature Geoscience* 2020; **13**(12): 769-79.

2485 173. IEA. World Energy Outlook 2020 Paris, France: International Energy Agency, 2021.

2486 174. WHO. World health statistics 2021: monitoring health for the SDGs, sustainable development  
2487 goals. Geneva, Switzerland: World Health Organization, 2021.

2488 175. Cozzi L, Contejean A, Samantar J, Dasgupta A, Rouget A, Arboleya L. The Covid-19 crisis is  
2489 reversing progress on energy access in Africa. Paris: IEA, 2020.

2490 176. Shupler M, Mwitari J, Gohole A, et al. COVID-19 Lockdown in a Kenyan Informal Settlement:  
2491 Impacts on Household Energy and Food Security. *medRxiv* 2020: 2020.05.27.20115113.

2492 177. EUROSTAT. People unable to keep their home adequately warm, 2019. 2020 2020.  
2493 [https://ec.europa.eu/eurostat/en/web/products-eurostat-news/-/ddn-20210106-](https://ec.europa.eu/eurostat/en/web/products-eurostat-news/-/ddn-20210106-1?redirect=/eurostat/en/news/whats-new)  
2494 [1?redirect=/eurostat/en/news/whats-new](https://ec.europa.eu/eurostat/en/news/whats-new) (accessed 14 April 2021).

2495 178. Kolokotsa D, Santamouris M. Review of the indoor environmental quality and energy  
2496 consumption studies for low income households in Europe. *Science of The Total Environment* 2015;  
2497 **536**: 316-30.

2498 179. Thomson H, Simcock N, Bouzarovski S, Petrova S. Energy poverty and indoor cooling: An  
2499 overlooked issue in Europe. *Energy and Buildings* 2019; **196**: 21-9.

2500 180. IEA. World Extended Energy Balances (2020 edition). Paris: IEA, 2021.

2501 181. Stoner O, Shaddick G, Economou T, et al. Global household energy model: a multivariate  
2502 hierarchical approach to estimating trends in the use of polluting and clean fuels for cooking. *Journal*  
2503 *of the Royal Statistical Society: Series C (Applied Statistics)* 2020; **69**(4): 815-39.

2504 182. WHO. Household Energy Database. Geneva: World Health Organization; 2021.

2505 183. WHO. Air quality guidelines – global update 2005. 2005.  
2506 [https://www.who.int/airpollution/publications/aqg2005/en/#:~:text=WHO%20Air%20Quality%20Gu-](https://www.who.int/airpollution/publications/aqg2005/en/#:~:text=WHO%20Air%20Quality%20Guidelines%20(AQG,they%20were%20revised%20in%201997)  
2507 [idelines%20\(AQG,they%20were%20revised%20in%201997](https://www.who.int/airpollution/publications/aqg2005/en/#:~:text=WHO%20Air%20Quality%20Guidelines%20(AQG,they%20were%20revised%20in%201997) (accessed 8 May 2021).

2508 184. Bennitt FB, Wozniak SS, Causey K, Burkart K, Brauer M. Estimating disease burden attributable  
2509 to household air pollution: new methods within the Global Burden of Disease Study. *The Lancet Global*  
2510 *Health* 2021; **9**: S18.

2511 185. Shupler M, Hystad P, Birch A, et al. Household and personal air pollution exposure  
2512 measurements from 120 communities in eight countries: results from the PURE-AIR study. *The Lancet*  
2513 *Planetary Health* 2020; **4**(10): e451-e62.

2514 186. Shupler M, Godwin W, Frostad J, Gustafson P, Arku RE, Brauer M. Global estimation of  
2515 exposure to fine particulate matter (PM<sub>2.5</sub>) from household air pollution. *Environment international*  
2516 2018; **120**: 354-63.

2517 187. Clougherty JE. Review A Growing Role for Gender Analysis in Air Pollution Epidemiology.  
2518 *Environmental Health Perspectives* 2010; **118**(2): 167-76.

2519 188. Oparaocha S, Dutta S. Gender and energy for sustainable development. *Current Opinion in*  
2520 *Environmental Sustainability* 2011; **3**: 265-71.

2521 189. Gordon SB, Bruce NG, Grigg J, et al. Respiratory risks from household air pollution in low and  
2522 middle income countries. *The Lancet Respiratory Medicine* 2014; **2**(10): 823-60.

2523 190. Kurata M, Takahashi K, Hibiki A. Gender differences in associations of household and ambient  
2524 air pollution with child health: Evidence from household and satellite-based data in Bangladesh. *World*  
2525 *Dev* 2020; **128**: 104779.

2526 191. European Commission. Revision of the Ambient Air Quality Directives. 2021.  
2527 [https://ec.europa.eu/environment/air/quality/revision\\_of\\_the\\_aaq\\_directives.htm](https://ec.europa.eu/environment/air/quality/revision_of_the_aaq_directives.htm) (accessed 18  
2528 May 2021).

2529 192. Dyer C. Air pollution from road traffic contributed to girl's death from asthma, coroner  
2530 concludes. *British Medical Journal Publishing Group*; 2020.

2531 193. Murray CJL, Aravkin AY, Zheng P, et al. Global burden of 87 risk factors in 204 countries and  
2532 territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet*  
2533 2020; **396**(10258): 1223-49.

2534 194. IEA. Transport. 2021. <https://www.iea.org/topics/transport> (accessed 14 April 2021).

2535 195. Hamilton I, Kennard H, McGushin A, et al. The public health implications of the Paris  
2536 Agreement: a modelling study. *The Lancet Planetary Health* 2021; **5**(2): e74-e83.

2537 196. Goel R, Goodman A, Aldred R, et al. Cycling behaviour in 17 countries across 6 continents:  
2538 levels of cycling, who cycles, for what purpose, and how far? *Transport Reviews* 2021: 1-24.

2539 197. Ferla M, Graham A. Women slowly taking off: An investigation into female  
2540 underrepresentation in commercial aviation. *Research in Transportation Business and Management*  
2541 2019; **31**(September): 100378.

2542 198. Adlakha D, Parra DC. Mind the gap: Gender differences in walkability , transportation and  
2543 physical activity in urban India. *Journal of Transport & Health* 2020; **18**(100875): 1-17.

2544 199. Mackett RL. The health implications of inequalities in travel. *Journal of Transport & Health*  
2545 2014; **1**(3): 202-9.

2546 200. Fraszczyk A, Piip J. A review of transport organisations for female professionals and their  
2547 impacts on the transport sector workforce. *Research in Transportation Business and Management*  
2548 2019; **31**(November): 100379.

2549 201. Olsen JR, Mitchell R, Mutrie N, Foley L, Ogilvie D, study M. Population levels of, and  
2550 inequalities in, active travel: A national, cross-sectional study of adults in Scotland. *Prev Med Rep* 2017;  
2551 **8**: 129-34.

2552 202. IEA. Global EV Outlook 2020. 2020. <https://www.iea.org/reports/global-ev-outlook-2020>  
2553 (accessed 15 April 2021).

2554 203. IEA. Tracking Transport 2020. 2020. <https://www.iea.org/reports/tracking-transport-2020>  
2555 (accessed 15 April 2020).

2556 204. IEA. Extended world energy balances. IEA World Energy Statistics and Balances. Paris, France:  
2557 IEA; 2021.

2558 205. IEA. Changes in transport behaviour during the Covid-19 crisis. 2021.  
2559 <https://www.iea.org/articles/changes-in-transport-behaviour-during-the-covid-19-crisis> (accessed 15  
2560 April 2021).

2561 206. IEA. The Covid-19 Crisis and Clean Energy Progress. 2021. [https://www.iea.org/reports/the-](https://www.iea.org/reports/the-covid-19-crisis-and-clean-energy-progress)  
2562 [covid-19-crisis-and-clean-energy-progress](https://www.iea.org/reports/the-covid-19-crisis-and-clean-energy-progress) (accessed 15 April 2021).

2563 207. Füzéki E, Schröder J, Carraro N, et al. Physical Activity during the First COVID-19-Related  
2564 Lockdown in Italy. *International Journal of Environmental Research and Public Health* 2021; **18**(5):  
2565 2511.

2566 208. Bechauf R. Cycling and COVID-19: why investments to boost cycling are important for a  
2567 sustainable recovery. 2020. [https://www.iisd.org/sustainable-recovery/cycling-and-covid-19-why-](https://www.iisd.org/sustainable-recovery/cycling-and-covid-19-why-investments-to-boost-cycling-are-important-for-a-sustainable-recovery/)  
2568 [investments-to-boost-cycling-are-important-for-a-sustainable-recovery/](https://www.iisd.org/sustainable-recovery/cycling-and-covid-19-why-investments-to-boost-cycling-are-important-for-a-sustainable-recovery/) (accessed 14 April 2021).

2569 209. Mbow CR, L. Barioni, T. Benton, M. Herrero, M. Krishnapillai, E. Liwenga, P. Pradhan, M.  
2570 Rivera-, Ferre TS, F. Tubiello, and Y. Xu, . Food security. In: Change IPoC, ed. Climate change and land:  
2571 an IPCC special report on climate change, desertification, land degradation, sustainable land  
2572 management, food security, and greenhouse gas fluxes in terrestrial ecosystems. London: Working  
2573 Group III Technical Support Unit; 2019.

2574 210. FAO. FAOSTAT. 2020. <http://www.fao.org/faostat/> (accessed 15 April 2021).

2575 211. Iannotti L, Tarawali S, Baltenweck I, et al. Livestock-derived foods and sustainable healthy  
2576 diets. Rome, Italy: UN Nutrition Secretariat, 2021.

- 2577 212. Gerber PJ, Steinfeld H, Henderson B, et al. Tackling climate change through livestock – A global  
2578 assessment of emissions and mitigation opportunities.  
2579 . Rome: FAO, 2013.
- 2580 213. Global Alliance for Improved Nutrition. GAIN Briefing Paper Series 2 - Animal-source foods for  
2581 human and planetary health. Geneva, Switzerland: Global Alliance for Improved Nutrition (GAIN),  
2582 2020.
- 2583 214. Food and Agriculture Organization of the United Nations. Food balance sheets. 2020.
- 2584 215. Wang H, Abbas KM, Abbasifard M, et al. Global age-sex-specific fertility, mortality, healthy life  
2585 expectancy (HALE), and population estimates in 204 countries and territories, 1950–2019: a  
2586 comprehensive demographic analysis for the Global Burden of Disease Study 2019. *The Lancet* 2020;  
2587 **396**(10258): 1160-203.
- 2588 216. Chai BC, Voort VDJR, Grofelnik K, Eliasdottir HG, Klöss I, Perez-cueto FJA. Which Diet Has the  
2589 Least Environmental Impact on Our Planet ? A Systematic Review of Vegan , Vegetarian and  
2590 Omnivorous Diets. *Sustainability* 2019; **11**(4110).
- 2591 217. Modlinska K, Adamczyk D, Maison D, Pisula W. Gender Differences in Attitudes to Vegans/  
2592 Vegetarians and Their Food Preferences, and Their Implications for Promoting Sustainable Dietary  
2593 Patterns – A Systematic Review. *Sustainability* 2020; **12**(6292): 1-17.
- 2594 218. Rosenfeld DL, Rothgerber H, Tomiyama AJ. Mostly Vegetarian , But Flexible About It :  
2595 Investigating How Meat-Reducers Express Social Identity Around Their Diets. *Social Psychological and*  
2596 *Personality Science* 2020; **11**(3): 406-15.
- 2597 219. Song S, Kim J, Kim J. Gender Differences in the Association between Dietary Pattern and the  
2598 Incidence of Hypertension in Middle-Aged and Older Adults. *Nutrients* 2018; **10**(252).
- 2599 220. European Academies Science Advisory Council, Federation of European Academies of  
2600 Medicine. Decarbonisation of the Health Sector: A Commentary by EASAC and FEAM. Brussels,  
2601 Belgium: European Academies Science Advisory Council, 2021.
- 2602 221. Health Care Without Harm. Race to Zero. 2021.  
2603 <https://healthcareclimateaction.org/racetozero> (accessed 24 June 2021).
- 2604 222. Hepburn C, O’Callaghan B, Stern N, Stiglitz J, Zenghelis D. Will COVID-19 fiscal recovery  
2605 packages accelerate or retard progress on climate change? *Oxford Review of Economic Policy* 2020;  
2606 **36**: S359-S81.
- 2607 223. International Monetary Fund. World Economic Outlook Update April 2021. Washington, DC,  
2608 USA: International Monetary Fund, 2021.
- 2609 224. Swiss Re. Sigma explorer. Zurich, Switzerland: Swiss Re; 2021.
- 2610 225. Stringhini S, Carmeli C, Jokela M. Socioeconomic status and the 25 x 25 risk factors as  
2611 determinants of premature mortality: a multicohort study and meta-analysis of 1.7 million men and  
2612 women (vol 389, pg 1229, 2017). *Lancet* 2017; **389**(10075): 1194-.
- 2613 226. ILO. ILOSTAT Statistics on wages. 2021. <https://ilostat.ilo.org/topics/wages/> (accessed 1st  
2614 April 2021).
- 2615 227. Muriithi MK, Mutegi RG, Mwabu G. Counting unpaid work in Kenya: Gender and age profiles  
2616 of hours worked and imputed wage incomes. *The Journal of the Economics of Ageing* 2020; **17**:  
2617 100120.
- 2618 228. Reddy AA, Mittal S, Roy NS, Kanjilal-Bhaduri S. Time Allocation between Paid and Unpaid Work  
2619 among Men and Women: An Empirical Study of Indian Villages. *Sustainability* 2021; **13**(5): 17.
- 2620 229. Sarker MR. Labor market and unpaid works implications of COVID-19 for Bangladeshi women.  
2621 *Gen Work Organ*: 8.
- 2622 230. World Bank. Global Economic Prospects. Washington DC; 2021.
- 2623 231. O’Callaghan B, Murdock E. Are We Building Back Better? Evidence from 2020 and Pathways  
2624 to Inclusive Green Recovery Spending. 2021.

2625 232. Vivid Economics. Greenness of Stimulus Index - February 2021. 2021.

2626 233. Leaton J. Unburnable Carbon – Are the world’s financial markets carrying a carbon bubble?  
2627 *Carbon Tracker* 2011.

2628 234. IEA. After steep drop in early 2020, global carbon dioxide emissions have rebounded strongly  
2629 - News - IEA.

2630 235. WHO Manifesto for a healthy recovery from COVID-19.

2631 236. IEA. Global Energy & CO2 Status Report 2019. 2019 (accessed April 15, 2021).

2632 237. IEA. World Energy Investment. Paris, France: International Energy Agency, 2021.

2633 238. Hendryx M, Zullig KJ, Luo J. Impacts of Coal Use on Health. *Annual Review of Public Health*  
2634 2020; **41**: 397-415.

2635 239. Garrett-Peltier H. Green versus brown: Comparing the employment impacts of energy  
2636 efficiency, renewable energy, and fossil fuels using an input-output model. *Economic Modelling* 2017;  
2637 **61**: 439-47.

2638 240. IRENA. Renewable Energy: A Gender Perspective. Abu Dhabi, 2019.

2639 241. Plantinga A, Scholtens B. The financial impact of fossil fuel divestment. *Climate Policy* 2021;  
2640 **21**: 107-19.

2641 242. Hunt C, Weber O. Fossil Fuel Divestment Strategies: Financial and Carbon-Related  
2642 Consequences. *Organization & Environment* 2019; **32**: 41-61.

2643 243. Phillips M. Exxon Mobil Defeated by Activist Investor Engine No. 1. 2021.  
2644 <https://www.nytimes.com/2021/06/09/business/exxon-mobil-engine-no1-activist.html> (accessed 29  
2645 June 2021).

2646 244. 350.org. Divestment Commitments. 2020.  
2647 <https://gofossilfree.org/divestment/commitments/> (accessed 30 March 2021).

2648 245. Ranger L. Unburnable Carbon 2013 : Wasted capital and stranded assets About the Grantham  
2649 Research Institute on. *Management of Environmental Quality: An International Journal* 2013; **24**: 1-  
2650 40.

2651 246. McGlade C, Ekins P. The geographical distribution of fossil fuels unused when limiting global  
2652 warming to 2°C. *Nature* 2015; **517**: 187-90.

2653 247. Caldecott B, Tilbury, J., Carey, C. Stranded Assets and Scenarios. 2014.

2654 248. Curtin J, McInerney C, Ó Gallachóir B, Hickey C, Deane P, Deeney P. Quantifying stranding risk  
2655 for fossil fuel assets and implications for renewable energy investment: A review of the literature.  
2656 *Renewable and Sustainable Energy Reviews* 2019; **116**: 109402.

2657 249. Leaton J, Fulton M, Spedding P, et al. The \$2 trillion stranded assets danger zone: How fossil  
2658 fuel firms risk destroying investor returns. *Carbon Tracker Initiative* 2015: 1-32.

2659 250. SEI I, ODI, E3G, UNEP. The Production Gap Report: 2020 Special Report. , 2020.

2660 251. Leaton J, Grant A. 2 degrees of separation: Transition risk for oil & gas in a low carbon world.  
2661 *Carbon Tracker* 2017.

2662 252. Coffin M. Absolute Impact: Why oil majors' climate ambitions fall short of Paris limits. *Carbon*  
2663 *Tracker* 2020: xi.

2664 253. IEA. Energy subsidies - tracking the impact of fossil fuel subsidies. 2021.  
2665 <https://www.iea.org/topics/energy-subsidies> (accessed 16th February 2021).

2666 254. OECD. OECD Inventory of support measures for fossil fuels. 2021.  
2667 [https://stats.oecd.org/Index.aspx?DataSetCode=FFS\\_AUS](https://stats.oecd.org/Index.aspx?DataSetCode=FFS_AUS) (accessed 6th April 2021).

2668 255. World Bank. World Bank Carbon Pricing Dashboard. 2021.  
2669 <https://carbonpricingdashboard.worldbank.org/> (accessed 3rd April 2021).

2670 256. WHO. World Health Organization Global Health Expenditure Database. 2021.  
2671 <https://apps.who.int/nha/database/Select/Indicators/en> (accessed 7th April 2021).

2672 257. Younger SD, Osei-Assibey E, Oppong F. Fiscal Incidence in Ghana. *Rev Dev Econ* 2017; **21**(4):  
2673 E47-E66.

2674 258. Inter-agency Task Force on Financing for Development. Financing for Sustainable  
2675 Development Report 2020. New York, NY, USA: UN Department of Economic and Social Affairs, 2020.

2676 259. Mi Z, Zheng J, Meng J, et al. Economic development and converging household carbon  
2677 footprints in China. *Nature Sustainability* 2020; **3**(7): 529-37.

2678 260. Stadler K, Wood R, Bulavskaya T, et al. EXIOBASE 3: Developing a time series of detailed  
2679 environmentally extended multi-regional input-output tables. *Journal of Industrial Ecology* 2018;  
2680 **22**(3): 502-15.

2681 261. Amann M, Kiesewetter G, Schöpp W, et al. Reducing global air pollution: the scope for further  
2682 policy interventions. *Philosophical Transactions of the Royal Society A* 2020; **378**(2183): 20190331.

2683 262. McCann G, Matenga C. COVID-19 and Global Inequality. *COVID-19 IN THE GLOBAL SOUTH*  
2684 2020: 161.

2685 263. UNDP. COVID-19 and Human Development: Assessing the Crisis, Envisioning the Recovery.  
2686 2020. [http://hdr.undp.org/sites/default/files/covid-19\\_and\\_human\\_development\\_0.pdf](http://hdr.undp.org/sites/default/files/covid-19_and_human_development_0.pdf) (accessed  
2687 March-21).

2688 264. UN\_News. Ahead of UN summit, leading scientists warn climate change ‘hitting harder and  
2689 sooner’ than forecast 2019. <https://news.un.org/en/story/2019/09/1046972> (accessed March-21).

2690 265. Intergovernmental\_Panel\_on\_Climate\_Change. Policymakers Summary. In:  
2691 Intergovernmental\_Panel\_on\_Climate\_Change, ed. Climate Change: the IPCC Response Strategies  
2692 Report of the working group III; 1990.

2693 266. Newman N, Fletcher R, Schulz A, Andi S, Nielsen RK. Digital News Report 2020.  
2694 <https://www.digitalnewsreport.org/> (accessed March-21).

2695 267. Barkemeyer R, Figge F, Hoepner A, Holt D, Kraak JM, Yu P-S. Media coverage of climate  
2696 change: An international comparison. 2017; **35**(6): 1029-54.

2697 268. Gavin NT. Addressing climate change: a media perspective. *Environmental Politics* 2009; **18**(5):  
2698 765-80.

2699 269. Boykoff M. Who Speaks for the Climate? Making Sense of Media Reporting on Climate Change:  
2700 Cambridge University Press; 2011.

2701 270. Rogstad I. Is Twitter just rehashing? Intermedia agenda setting between Twitter and  
2702 mainstream media. *Journal of Information Technology & Politics* 2016; **13**(2): 142-58.

2703 271. Duan R, Miller S. Climate change in China: A study of news diversity in party-sponsored and  
2704 market-oriented newspapers. *Journalism* 2019: 1464884919873173.

2705 272. Wang H, Sparks C, Huang Y. Measuring differences in the Chinese press: A study of People’s  
2706 Daily and Southern Metropolitan Daily. *Global Media and China* 2018; **3**(3): 125-40.

2707 273. Auerbach Y, Bloch-Elkon Y. Media Framing and Foreign Policy: The Elite Press vis-à-vis US  
2708 Policy in Bosnia, 1992–95. *Journal of Peace Research* 2005; **42**(1): 83-99.

2709 274. Billett S. Dividing climate change: global warming in the Indian mass media. *Climatic change*  
2710 2010; **99**(1-2): 1-16.

2711 275. Boykoff MT, Boykoff JM. Balance as bias: global warming and the US prestige press. *Global*  
2712 *Environmental Change* 2004; **14**(2): 125-36.

2713 276. Amazon\_Alexa. The top 500 sites on the Web. 2020. <https://www.alexa.com/topsites>.  
2714 (accessed March-21).

2715 277. Schroeder R, Taylor L. Big data and Wikipedia research: social science knowledge across  
2716 disciplinary divides. *Information, Communication & Society* 2015; **18**(9): 1039-56.

2717 278. Mesgari M, Okoli C, Mehdi M, Nielsen FÅ, Lanamäki A. “The sum of all human knowledge”: A  
2718 systematic review of scholarly research on the content of Wikipedia. *Journal of the Association for*  
2719 *Information Science and Technology* 2015; **66**(2): 219-45.



2720 279. Bornmann L. Scientific peer review. *Annual review of information science and technology*  
2721 2011; **45**(1): 197-245.

2722 280. United Nations General Debate of the 75th session of the General Assembly. 2021.

2723 281. Peterson M. General Assembly. In: Daws S, Weiss T, eds. *The Oxford Handbook on the United*  
2724 *Nations* (2nd edition); 2018.

2725 282. Change UNFCCC. The Paris Agreement. [https://unfccc.int/process-and-meetings/the-paris-](https://unfccc.int/process-and-meetings/the-paris-agreement/the-paris-agreement)  
2726 [agreement/the-paris-agreement](https://unfccc.int/process-and-meetings/the-paris-agreement/the-paris-agreement) (accessed March-21).

2727 283. Dasandi N, Graham H, Lampard P, Jankin Mikhaylov S. Engagement with health in national  
2728 climate change commitments under the Paris Agreement: a global mixed-methods analysis of the  
2729 nationally determined contributions. *The Lancet Planetary Health* 2021; **5**(2): e93-e101.

2730 284. Dasandi N, Graham H, Lampard P, Jankin Mikhaylov S. Intergovernmental engagement on  
2731 health impacts of climate change. *Bull World Health Organ* 2021; **99**(2): 102-11B.

2732 285. Wright C, Nyberg D. *Climate Change, Capitalism, and Corporations: Processes of Creative Self-*  
2733 *Destruction*. Cambridge: Cambridge University Press; 2015.

2734 286. Covington H, Thornton J, Hepburn C. Global warming: Shareholders must vote for climate-  
2735 change mitigation. *Nature News* 2016; **530**(7589): 156.

2736 287. Ekwurzel B, Boneham J, Dalton M, et al. The rise in global atmospheric CO<sub>2</sub>, surface  
2737 temperature, and sea level from emissions traced to major carbon producers. *Climatic Change* 2017;  
2738 **144**(4): 579-90.

2739 288. United\_Nations. The United Nations Global Compact. <https://www.unglobalcompact.org/>  
2740 (accessed March-21).

2741 289. Margaretha Jastram S, Klingenberg J. Assessing the Outcome Effectiveness of Multi-  
2742 Stakeholder Initiatives in the Field of Corporate Social Responsibility – The Example of the United  
2743 Nations Global Compact. *Journal of Cleaner Production* 2018; **189**: 775-84.

2744 290. Preet R, Nilsson M, Schumann B, Evengård B. The gender perspective in climate change and  
2745 global health. *Global Health Action* 2010; **3**(1): 5720.

2746 291. Holmberg K, Hellsten I. Gender differences in the climate change communication on Twitter.  
2747 *Internet Research* 2015.

2748 292. Mavisakalyan A, Tarverdi Y. Gender and climate change: Do female parliamentarians make  
2749 difference? *European Journal of Political Economy* 2019; **56**: 151-64.

2750 293. Pearse R. Gender and climate change. *WIREs Climate Change* 2017; **8**(2): e451.

2751 294. Koch-Baumgarten S, Voltmer K. Introduction: Mass Media and Public Policy – Is There a Link?  
2752 In: Koch-Baumgarten S, Voltmer K, eds. *Public Policy and the Mass Media The Interplay of Mass*  
2753 *Communication and Political Decision Making*.

2754 295. World\_Health\_Organization. Sixty-first World Health Assembly.  
2755 <https://www.who.int/mediacentre/events/2008/wha61/en/> (accessed March-21).

2756 296. Nagarathinam S, Bhatta A. Coverage of climate change issues in Indian newspapers and policy  
2757 implications. *Current Science* 2015; **108**(11): 1972-3.

2758 297. Pierre-Louis K. Heat Waves in the Age of Climate Change: Longer, More Frequent and More  
2759 Dangerous. *The New York Times*. 2020 June 18.

2760 298. HTCorrespondent. Need multi-sectoral coordination to tackle health impacts of climate  
2761 change. *Hindustan Times*. 2020 February 25.

2762 299. Guterres A. A Time to Save the Sick and Rescue the Planet. *The New York Times*. 2020 April  
2763 28.

2764 300. Giles J. Internet encyclopaedias go head to head. *Nature* 2005; **438**(7070): 900-1.

2765 301. Yoshida M, Arase Y, Tsunoda T, Yamamoto M. Wikipedia Page View Reflects Web Search  
2766 Trend. *Proceedings of the ACM Web Science Conference*. Oxford, United Kingdom: Association for  
2767 Computing Machinery; 2015. p. Article 65.

2768 302. Wikipedia. Most popular edition of Wikipedia by country.  
 2769 <https://commons.wikimedia.org/w/index.php?curid=99613651> (accessed March-21.  
 2770 303. Wikipedia. Wikipedia page views by language over time.  
 2771 <https://commons.wikimedia.org/w/index.php?curid=99654507> (accessed March-21.  
 2772 304. Molek-Kozakowska K. Popularity-driven science journalism and climate change: A critical  
 2773 discourse analysis of the unsaid. *Discourse, Context & Media* 2018; **21**: 73-81.  
 2774 305. Baturo A, Dasandi N, Mikhaylov SJ. Understanding state preferences with text as data:  
 2775 Introducing the UN General Debate corpus. *Research & Politics* 2017; **4**(2): 2053168017712821.  
 2776 306. General Assembly of the United Nations. General Debate of the 75th Session 22 to 26  
 2777 September and 29 September 2020. 2020. <https://gadebate.un.org/generaldebate75/en/> (accessed  
 2778 13 April 2021).  
 2779 307. UNFCCC. Paris Agreement. 2015.  
 2780 308. UNFCCC. UNFCCC NDC Registry. 2021.  
 2781 <https://www4.unfccc.int/sites/ndcstaging/Pages/Home.aspx> (accessed 1 April 2021).  
 2782 309. Voegtlin C, Pless NM. Global Governance: CSR and the Role of the UN Global Compact. *Journal*  
 2783 *of Business Ethics* 2014; **122**(2): 179-91.

2784