



Autotrophic fixation of geogenic CO₂ by microorganisms contributes to soil organic matter formation and alters isotope signatures in a wetland mofette

M. E. Nowak¹, F. Beulig², J. von Fischer³, J. Muhr¹, K. Küsel², and S. E. Trumbore¹

¹Department for Biogeochemical Processes, Max Planck Institute for Biogeochemistry, Hans-Knöll Str. 10, 07745 Jena, Germany

²Aquatic Geomicrobiology, Institute of Ecology, Friedrich Schiller University Jena, Dornburger Str. 159, 07743 Jena, Germany

³Department of Biology, Colorado State University, Fort Collins, CO 80523, USA

Correspondence to: M. Nowak (mnowak@bgc-jena.mpg.de)

Received: 24 July 2015 – Published in Biogeosciences Discuss.: 4 September 2015

Revised: 17 November 2015 – Accepted: 20 November 2015 – Published: 8 December 2015

Abstract. To quantify the contribution of autotrophic microorganisms to organic matter (OM) formation in soils, we investigated natural CO₂ vents (mofettes) situated in a wetland in northwest Bohemia (Czech Republic). Mofette soils had higher soil organic matter (SOM) concentrations than reference soils due to restricted decomposition under high CO₂ levels. We used radiocarbon ($\Delta^{14}\text{C}$) and stable carbon ($\delta^{13}\text{C}$) isotope ratios to characterize SOM and its sources in two mofettes and compared it with respective reference soils, which were not influenced by geogenic CO₂.

The geogenic CO₂ emitted at these sites is free of radiocarbon and enriched in ¹³C compared to atmospheric CO₂. Together, these isotopic signals allow us to distinguish C fixed by plants from C fixed by autotrophic microorganisms using their differences in ¹³C discrimination. We can then estimate that up to 27 % of soil organic matter in the 0–10 cm layer of these soils was derived from microbially assimilated CO₂.

Isotope values of bulk SOM were shifted towards more positive $\delta^{13}\text{C}$ and more negative $\Delta^{14}\text{C}$ values in mofettes compared to reference soils, suggesting that geogenic CO₂ emitted from the soil atmosphere is incorporated into SOM. To distinguish whether geogenic CO₂ was fixed by plants or by CO₂ assimilating microorganisms, we first used the proportional differences in radiocarbon and $\delta^{13}\text{C}$ values to indicate the magnitude of discrimination of the stable isotopes in living plants. Deviation from this relationship was taken to indicate the presence of microbial CO₂ fixation, as

microbial discrimination should differ from that of plants. ¹³CO₂-labelling experiments confirmed high activity of CO₂ assimilating microbes in the top 10 cm, where $\delta^{13}\text{C}$ values of SOM were shifted up to 2 ‰ towards more negative values. Uptake rates of microbial CO₂ fixation ranged up to $1.59 \pm 0.16 \mu\text{g g}_{\text{dw}}^{-1} \text{d}^{-1}$. We inferred that the negative $\delta^{13}\text{C}$ shift was caused by the activity of autotrophic microorganisms using the Calvin–Benson–Bassham (CBB) cycle, as indicated from quantification of *cbbL*/*cbbM* marker genes encoding for RubisCO by quantitative polymerase chain reaction (qPCR) and by acetogenic and methanogenic microorganisms, shown present in the mofettes by previous studies. Combined $\Delta^{14}\text{C}$ and $\delta^{13}\text{C}$ isotope mass balances indicated that microbially derived carbon accounted for 8–27 % of bulk SOM in this soil layer.

The findings imply that autotrophic microorganisms can recycle significant amounts of carbon in wetland soils and might contribute to observed radiocarbon reservoir effects influencing $\Delta^{14}\text{C}$ signatures in peat deposits.

1 Introduction

Microbial assimilation of CO₂ is a ubiquitous process in soils, and can be accomplished by a wide variety of microorganisms using different metabolic pathways (Berg, 2011; Wood et al., 1941). RubisCO, the most important carboxy-

lating enzyme for obligate and facultative chemo- or photoautotrophic microorganisms that fix CO₂ using the Calvin–Benson–Bassham (CBB) cycle has been shown to be highly abundant in agricultural, forest, and volcanic soils (Nanba et al., 2004; Tolli and King, 2005; Selesi et al., 2007). Direct uptake of CO₂ into microbial biomass (MB) and soil organic matter (SOM) by photoautotrophic and chemoautotrophic organisms has been measured in paddy rice and agricultural upland soils (Liu and Conrad, 2011; Wu et al., 2015, 2014) as well as under manipulating experimental conditions, such as H₂ amendment (Stein et al., 2005) or addition of reduced sulfur compounds (Hart et al., 2013). Autotrophic acetogenic organisms, using the Wood–Ljungdahl pathway for CO₂ fixation, are important groups in wetland and forest soils (Küsel and Drake, 1995; Ye et al., 2014). In addition, many heterotrophic soil microorganisms fix CO₂ in order to maintain their metabolic cycle by anaplerotic reactions, either to form new sugars for cell wall synthesis or to excrete organic acids for nutrient mobilization (Feisthauer et al., 2008; Miltner et al., 2005; Santruckova et al., 2005). Global estimates of microbial CO₂ fixation in soils range between 0.9 and 5.4 PgC yr⁻¹ (Yuan et al., 2012). However, it still remains unclear how much of assimilated CO₂ is stored and contributes to the formation of soil organic matter (SOM).

Microbial utilization of CO₂ and its incorporation into SOM is also potentially an important mechanism influencing the isotope signatures of SOM (Ehleringer et al., 2000; Kramer and Gleixner, 2008). Stable carbon ($\delta^{13}\text{C}$) and radiocarbon ($\Delta^{14}\text{C}$) isotope signatures are important tools for determining turnover of soil organic matter and dating ancient sediments (Balesdent et al., 1987; Hughen et al., 2004; Trumbore, 2000).

Stable isotope variations in soil reflect mass-dependent fractionation processes (Werth and Kuzyakov, 2010). In many well-drained soils, there is a well-documented increase in $\delta^{13}\text{C}$ with depth that has been variously attributed to selective preservation/decomposition of different components of organic matter, recent declines in atmospheric $\delta^{13}\text{C}$ due to the Suess effect, or microbial fractionation (summarized in Ehleringer et al., 2000). Enzymatic fractionation during assimilation of CO₂ can also lead to changes in $\delta^{13}\text{C}$ values of synthesized organic matter (Hayes, 2001; Robinson and Cavanaugh, 1995; Whiticar, 1999). Carboxylation processes by heterotrophic microorganisms have been hypothesized to be responsible for the increase in $\delta^{13}\text{C}$ values with depth in aerated upland soils (Ehleringer et al., 2000).

Radiocarbon signatures reflect the time elapsed since the C being measured was fixed from the atmosphere, and are corrected (using measured $\delta^{13}\text{C}$ values) to remove mass-dependent fractionation effects. The radiocarbon signature of CO₂ in soil pore space can be depleted or enriched in ¹⁴C compared to organic matter found at the same depth, depending on the age of C being mineralized (Trumbore, 2006). Because soil pore space CO₂ can have quite different isotopic signatures compared to SOM at the same depth, microbial

assimilation of CO₂ may influence SOM ¹⁴C signatures and therefore bias estimates of carbon turnover and radiocarbon age by generating reservoir effects (Pancost et al., 2000).

In turn, comparing both, radiocarbon and stable isotope values of SOM, MB and their sources might allow for quantifying the potential contribution of autotrophic microorganisms to SOM, because a mismatch of both isotopes in quantifying SOM sources indicates either fractionation of ¹³C by carboxylation processes of different enzymes or depletion or enrichment of ¹⁴C by the use of soil CO₂ (Kramer and Gleixner, 2006).

In order to test the hypothesis that microbial CO₂ fixation contributes to SOM formation and alters isotope signatures in soil depth profiles, we investigated wetland mofettes in northwest Bohemia. Mofettes are cold exhalations of geogenic CO₂ from wetland soils with high CO₂ concentrations. The exhaling volcanic-derived CO₂ has a distinct isotopic signature, is enriched in $\delta^{13}\text{C}$ by about 5 ‰ and free of radiocarbon compared to atmospheric CO₂. This unique feature allows us to use geogenic CO₂ as a natural isotopic tracer, because CO₂ assimilating microorganisms take up an isotopically different CO₂ source compared to plants growing in the area, which use a mixture of geogenic and atmospheric CO₂. We used three approaches to evaluate the importance of CO₂ fixation for SOM generation in mofettes and its impact on carbon isotope values:

1. We measured natural abundance ¹³C and radiocarbon signatures of SOM, CO₂, and plant material in mofette and reference soils, in order to identify areas where C derived from microbial CO₂ fixation altered isotope signatures of bulk SOM from expected plant signals and quantified C derived from microbial CO₂ fixation by isotope mass balances.
2. We conducted isotope-labelling experiments with ¹³CO₂ in order to quantify the rate of CO₂ fixation by microorganisms in soil profiles of two CO₂ vents and compared these to reference soils away from the vents.
3. We complemented existing data about microbial community and activity in wetland mofettes (Beulig et al., 2014), by assessing the importance of microorganisms using the CBB cycle for CO₂ fixation. This was especially important to infer whether differences in kinetic isotope effects compared to plants were feasible given the pathways of microbial C fixation. Therefore, we quantified *cbbL* and *cbbM* marker genes encoding for form I and II RubisCO, respectively. Form I RubisCO consists of eight small and eight large subunits. It can be subdivided into two groups, the *red* and *green* like groups, which can be further subdivided into form 1A and 1B, as well as 1C and 1D, respectively (Yuan et al., 2012; Tolli and King, 2005). Form II RubisCO consists only of large subunits. Because of its low CO₂ affinity and high O₂ sensitivity, it represents

an early form, evolved under anaerobic conditions and high CO₂ concentrations (Alfreider et al., 2003). Form II RubisCO might be favorable under conditions prevailing in mofettes. cbbL 1A was identified mainly in obligate autotrophic bacteria and cbbL 1C in facultative autotrophic bacteria (Tolli and King, 2005). cbbM encodes for autotrophic organisms living under anaerobic conditions (Selesi et al., 2005).

Using this information, we aimed to quantify the amount of C derived from microbial assimilation of CO₂ into soil organic matter within soil profiles, and assess its potential to alter isotope signatures of SOM.

2 Materials and methods

2.1 Site description

The study site (50°08′48″ N, 12°27′03″ E) is located in the northwestern part of the Czech Republic (Bohemia). The area is part of a continental rift system, where deep tectonic faults provide pathways for ascending gases and fluids from upper earth's mantle (Kämpf et al., 2013). Mofettes are surficial, low-temperature exhalations of mantle-derived CO₂. Macroscopically, they form a complex of landscape features. At the center is a spot of typically 0.5–1 m bare soil. From this central spot, almost pure CO₂ emanates to the atmosphere. The mofette center is surrounded by a raised hummock that extends 1–20 m away from the spot. The investigated mofettes are situated on the floodplain of the river Plesná and are part of a wetland. Geogenic CO₂ emanates with an average discharge of up to 0.62 tons CO₂ d⁻¹ per spot (Kämpf et al., 2013). The surrounding hummock is built up by different vascular plant communities. *Eriophorum vaginatum* and *Deschampsia cespitosa* are dominating plant species in the immediate proximity of the central vent and hummock structure, respectively. *Filipendula ulmaria* represents typical floodplain vegetation.

We investigated two mofettes that differed in size. Mofette 1 had a spot diameter of 0.6 m, whereas the diameter of mofette 2 was 1.5 m. We also sampled soils away from the influence of the mofette-exhaled CO₂ (deemed reference soils). These soils are vegetated and experience periodic anoxic conditions due to waterlogging, as evidenced by gleyed soil features and porewater geochemistry (Mehlhorn et al., 2014). In mofettes 1 and 2, the local water table is elevated by ascending CO₂ and O₂ is mainly displaced by the CO₂ stream, leading to anoxic (but not necessarily water logged) conditions (Bräuer et al., 2011). According to the World Reference Base for soil resources (WRB, 2007), mofette soils are characterized as Histosols with pronounced reductomorphic features (reduced Y horizons) due to the influence of up-streaming CO₂. Reference soils are classified as gleyic Fluvisols (Beulig et al., 2014).

2.2 Sampling of soils, plants and gases for bulk geochemical and isotope measurements

For bulk δ¹³C and radiocarbon analyses, soil cores were taken from the central, un-vegetated part of the mofette structure and reference soils. Reference soils lacking CO₂ emissions were identified with a portable landfill gas analyzer (Visalla GM70 portable CO₂ sensor) in close proximity to each vent structure. Reference soils 1 and 2 were defined at 5 and 18 m distant from the central vent structures, respectively. Samples for bulk stable isotope and radiocarbon analyses were taken in November 2013. In order to account for soil heterogeneity, three soil cores (ID=5 cm) were taken from a plot of 50 cm × 50 cm from mofette and reference soils. Because mofette and reference soils were characterized by very different soil features, soil cores were not divided according to horizons, but depth intervals. Based on visual inspection, soil cores were divided into depth intervals of 0–10, 10–25, and 25–40 cm. Replicates of the respective depth intervals were mixed and sieved to 2 mm. Roots and plant debris were removed by hand. The sieved soil was subsequently dried at 40° and prepared for stable isotope, radiocarbon, and C / N analysis.

In April 2014, vegetation samples were taken from the same plot as soil cores, in order to characterize the isotopic composition of the plant material contributing to mofette SOM. Vegetation samples in the direct proximity of both mofettes were represented by *Eriophorum vaginatum*. Vegetation samples were also taken along a transect that crossed mofette 2, allowing us to test how the isotope signatures (δ¹³C and Δ¹⁴C) of plants changed with different mixtures of ambient and geogenic CO₂. Mofette 2 is an exposed hummock, dominated by an un-vegetated central region of CO₂ exhalation. At 1 to 2 m distant from the central exhalation, the dominant plant species was *Deschampsia cespitosa*, and at greater distances the dominant plant was *Filipendula ulmaria*. All vegetation samples were taken by clipping plants at 2 cm height and in 2 m intervals along the transect. The collected samples were dried at 40 °C, ground, and prepared for stable isotope, radiocarbon, and C / N analysis.

CO₂ was sampled from the center of each mofette by filling 250 mL evacuated stainless steel cylinders through a perforated lance from four different soil depths (5, 15, 25, 40 cm), in order to determine its radiocarbon and stable isotope signature.

2.3 Soil sampling for ¹³CO₂-labelling experiments

Mofette soils were sampled for two labelling experiments in November 2013 and September 2014. For the first experiment, 10 cm × 10 cm soil monoliths, extending to 10 cm depth were sampled from each soil in November 2013. After removing the top of the O horizon (about 1 cm thickness), the remaining material was divided into three subsamples. Each replicate was homogenized within a sterilized plastic

bag, put under an anoxic N₂ atmosphere and cooled at 4°C until further processing in the lab within the same day.

For a second experiment, three soil cores (ID = 5 cm) were taken from 0 to 40 cm of each mofette and reference soil and subsampled at 0–5, 5–10, 10–20, 20–30, and 30–40 cm. 5 g subsamples from each core were transferred immediately after core recovery to a sterilized 12 mL Labco® Exetainer, flushed with N₂ to preserve anoxia, sealed, and brought to the laboratory at 4°C for further processing. To obtain background (i.e., with no influence of added label) values for isotopic composition, one set of subsamples was dried and prepared for total organic carbon (TOC), C/N, pH, and δ¹³C analyses as described above.

2.4 Sampling for DNA extraction

Samples for DNA extraction were taken in May 2014 from mofette 1 and reference 1. Samples were taken at 0–5, 5–10, 10–20, 20–30, and 30–40 cm. Three replicates of 30 g were sampled from each depth, and homogenized under anoxic conditions. Subsequently, subsamples of 5 g were transferred to 50 mL tubes, cooled with dry ice and transported under an Ar atmosphere to the laboratory for molecular analyses.

2.5 Analyses of geochemical parameters and natural abundance isotope signatures of vegetation and soil samples

Soil pH was determined in a 0.01 M CaCl₂ solution with a soil:solution ratio of 1:2.5 using a WTW pH meter (pH 330, Weilheim, Germany). The precision of pH measurements was better than 0.1 (*n* = 3). Total C and N concentration of soil and plant samples were determined on a Vario EL (Elementar Analysensysteme GmbH, Germany). Gravimetric water content was determined after drying the soils for 48 h at 105 °C and C and N content are reported per g dry soil weight.

Stable C isotope signatures of bulk soil and plant samples were determined on an isotope ratio mass spectrometer (IRMS; DELTA+XL, Finnigan MAT, Bremen, Germany) coupled to an elemental analyzer (NA 1110, CE Instruments, Milan, Italy) via a modified ConFloII™ interface. Stable carbon isotope ratios are reported in the delta notation that expresses ¹³C/¹²C ratios as δ¹³C values in per mil (‰) relative to the international reference material Vienna Pee Dee Belemnite (V-PDB; Coplen et al., 2006):

$$\delta^{13}\text{C} = \left[\frac{\left(\frac{^{13}\text{C}}{^{12}\text{C}}_{\text{sample}} \right)}{\left(\frac{^{13}\text{C}}{^{12}\text{C}}_{\text{reference}} \right)} - 1 \right] \times 1000. \quad (1)$$

Analytical precision of all samples was better than 0.1 ‰.

For discussing microbially mediated isotope effects the isotope discrimination value Δ is used, which expresses the isotopic difference between two compounds in ‰:

$$\Delta_{x-y} = \delta_x - \delta_y. \quad (2)$$

Where δ_x and δ_y refer to δ¹³C values of the product and reactant, respectively.

The radiocarbon content of soil and plant samples was determined by accelerator mass spectrometry at the Jena ¹⁴C facilities (Steinhof et al., 2004). Subsamples of soil containing 1 mg of carbon were combusted quantitatively and the developed CO₂ was catalytically reduced to graphite at 625 °C by H₂ reduction. To simplify comparison with stable isotope ratios, radiocarbon activities are reported in Δ¹⁴C, which is the ‰ deviation of the ¹²C/¹⁴C ratio from the international oxalic acid universal standard. The Δ¹⁴C value of the sample is corrected for mass-dependent isotope fractionation to a common value of −25 ‰ (Mook and van der Plicht, 1999). The standard is corrected for radioactive decay between 1950 and the year (y) of the measurement (2014).

$$\Delta^{14}\text{C} = \left[\frac{\frac{^{14}\text{C}}{^{12}\text{C}}_{\text{sample}, -25}}{0.95 \frac{^{14}\text{C}}{^{12}\text{C}}_{\text{Ox1}, -19} \times \exp\left(y - \frac{1950}{8267}\right)} \right] \times 1000 \quad (3)$$

Errors reported for radiocarbon measurements represent the analytical error of homogenized mixed samples in ‰. Analytical precision of all radiocarbon measurements was better than 3 ‰.

2.6 Labelling experiments

The first labelling experiment traced the flow of fixed CO₂ directly into MB, evaluated rates of CO₂ uptake associated with biological activity, and compared the proportion of labelled MB in mofettes with reference soils. From each field replicate sample, 20 g aliquots were taken and put into sterilized 120 mL borosilicate bottles with butyl rubber stoppers inside a glove box containing an N₂ atmosphere. From these subsamples, three replicates were prepared for incubation with ¹³CO₂. In order to obtain control samples without biological activity, an additional aliquot of each sample was prepared and autoclaved for 2 h at 160 °C and 60 bar.

Soil samples were incubated under anoxic conditions with ¹³CO₂ at N₂:CO₂ ratios equivalent to those experienced by the soils in the field: mofette soils were incubated with a 100 vol. % ¹³CO₂ atmosphere using sterile techniques and reference soils were incubated with a 10 vol. % ¹³CO₂ and 90 vol. % N₂ atmosphere. In order to account for soil respiration and to maintain a constant label, the headspace of every sample was removed and renewed every 3 days. The samples were incubated for 14 days in the dark at 12 °C. Living and autoclaved control samples were treated identically.

After 14 days, the jars were flushed with N₂ and the soil samples were homogenized and split. One part was air dried for bulk ¹³C analysis and the other part was prepared for extraction of the microbial biomass C by chloroform fumigation extraction (CFE) (Vance et al., 1987). CFE extracts microbial biomass C by lysing the cells with chloroform and releasing the products of cell lysis into a salt solution as dissolved organic carbon (DOC). In order to enhance extraction

efficiency and to minimize the losses for extracted C by microbial degradation, the protocol from Vance et al. (1987) was slightly modified (Malik et al., 2013). The concentration of dissolved microbial biomass C (MB-DOC) and its stable carbon isotope ratio were determined by a high-performance liquid chromatography system coupled to an IRMS (HPLC–IRMS) system (Scheibe et al., 2012). This method allows direct determination of concentration and carbon isotopic value of DOC in the liquid phase by coupling a LC-IsoLink system (Thermo Electron, Bremen, Germany) to a Delta+XP IRMS (Thermo Fisher Scientific, Germany). A detailed description of the apparatus and measurement procedure is given in Scheibe et al. (2012).

The amount of microbial biomass was determined by subtracting the amount of MB-DOC of un-fumigated samples from MB-DOC of fumigated samples and dividing with a proportionality factor K_c that accounts for the extraction efficiency:

$$C_{\text{mic}} = \frac{\text{DOC}_{\text{fum}} - \text{DOC}_{\text{unfum}}}{K_c} \quad (4)$$

A value of 0.45 was used for K_c according to Amha et al. (2012). The isotope ratio of microbial biomass C can be derived by applying an isotope mass balance:

$$\delta^{13}\text{C}_{\text{MB}} = \frac{(\delta^{13}\text{C}_{\text{fum}} \times C_{\text{fum}} - \delta^{13}\text{C}_{\text{unfum}} \times C_{\text{unfum}})}{C_{\text{fum}} - C_{\text{unfum}}} \quad (5)$$

The net CO₂ fixation rate was calculated by determining the increase in ¹³C from the label compared to the unlabelled control, and is normalized for C content (either total soil or microbial C). The excess ¹³C can be derived from the ¹³C/¹²C ratio of the sample before and after the labelling:

$$\text{Excess C [mg]} = \frac{^{13}\text{C}_{\text{after labelling}}}{^{12}\text{C}_{\text{after labelling}}} \cdot C_{\text{sample}} [\text{mg}] - \frac{^{13}\text{C}_{\text{before labelling}}}{^{12}\text{C}_{\text{before labelling}}} \cdot C_{\text{sample}} [\text{mg}] \quad (6)$$

The ¹³C/¹²C ratio can be obtained from the measured $\delta^{13}\text{C}$ as follows:

$$\frac{^{13}\text{C}}{^{12}\text{C}}_{\text{sample}} = \left(\frac{\delta^{13}\text{C}_{\text{measured}}}{1000} + 1 \right) \cdot 0.011237, \quad (7)$$

where 0.01123 is the ¹³C/¹²C ratio of the international V-PDB standard (Craig, 1957).

A second labelling experiment was performed in order to obtain uptake rates as a function of depth for mofette and reference soils. After sampling 5 g of soil into 12 mL Labco® Exetainers as described above, mofette samples were flushed with 100 vol. % ¹³CO₂, and reference soils with 10 vol. % ¹³CO₂ and 90 vol. % N₂. Soils were incubated for 7 days in the dark at 12 °C. The headspace of all samples was exchanged after 3 days of incubation. After 7 days, vials were

opened and flushed with N₂ for 2 min and evacuated to remove any sorbed or dissolved ¹³CO₂. Soil samples were subsequently air dried at 60 °C and prepared for bulk ¹³C analysis as described above. The measured enrichment in ¹³C was used to measure uptake rates according Eq. (6).

2.7 DNA extraction and quantitative polymerase chain reaction

Total nucleic acid extractions of 0.7 g homogenized soil from mofette 1 and reference 1 were performed in triplicates according to the protocol of Lueders et al. (2004). Co-extracted organic soil compounds were removed by sequential purification with gel columns (S-400 HR; Zymo Research, Irvine, CA, USA) and silica columns (Powersoil Total RNA Kit in combination with the DNA Elution Accessory kit; MO BIO Laboratories, Carlsbad, CA, USA). Nucleic acid extraction efficiency was checked by agarose gel electrophoresis.

Copy numbers of 16S ribosomal ribonucleic acid (rRNA), *cbbL* 1A, *cbbL* 1C, and *cbbM* genes in extracted DNA were determined using quantitative polymerase chain reaction (qPCR). qPCR was performed on a Mx3000P instrument (Agilent, Santa Clara, CA, USA) using Maxima SYBR Green Mastermix (Thermo Scientific) and the primer combinations Uni-338 F-RC and Uni-907 R (16S rRNA; Weisburg et al., 1991), F-*cbbM* and R-*cbbM* (*cbbM*; Alfreider et al., 2003), F-*cbbL* and R-*cbbL* (*cbbL* 1A; Alfreider et al., 2003) as well as F-*cbbL* 1C and R-*cbbL* 1C (*cbbL* 1C; Alfreider et al., 2003) as described by Herrmann et al. (2012). Cycling conditions for 16S rRNA genes as well as *cbbL* and *cbbM* genes consisted of denaturation for 10 min at 95 °C, followed by 50 cycles with four temperature steps (1: 95 °C at 30 s; 2: 55 and 57 °C at 30 s for *cbbL* and *cbbM*/16S rRNA genes, respectively; 3: 72 °C at 45 s; 4: data acquisition at 78 °C and 15 s). Standard curves were constructed using plasmid CB54 for 16S rRNA and standard curves for *cbbL* and *cbbM* marker genes were constructed from 10 times dilution series of mixtures of plasmids containing *cbbL* and *cbbM* inserts as described in Herrmann et al. (2015). PCR inhibitors were tested by 10 times dilution series of representative samples. For the investigated samples 5 µL of DNA was taken as template for gene copy quantification of 16S rRNA, *cbbL*, and *cbbM*.

2.8 Mass-balance calculations

The unique isotopic composition of geogenic CO₂ and combined measurements of radiocarbon and stable isotopes allows identification of plant and microbial end-members for quantifying the importance of these two sources of SOM. Geogenic CO₂ ($\Delta^{14}\text{C} = -1000$ ‰, $\delta^{13}\text{C} = -2$ ‰) is quite different from atmospheric CO₂ ($\Delta^{14}\text{C} \sim +20$ ‰, $\delta^{13}\text{C} = -7$ ‰) in both isotopes. Therefore, $\Delta^{14}\text{C}$ values can be used to determine the overall fraction of geogenic CO₂ that is assimilated by plants or microorganisms in the mofette

Table 1. Geochemical soil properties of mofette and reference soils. $\delta^{13}\text{C}$ and geochemical data represent background (i.e., without addition of label) data obtained from sampling in September 2014. Radiocarbon data were obtained in November 2013. Uncertainties in geochemical and $\delta^{13}\text{C}$ data represent $\pm 1\sigma$ standard deviation ($n = 3$). Uncertainties in radiocarbon values represent analytical precision of a homogenized mixed sample.

	pH	TOC [wt %]	C/N	Water content [%]	$\delta^{13}\text{C}$	$\Delta^{14}\text{C}$
Mofette 1						
0–5	3.68	19.64 \pm 1.20	15.95	53	–26.90 \pm 0.15	–554.3 \pm 2.0
5–10	3.59	26.54 \pm 0.08	16.52	52	–27.55 \pm 0.21	
10–20	3.68	11.53 \pm 0.18	15.12	57	–26.71 \pm 0.18	–559.7 \pm 2.1
20–30	3.43	16.33 \pm 0.59	21.65	51	–26.79 \pm 0.12	
30–40	3.40	34.00 \pm 1.25	31.40	56	–27.01 \pm 0.23	–640.2 \pm 1.9
Reference 1						
0–5	4.13	25.85 \pm 1.72	14.37	69	–27.98 \pm 0.32	–117.5 \pm 2.8
5–10	4.07	12.40 \pm 0.60	14.18	49	–28.10 \pm 0.24	
10–20	4.00	3.16 \pm 0.26	14.52	42	–27.80 \pm 0.13	–236.3 \pm 2.7
20–30	3.91	3.14 \pm 0.13	12.93	31	–27.79 \pm 0.16	
30–40	3.69	2.81 \pm 0.50	15.88	30	–28.23 \pm 0.09	–280.2 \pm 2.5
Mofette 2						
0–5	3.80	8.66 \pm 0.69	8.95	52	–26.01 \pm 0.14	–648.1 \pm 1.2
5–10	3.76	5.87 \pm 1.11	8.97	53	–26.26 \pm 0.24	
10–20	3.79	11.41 \pm 0.95	9.72	50	–26.76 \pm 0.19	–618.7 \pm 1.3
20–30	3.52	28.72 \pm 1.42	19.74	56	–27.10 \pm 0.59	
30–40	–	–	–	–	–	–
Reference 2						
0–5	4.50	12.48 \pm 0.31	12.16	45	–27.91 \pm 0.12	–34.1 \pm 2.2
5–10	4.51	7.59 \pm 0.21	11.52	42	–28.85 \pm 0.21	
10–20	4.48	2.94 \pm 0.15	10.30	46	–28.11 \pm 0.05	–114.7 \pm 1.9
20–30	4.46	1.91 \pm 0.10	11.85	40	–27.82 \pm 0.30	
30–40	4.43	1.80 \pm 0.04	10.19	35	–28.23 \pm 0.06	–162.9 \pm 1.9

by using the end-members $\Delta^{14}\text{C}_{\text{geogenic CO}_2}$ and $\Delta^{14}\text{C}_{\text{air}}$. A conventional-mixing model for determining the fraction of geogenic CO₂ in SOM can be calculated according to

$$\text{geogenic CO}_2 \text{ in SOM}(\%) = \frac{\Delta^{14}\text{C}_{\text{SOM}} - \Delta^{14}\text{C}_{\text{air}}}{\Delta^{14}\text{C}_{\text{geogenic CO}_2} - \Delta^{14}\text{C}_{\text{air}}} \times 100. \quad (8)$$

This mass balance assumes that changes in $\Delta^{14}\text{C}_{\text{SOM}}$ caused by radioactive decay of ^{14}C are small compared to contributions from geogenic CO₂.

The same mass balance can be applied for calculating the fraction of geogenic CO₂ with stable isotope values. The end-members for this calculation are $\delta^{13}\text{C}$ values of plants, which grew solely on geogenic CO₂ or solely on ambient air CO₂. Plant $\delta^{13}\text{C}$ values are expected to be around 20‰ depleted in ^{13}C compared to the respective CO₂ source due to enzymatic fractionation, which has to be considered in determining the $\delta^{13}\text{C}$ end-member value.

We used the correlations between $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ of plant material to prove that enzymatic discrimination of plants is constant in the vicinity of the mofette, despite potentially fluctuating CO₂ concentrations. If $\Delta^{14}\text{C}$ and $\delta^{13}\text{C}$ values of

plants show a linear correlation, $\Delta^{14}\text{C}$ values of SOM can be used to derive $\delta^{13}\text{C}$ values that should be expected, if the organic matter is solely derived from plants according to the mixing model:

$$\delta^{13}\text{C}_{\text{model}} = \delta^{13}\text{C}_{\text{plant geo}} \cdot (\Delta^{14}\text{C}_{\text{SOM mofette}} \cdot m + t) + \delta^{13}\text{C}_{\text{plant air}} \cdot (1 - (\Delta^{14}\text{C}_{\text{SOM mofette}} \cdot m + t)), \quad (9)$$

where $\delta^{13}\text{C}_{\text{plant geo}}$ and $\delta^{13}\text{C}_{\text{plant air}}$ are the measured plant input end-members exhibiting the most depleted (i.e., highest exposure to geogenic CO₂) and most enriched (exposure to atmospheric CO₂) $\Delta^{14}\text{C}$ values, respectively. $\Delta^{14}\text{C}_{\text{SOM mofette}}$ are measured radiocarbon values at a certain depth within the mofette soil. m and t are the slope and intercept of the regression between measured $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ plant values. The model calculates the $\delta^{13}\text{C}_{\text{SOM}}$ that corresponds to measured $\Delta^{14}\text{C}_{\text{SOM}}$ values, if all SOM would be derived from plant material. Deviation from the model indicates input of C sources other than plants with distinct isotopic compositions.

2.9 Statistical analyses

Reported results (e.g., $\delta^{13}\text{C}$ values, microbial biomass), represent the mean of three independent replicates. Uncertainties reported for radiocarbon data represent analytical precision of a homogenized sample comprised of three independent soil cores. Differences of $\delta^{13}\text{C}$ in mofette and reference soils as well as between soil depth intervals were analyzed using Student's *t* test. Significant differences are reported at $p < 0.05$.

3 Results

3.1 pH, bulk TOC, and C/N

Soil pH ranges from 3.0 to 3.5 in mofette soils and is higher in reference soils (averaging 4.4), without significant trends with depth (Table 1). TOC contents are high (~ 12 – 20% C) in the surface 5 cm of both mofette and reference soils. In the reference soil, TOC decreases with depth to concentrations of 3% C below 20 cm. In contrast, TOC concentrations in both mofettes decrease below 5 cm (~ 6 – 16%) and increase subsequently to more than 30% below 20 cm.

Organic matter quality as indicated by C/N ratio also highlights differences between mofette and reference soils. High C/N ratios ranging from 25 to 30 are found below 20 cm depth in both mofettes, whereas C/N ratios decrease rapidly as low as 16.5–9 (for mofette 1 and 2, respectively) in the upper 10 cm (Table 1). In both reference soils, C/N ratios remain constant throughout the profile at 10–14 (Table 1).

3.2 Radiocarbon and stable isotope ratios of bulk SOM, plants, and CO₂

Consistent with our expectation, we found that geogenic CO₂ is free of radiocarbon (-1000%) and has an average $\delta^{13}\text{C}$ value of $-2.36 \pm 0.6\%$.

Radiocarbon concentrations of SOM in both mofettes are generally more depleted by several hundred ‰ compared to reference soils (Table 1). In reference soils, $\Delta^{14}\text{C}$ values decrease uniformly with depth from -60 and -34% in the top 10 cm to values of -280 and -163% at 40 cm depth in reference soil 1 and 2, respectively, reflecting radioactive decay (Table 1).

$\delta^{13}\text{C}_{\text{SOM}}$ in mofettes has average values of $-26.99 \pm 0.33\%$ and $-26.38 \pm 0.54\%$ in mofette 1 and 2, respectively. In both mofettes $\delta^{13}\text{C}_{\text{SOM}}$ decreases slightly (but not significantly) below 20 cm depth ($p = 0.39$ and 0.49 in mofette 1 and 2, respectively) (Table 1). Both reference soils have $\delta^{13}\text{C}_{\text{SOM}}$ of $-28.08 \pm 0.4\%$ with no distinct depth trend in reference 1 ($p = 0.96$) and a slight but not significant decrease in reference 2 ($p = 0.35$) below 20 cm. At every depth, reference soils are 1–2 ‰ depleted




Figure 1. Correlation between $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ of plants growing around the mofette structure. Dependent on exposure plants incorporate different amounts of geogenic CO₂, which complicates isotope mass-balance calculations for mofette SOM. However, both isotopes are highly correlated in sampled plant material, which allows for prediction of $\delta^{13}\text{C}$ SOM isotope values from plant $\Delta^{14}\text{C}$. Most data points measured from mofette SOM fall outside 95% confidence levels of the regression, which suggests a deviation of mofette SOM $\delta^{13}\text{C}$ values from a pure vegetation signal. Reference SOM $\delta^{13}\text{C}$ values fall mainly within the observed plant $\delta^{13}\text{C}$ values. Parameters of the regression are used to predict the $\delta^{13}\text{C}_{\text{SOM}}$ values expected in mofette soils that correspond to measured radiocarbon values, assuming that all carbon would be plant derived (Eq. 9).

in ^{13}C compared to mofette $\delta^{13}\text{C}_{\text{SOM}}$ throughout the soil profile ($p < 0.05$) (Table 1).

Carbon isotope signatures in vegetation samples surrounding the mofette range from -29.95 ± 0.16 to $-23.81 \pm 0.30\%$ in $\delta^{13}\text{C}$ and from -10.3 to -807.7% in $\Delta^{14}\text{C}$. Variations in the two isotopes are highly correlated, and plants with most positive $\delta^{13}\text{C}$ and most negative $\Delta^{14}\text{C}$ were found closest to the mofette and vice versa (Fig. 1). The linear fit to the strong ($R^2 = 0.86$) relationship between ^{13}C and ^{14}C found in vegetation material (Fig. 1) is used to determine parameters for the mixing model (Eq. 9). The intercept of the line with the y axis yields a value of -22.79% and represents the $\delta^{13}\text{C}$ end-member value of plant material, which is fully labelled with geogenic CO₂ ($\delta^{13}\text{C}_{\text{plant geo}}$, or *t* in Eq. 9). For the other end-member, $\delta^{13}\text{C}_{\text{plant air}}$, we used the $\delta^{13}\text{C}$ value of plants from the reference site that exhibited the most positive $\Delta^{14}\text{C}$ value, which yields $\delta^{13}\text{C}_{\text{plant air}}$ of -29.15% . The corresponding $\Delta^{14}\text{C}$ value, i.e., the value closest to atmospheric radiocarbon concentrations, was -10.3% (i.e., $\delta^{13}\text{C}_{\text{plant air}}$). This is less than $\Delta^{14}\text{C}$ measured in CO₂ in clean background air in the year of sampling ($\sim +20\%$) and indicates either that the reference site experiences some influence of geogenic CO₂ or the influence of local fossil fuel release in the region.




Figure 2. Depth profile of ¹⁴C and ¹³C signatures of SOM in mofette and reference soils. **(a)** Radiocarbon values in mofette soils are more depleted than reference soils, reflecting incorporation of geogenic CO₂ either by plants or by microorganisms. Error bars reflect analytical precision because only one homogenized sample was analyzed. **(b)** $\delta^{13}\text{C}$ values in both mofettes are also shifted towards geogenic CO₂, but to a smaller extent than radiocarbon values. Gray squares in $\delta^{13}\text{C}$ depth profiles show values of $\delta^{13}\text{C}$ in mofette SOM estimated using Eq. (9). Measured $\delta^{13}\text{C}$ values are more depleted than estimated values at all depths. **(c)** Estimated $\delta^{13}\text{C}$ values, assuming Eq. (9) but with $\Delta^{14}\text{C}$ values that have been corrected for radioactive decay assuming that SOM ages with depth in the same way as the reference soil. These estimated $\delta^{13}\text{C}$ values agree with measured values below 20 cm depth but remain depleted compared to what is expected from a pure plant SOM source in the top 10 cm. This suggests that the observed depletion in the top 10 cm of both mofette soils is caused by addition of ¹³C depleted microbial carbon, derived from fixed CO₂. In contrast, the mismatch between estimated and measured values below 20 cm depth in **(b)** can be explained by radioactive decay.

The slope of the relationship fit to plant samples (m in Eq. 9) is what would be expected for a linear mixture of plant material of the two end-member atmospheres (pure geogenic and pure air). Plant-derived SOM would be expected to fall within this mixing line. The majority (71 %) of reference soil values are within the 95 % confidence interval of this expected slope (Fig. 1). In reference soils, ¹⁴C declines with soil depth, while ¹³C remains nearly constant (Fig. 2a). Mofette SOM generally has lower ¹³C values than would be expected if they had the same linear relationship as plant material, and ¹⁴C signatures are all much lower than those of the reference soil (Fig. 1). Only 5 % of mofette SOM values fall within the 95 % confidence interval of the regression line.

3.3 Mass-balance calculations

Radiocarbon signatures of SOM indicate that, on average, 55–65 % of carbon accumulated in the mofette is derived from geogenic CO₂ (assuming end-members of –10 ‰ for $\Delta^{14}\text{C}$ air and –1000 ‰ for $\Delta^{14}\text{C}$ geogenic CO₂). The calculated proportion increases with depth. By doing the same mass-balance calculation with $\delta^{13}\text{C}$ values (with –22.47 ‰ as geogenic CO₂ end-member and –29.15 ‰ as reference end-member), one obtains lower proportions of 34–44 % geogenic C compared to the radiocarbon mass balance. This mismatch in quantifying the proportion of geogenic C suggests that $\delta^{13}\text{C}_{\text{SOM}}$ values differ from what we would expect if they were completely derived from plant inputs.

Table 2. Microbial biomass C and comparison of uptake rates determined during experiment 1 with CFE and bulk measurements. Uncertainties represent $\pm 1\sigma$ standard deviation ($n = 3$).

	$\delta^{13}\text{C}$ extract (after fumigation) [‰]	$\delta^{13}\text{C}$ control [‰]	Uptake rate [$\mu\text{g gdw}^{-1} \text{d}^{-1}$]	Uptake rate CFE [$\mu\text{g g MB}^{-1} \text{d}^{-1}$]	% labelled MB
Mofette 1					
CFE 0–10 cm	233.24 \pm 11.19	–25.94 \pm 0.36	0.17 \pm 0.03	287 \pm 85	0.88 \pm 0.33
Bulk 0–10 cm	–21.19 \pm 0.62	–26.28 \pm 0.10	0.77 \pm 0.23	–	–
Reference 1					
CFE 0–10 cm	182 \pm 5.44	–23.65 \pm 0.54	0.59 \pm 0.05	139 \pm 32	0.40 \pm 0.13
Bulk 0–10 cm	–12.82 \pm 0.95	–27.55 \pm 0.14	2.65 \pm 0.36	–	–
Mofette 2					
CFE 0–10 cm	124.51 \pm 10.66	–24.10 \pm 0.38	0.06 \pm 0.02	271 \pm 58	0.8 \pm 0.16
Bulk 0–10 cm	–21.37 \pm 0.99	–26.49 \pm 0.08	0.66 \pm 0.15	–	–
Reference 2					
CFE 0–10 cm	158.05 \pm 4.01	–26.46 \pm 0.21	0.25 \pm 0.09	99 \pm 36	0.20 \pm 0.10
Bulk 0–10 cm	–17.44 \pm 0.81	–27.21 \pm 0.22	0.71 \pm 0.16	–	–

Table 3. Quantification of 16S rRNA, cbbL, and cbbM marker genes. Uncertainties represent $\pm 1\sigma$ standard deviation ($n = 3$).

	Depth [cm]	16S rRNA	cbbM	cbbL 1A	cbbL 1C	cbbL 1C/16sRNA
Mofette 1	0–5	7.50E+10 \pm 1.42E+07	5.70E+08 \pm 3.21E+08	9.45E+08 \pm 4.86E+08	9.23E+09 \pm 4.55E+09	0.12 \pm 0.06
	5–10	1.65E+10 \pm 5.35E+06	2.21E+08 \pm 1.28E+08	1.40E+08 \pm 1.69E+08	1.46E+09 \pm 1.20E+09	0.11 \pm 0.04
	10–20	3.35E+09 \pm 0.51E+06	1.49E+07 \pm 8.45E+06	1.83E+07 \pm 1.22E+07	6.02E+08 \pm 1.25E+08	0.17 \pm 0.03
	20–30	5.94E+09 \pm 9.02E+05	1.62E+07 \pm 1.23E+07	1.12E+07 \pm 4.07E+06	3.98E+08 \pm 1.53E+08	0.07 \pm 0.03
	30–40	7.62E+08 \pm 9.39E+04	8.53E+05 \pm 3.02E+05	1.71E+06 \pm 5.23E+05	7.91E+07 \pm 2.18E+07	0.10 \pm 0.03
Reference 1	0–5	4.63E+10 \pm 3.01E+07	3.43E+08 \pm 3.18E+08	1.14E+09 \pm 4.74E+08	1.58E+10 \pm 7.20E+09	0.37 \pm 0.23
	5–10	2.98E+10 \pm 2.02E+07	2.01E+08 \pm 5.98E+07	2.69E+08 \pm 1.52E+08	7.78E+09 \pm 8.12E+08	0.28 \pm 0.08
	10–20	2.81E+10 \pm 4.83E+07	1.31E+08 \pm 4.73E+07	3.06E+08 \pm 1.59E+08	5.95E+09 \pm 1.50E+09	0.21 \pm 0.06
	20–30	1.24E+10 \pm 4.37E+07	9.75E+07 \pm 3.99E+07	9.11E+07 \pm 3.90E+07	2.25E+09 \pm 6.84E+08	0.18 \pm 0.03
	30–40	4.65E+09 \pm 9.61E+07	1.57E+08 \pm 9.26E+07	3.47E+07 \pm 2.20E+07	5.95E+08 \pm 1.78E+08	0.10 \pm 0.06

Equation (9) can be used to predict $\delta^{13}\text{C}$ SOM values corresponding to measured radiocarbon values, assuming that all carbon would be derived from unaltered plant material. Calculated $\delta^{13}\text{C}_{\text{SOM}}$ values are 1–2‰ more positive at all depths ($p < 0.05$) compared to observations (Fig. 2b), i.e., measured $\delta^{13}\text{C}_{\text{SOM}}$ values are depleted in ^{13}C compared to a signal that would be expected, if SOM preserved its original plant $\delta^{13}\text{C}$ signature.

3.4 Quantification of microbial CO₂ fixation activity

The analysis of bulk SOM and plant material revealed that mofette and reference soils are distinct in their radiocarbon as well as stable isotope values, indicating incorporation of geogenic CO₂ into mofette SOM either by plants or by microorganisms. Both isotopes show a bias in quantifying the amount of SOM derived from geogenic CO₂ by the same isotope mass balance, which suggests the presence of another source of carbon than plants, presumably microorganisms, that deplete $\delta^{13}\text{C}$ values. CO₂ fixing microorganisms might

be a potential source with a distinct $\delta^{13}\text{C}$ value. In order to assess the activity of CO₂ fixing microorganisms as well as their spatial distribution along the soil profile, we conducted two isotope-labelling experiments.

In the first experiment we traced $^{13}\text{CO}_2$ directly into MB within the first 10 cm of the soil profile. After incubating the soils with $^{13}\text{CO}_2$, MB within all soils showed high enrichment in ^{13}C , except in autoclaved control soils. Microbial biomass extracts of autoclaved controls had $\delta^{13}\text{C}$ values ranging between -24.10 ± 0.38 and -27.55 ± 0.14 ‰, in both fumigated and un-fumigated samples, which is close to $\delta^{13}\text{C}$ values obtained from bulk soil measurements (Table 2). This confirms that mainly biological processes mediated CO₂ incorporation. In un-sterilized samples, un-fumigated extracts showed enrichment in ^{13}C in all mofette and reference soils. The $\delta^{13}\text{C}$ of un-fumigated samples ranged from -14.29 ± 0.8 to $+80.47 \pm 9.46$ ‰ and are therefore enriched in ^{13}C compared to controls ($p < 0.05$). However, in all cases ^{13}C enrichment was higher after fumigation ($p < 0.05$). $\delta^{13}\text{C}$

values of fumigated samples ranged between 143.76 ± 3.93 and 227.04 ± 2.63 ‰.

The calculated rate of CO₂ uptake expressed per gram microbial biomass in the top 10 cm of soil (Table 2) was higher in mofettes compared to reference soils ($p < 0.05$) ranging between 287 ± 85 and $271 \pm 58 \mu\text{g}^{-1} \text{gMB}^{-1} \text{d}^{-1}$ in mofettes compared to 139 ± 32 and $99 \pm 36 \mu\text{g}^{-1} \text{gMB}^{-1} \text{d}^{-1}$ in reference soils (Table 2).

The second labelling experiment measured CO₂ fixation activity along the whole soil profile with samples taken from depth intervals between 1 and 40 cm. Tracer uptake was measured only in bulk SOM. In both soils, uptake rates decrease with depth (Fig. 3a). In the top 5 cm, uptake rates were higher in mofette soils compared to reference soils. Below 20 cm, rates decrease to values of $0.14 \pm 0.03 \mu\text{g g}_{\text{dw}}^{-1} \text{d}^{-1}$ in both mofettes and $0.09 \pm 0.02 \mu\text{g g}_{\text{dw}}^{-1} \text{d}^{-1}$ in reference soils. Normalizing the uptake rates to soil carbon content ($\mu\text{g gC}^{-1} \text{d}^{-1}$) instead of soil mass, removes the depth dependence of uptake rates in reference soils ($p < 0.05$), but not in mofette soils (Fig. 3b).

3.5 Quantification of 16S rRNA and marker genes for RubisCO

Results of 16S rRNA and RubisCO encoding marker genes are listed in Table 3. The abundance of 16S rRNA genes per gram soil is a measure of the total abundance of microorganisms in the soil (Fierer et al., 2005). Gene copy numbers per gram soil of 16S rRNA genes were more abundant in the top 5 cm of the mofette soil. They decrease with depth, in both, mofette and reference soil ($p < 0.05$), but the decrease is more rapid in the mofette. The same holds true for marker genes encoding for RubisCO. *cbbL* IC is the most abundant marker gene in both soils, whereas it is more abundant in the reference soil compared to the mofette. *cbbL* 1C is 1 order of magnitude more abundant than *cbbL* 1A and *cbbM* in both, reference and mofette soils. *cbbL*:16S rRNA ratios range between 0.07 ± 0.03 and 0.19 ± 0.04 in the mofette soil and stay fairly constant with depth ($p = 0.61$). In the reference soil, the ratio decreases slightly with depth from 0.37 ± 0.16 to 0.17 ± 0.04 , but values are consistently greater than in the mofette soil.

4 Discussion

4.1 Carbon sources in mofette soils

Low C/N ratios, as found in the top 10 cm of both mofettes, reflect microbially degraded organic matter (OM) (Rumpel and Kogel-Knabner, 2011) and C/N ratios as low as 9 (top 10 cm of mofette 2) suggest a high contribution of microbial biomass to bulk SOM (Wallander, 2003). A significant contribution of microbial biomass carbon at these depths is




Figure 3. CO₂ uptake rates along depth profiles of mofette and reference soils as determined by bulk measurements from experiment 2. In both mofettes, uptake rates are the highest in the top 10 cm and show a trend towards decreasing values at lower depths, especially below 20 cm. Uptake rates in reference soils also decrease with depth, but are nearly constant if normalized to organic carbon content. In contrast, uptake rates per organic carbon decline with depth in the mofette soils. This suggests increasing importance of autotrophic organisms with soil depth in the reference soil.

also supported by very high 16S rRNA gene copy numbers, extracted from mofette 1, which are 1 order of magnitude higher than known from other soils (Fierer et al., 2005). Also numbers of RubisCO encoding genes are 2 orders of magnitude more abundant than in agricultural soils (Selesi et al., 2007) and twice as high as in organic rich paddy rice fields (Wu et al., 2015), suggesting microbial carbon derived from CO₂ assimilation as an important carbon source. Further evidence is given by the isotope data, as mofette SOM at 0–10 cm differs from a pure plant signal. The deviation of $\delta^{13}\text{C}_{\text{SOM}}$ towards more negative values compared to plant signatures suggests that microbially derived carbon in shallower depths is fractionated against ¹³C, which provides further evidence that autotrophic microorganisms contribute significantly to mofette SOM.

Below 20 cm, increasing C contents in both mofettes are accompanied by a steep increase in C/N, which is attributed to lower proportions of microbial carbon and accumulation of undecomposed plant organic matter, as suggested from studies at other mofette sites (Rennert et al., 2011).

4.2 Quantification of SOM isotope shifts by combined $\Delta^{14}\text{C}$ and $\delta^{13}\text{C}$ mass balances

TOC, C/N ratios, and the abundance of 16S rRNA genes in mofette soils all suggest that microbial carbon might constitute a significant part of bulk SOM. The isotope mass-balance model can be used to assess the contribution of plant vs. microbial-derived carbon. The approach assumes that microbially derived carbon is distinct either in its ¹⁴C or its ¹³C isotope ratio compared to plant carbon. The isotope mass-balance model derived from Eq. (9) shows that microbial carbon that is added to SOM has to be depleted in $\delta^{13}\text{C}$ compared to plant inputs, leading to an overall negative $\delta^{13}\text{C}$ shift in bulk SOM of 1–2 ‰ compared to a pure plant signal at all depths (Fig. 2b).

However, the model assumes that the radiocarbon content of mofette SOM solely depends on the amount of fixed geogenic CO₂ and does not consider radioactive decay. ¹⁴C depletion by radioactive decay, especially with soil depth, can lead to an overestimation of the shift in $\delta^{13}\text{C}$ values and consequently to an overestimation of fixed geogenic CO₂. In order to account for ¹⁴C depletion by radioactive decay, $\Delta^{14}\text{C}$ values of reference soil SOM can be subtracted from $\Delta^{14}\text{C}_{\text{SOM}_{\text{mofette}}}$ in Eq. (9).

After correcting the model for radioactive decay, the calculated $\delta^{13}\text{C}_{\text{SOM}}$ depletion still matches the data for the first 10 cm of both mofettes, where measured $\delta^{13}\text{C}$ values are more negative than calculated ones (Fig. 2c). Below 10 cm, the calculated $\delta^{13}\text{C}_{\text{SOM}}$ coincides with measured values in both mofettes, suggesting that SOM $\delta^{13}\text{C}$ preserved the signal of the plant source and only radioactive decay lead to the initial $\delta^{13}\text{C}$ shift in the model (Fig. 2c). This supports findings from previous studies, where carbon accumulation accompanied with high C/N ratios was attributed to accumu-

lation of poorly decomposed plant material (Rennert et al., 2011). The only exception from this pattern is at 30–40 cm in mofette 2, where measured $\delta^{13}\text{C}$ values are still more negative than calculated ones, even after correction for radioactive decay (Fig. 2c). This might be caused by extremely low carbon dynamics, e.g., due to permanently waterlogged conditions, which would lead to an overestimation of the $\delta^{13}\text{C}$ isotope shift in the model. Although water levels fluctuate in the floodplain, permanently waterlogged conditions are likely to occur deeper in mofette 2, where high CO₂ discharge rates might lead to an elevation of the water table. Waterlogged conditions lead to low carbon turnover, and correction of radioactive decay with reference soil values might not be sufficient, because reference soils at these depths are only temporally waterlogged. This might explain the mismatch of measured and calculated $\delta^{13}\text{C}$ values at the deepest sampling point in mofette 2 and would indicate a potential bias of modeled C-isotope signatures towards too positive $\delta^{13}\text{C}$ values.

Another source of error in the model is accumulation of recalcitrant compounds within the SOM pool, like lignin or lipids, which might also lead to a shift in $\delta^{13}\text{C}$ values compared to the original bulk plant material (Benner et al., 1987; Werth and Kuzyakov, 2010). The accumulation of phenolic compounds is usually accompanied by an increase in C/N ratios (Hornibrook et al., 2000; Werth and Kuzyakov, 2010), which is not the case in the top 10 cm of the mofette soil. Therefore, lignin accumulation is not likely to have caused the depletion in the top 10 cm of both mofettes. Nevertheless, increased lignin accumulation might also be the reason for the observed depletion in $\delta^{13}\text{C}$ below 20 cm depth in mofette 2.

Therefore, the model shows that $\delta^{13}\text{C}$ values in the top 10 cm of both mofettes are significantly lower than expected for SOM derived from plants alone, indicating significant addition of $\delta^{13}\text{C}$ depleted carbon. Below 10 cm depth, the calculated and measured $\delta^{13}\text{C}$ values agree after correcting for possible sources of error, like radioactive decay and alteration of $\delta^{13}\text{C}$ due to decomposition processes.

Microbial carbon that is added to mofette SOM by several CO₂ fixation pathways is likely to be depleted in $\delta^{13}\text{C}$ because of enzymatic fractionation processes (Fuchs, 2011). The deviation in $\delta^{13}\text{C}$ in the top 10 cm of both mofettes also is in accord with high CO₂ fixation rates and the abundance of functional marker genes for CO₂ fixation at this depth (Fig. 4). This implies that microbial carbon derived from CO₂ assimilating organisms is a major driver of the observed $\delta^{13}\text{C}_{\text{SOM}}$ depletion.

4.3 Quantification of microbial carbon C derived from CO₂ fixation

In order to quantify the proportion of CO₂-derived microbial carbon from the observed isotope shift, it is important to know the metabolic pathway that was used for CO₂ fixation and its corresponding isotope fractionation factor. Beulig et




Figure 4. Correlation of marker genes encoding for RubisCO and measured uptake rates in mofette soil 1 and reference soil 1 in the soil depth profile from 0 to 40 cm depth. The good correlation in the reference soil indicates high contribution of autotrophic microorganisms to measured uptake rates. In the mofette soil R^2 is considerably lower, most probably, because also other CO₂ fixation cycles than the CBB cycle, like the acetyl-CoA cycle, are important pathways in these soils.

al. (2016) investigated by metatranscriptomic and metagenomic approaches microbial key processes in mofette soil 1. Consistent with our quantification of cbbL/cbbM marker genes, Beulig et al. (2016) detected high frequencies of transcripts encoding key enzymes for the CBB as well as the reductive acetyl-CoA (coenzyme A) cycle. The acetyl-CoA Cycle is used by acetogens, methanogens, and sulfate reducers for catabolism and anabolism (Drake et al., 2006). According to Beulig et al. (2016), transcripts of key enzymes for the acetyl-CoA pathway in the mofette soil are also related to these groups. Most transcripts encoding for the CBB were related to chemoautotrophic bacteria and algae, living under anaerobic restrictions. The activity of autotrophic bacteria using the CBB is also supported by our data, as shown by the good correlation of cbbL/cbbM marker genes and uptake rates (Fig. 4).

Carbon that is fixed by autotrophs or algae using form I RubisCO, the dominant form in the mofette, is depleted by -27 to -30 ‰ compared to the source CO₂ ($\Delta \approx -27$ to -30 ‰) (Hayes, 2001; Pancost and Damste, 2003). A similar value can be expected for acetate formed from geogenic CO₂ during acetogenesis. In systems where acetate is not limiting, depletion is less pronounced ($\Delta \approx -32$ ‰) than in acetate-limited systems ($\Delta \approx -58.6$ ‰) (Conrad, 2005; Gelwicks et al., 1989). A value of -32 ‰ is in accordance with acetate $\delta^{13}\text{C}$ values measured by Beulig et al. (2014) in a mofette study from the same area. Therefore, given a $\delta^{13}\text{C}$ value of geogenic CO₂ of around -2 ‰, the C end-member derived from microbial CO₂ fixation adds carbon with an average $\delta^{13}\text{C}$ value of -30 to -34 ‰ to bacterial biomass and SOM in mofettes. Taking the differences between measured and calculated $\delta^{13}\text{C}$ (with and without correction for radioactive decay, respectively) for mass-balance calculation according to Eq. (8), microbially fixed geogenic CO₂ carbon in the top

10 cm of the mofette soil can make up between 8 ± 2 and 15 ± 4 % in mofette 1 and between 23 ± 4 and 27 ± 5 % in mofette 2.

4.4 Importance of microbial CO₂ fixation for isotope ratios in peat soils

Our data provide evidence that assimilation of CO₂ by several groups of autotrophic microorganisms contribute to SOM formation. Recycling of CO₂ in peat deposits has been proposed to cause reservoir effects in radiocarbon, biasing dating of peat (Kilian et al., 1995). As an explanation, Pancost et al. (2000) proposed recycling of $\Delta^{14}\text{C}$ depleted methane that diffuses from the catotelm layer up the peat profile, where it is oxidized by methanotrophic organisms and subsequently assimilated by mycorrhizal fungi living in association with *Ericaceae* rootlets. However, the authors could not find evidence from biomarker analyses of methanotrophic or fungal organisms and attributed recycling of ^{14}C depleted CO₂ to plants. Our findings suggest that other groups besides fungi are involved in CO₂ recycling, namely, CO₂ utilizing autotrophic microorganisms. Pancost et al. (2000) estimated that 20 % of C in the investigated peat is derived from this recycling process. This proportion is very similar to our estimates for autotrophic fixation of CO₂ in the 0–10 cm of mofette soil. Hence, we would propose that direct fixation of CO₂ could be a major process influencing peat radiocarbon signatures.

4.5 Importance of CO₂ fixation for soil carbon in reference soils

When normalized for the mass of carbon (as opposed to mass of soil), rates of CO₂ fixation in the reference soil at depth remain similar to values at the surface (Fig. 3). We cannot use the isotope-mixing model to estimate the amount of C derived from CO₂ fixation in the reference soil, because the soil atmosphere as well as plants at the reference soil are not directly influenced by geogenic CO₂. However, the rate measurements suggest increasing importance of CO₂ assimilating microorganisms for carbon stocks with depth. In addition, the high relative abundance of RubisCO marker genes relative to 16S rRNA genes suggest that autotrophic organisms constitute a substantial part of the microbial community throughout the soil profile. Their activity is also indicated by the strong correlation between RubisCO marker genes and uptake rates ($R^2 = 0.94$; $p < 0.05$) (Fig. 4). Higher CO₂ concentrations, which are usually observed with depth, might also lead to an increase of CO₂ assimilation, because of higher substrate availability for RubisCO or other carboxylases with depth.

In contrast to the mofette soil, which is characterized as an organic rich Histosol, reference soils are classified as Gleysols, with high organic carbon contents only in the A horizon. They are characterized by frequently changing

redox conditions due to groundwater fluctuations, which might provide sufficient electron donors and acceptors for chemolithoautotrophic microorganisms (Akob and Küsel, 2011).

Beulig et al. (2014) characterized the microbial community of a reference soil at the same study site. The authors found that Proteobacteria constituted a substantial part of the microbial community. Many Proteobacteria are facultative autotrophs using the CBB cycle and have a facultative anaerobe metabolism (Badger and Bek, 2008). They would be therefore able to assimilate CO₂ also under the experimental conditions.

A contribution of phototrophic and chemoautotrophic microorganisms to SOM has been demonstrated already by other studies (Hart et al., 2013; Yuan et al., 2012), but solely for top soils. Wu et al. (2014, 2015) investigated soil depth profiles up to 15 cm depth, but found no significant incorporation below 5 cm depth in upland and paddy soils under not manipulating experimental conditions, like illumination.

Our data suggest that autotrophic microorganisms are active even in the reference subsoil. Microorganisms using the CBB cycle would add ¹³C-depleted carbon to SOM. Indeed, $\delta^{13}\text{C}$ profiles of both reference soils do not show shifts towards more positive values with depth, as is usually observed from other Gleysols, although radiocarbon data indicate that SOM becomes older with depth (Alewell et al., 2011; Bol et al., 1999). Further, both reference soils have C/N ratios close to 10 throughout the soil profile, which normally indicates a higher contribution of microbial C to SOM (Rumpel and Kogel-Knabner, 2011). This strongly suggests a contribution of autotrophic microorganisms to carbon stocks in the subsoil, though ultimately its influence on the C isotopic signature of SOM at depth must be further evaluated.

5 Conclusions

$\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ values of SOM in wetland mofettes are influenced by incorporation of geogenic CO₂ fixed not only by plants but also by microbes, as indicated by deviation of $\delta^{13}\text{C}$ values from those expected if plant C inputs were the sole source of SOM-C. The unique isotopic composition of geogenic CO₂ and the different enzymatic fractionation of plants and microorganisms allows us to quantify microbially derived C using combined ¹⁴C and ¹³C mass balances, because microbial carbon is more depleted than plant C. Other parameters, like C/N ratio, 16S rRNA, and cbbL gene abundance also indicate addition of C fixed from geogenic CO₂ by microbes. According to the isotope mass balances, microbial carbon derived from CO₂ fixation accounts for 8–27% of bulk SOM in mofette soils. The significant contribution of autotrophic microorganisms to SOM also implies that they might be able to cause reservoir effects in radiocarbon by recycling of old CO₂, as has been already suggested for peat soils.

Further, high CO₂ fixation rates, especially in mineral horizons of the reference soil, as well as the high of RubisCO marker genes indicate a significant contribution of autotrophic microorganisms to subsoil carbon.

Acknowledgements. We thank Heike Geilmann and Stefan Rühlow for assistance with $\delta^{13}\text{C}$ analysis of bulk soil and CFE extracts. We thank Heike Machts and Axel Steinhof for radiocarbon analysis of soil and plant samples. Further, we thank Iris Kuhlmann for assistance in CFE extractions, as well as Julia Kuhr for helping in DNA extraction and soil sampling. We kindly acknowledge Gerd Gleixner for helpful discussions and comments on the manuscript. We would also like to acknowledge two anonymous reviewers, whose comments improved the manuscript. This project was supported by the graduate research training group “Alteration and element mobility at the microbe–mineral interface” (GRK 1257), which is part of the Jena School for Microbial Communication (JSMC) and funded by the Deutsche Forschungsgemeinschaft (DFG).

The article processing charges for this open-access publication were covered by the Max Planck Society.

Edited by: I. Trebs

References

- Akob, D. M. and Küsel, K.: Where microorganisms meet rocks in the Earth’s Critical Zone, *Biogeosciences*, 8, 3531–3543, doi:10.5194/bg-8-3531-2011, 2011.
- Alewell, C., Giesler, R., Klaminder, J., Leifeld, J., and Rollog, M.: Stable carbon isotopes as indicators for environmental change in peatlands, *Biogeosciences*, 8, 1769–1778, doi:10.5194/bg-8-1769-2011, 2011.
- Alfreider, A., Vogt, C., Hoffmann, D., and Babel, W.: Diversity of ribulose-1,5-bisphosphate carboxylase/oxygenase large-subunit genes from groundwater and aquifer microorganisms, *Microb. Ecol.*, 45, 317–328, 2003.
- Amha, Y., Bohne, H., and Alsanius, B.: Comparison of physiological and biochemical methods for assessing microbial activity and biomass of peats, in: *Peat: Formation, Uses and Biological Effects*, 35–56, 2012.
- Badger, M. R. and Bek, E. J.: Multiple Rubisco forms in proteobacteria: their functional significance in relation to CO₂ acquisition by the CBB cycle, *J. Exp. Bot.*, 59, 1525–1541, 2008.
- Balesdent, J., Mariotti, A., and Guillet, B.: Natural C-13 abundance as a tracer for studies of soil organic-matter dynamics, *Soil Biol. Biochem.*, 19, 25–30, 1987.
- Benner, R., Fogel, M. L., Sprague, E. K., and Hodson, R. E.: Depletion of C-13 in Lignin and Its Implications for Stable Carbon Isotope Studies, *Nature*, 329, 708–710, 1987.
- Berg, I. A.: Ecological aspects of the distribution of different autotrophic CO₂ fixation pathways, *Appl. Environ. Microb.*, 77, 1925–1936, 2011.
- Beulig, F., Heuer, V. B., Akob, D. M., Viehweger, B., Elvert, M., Herrmann, M., Hinrichs, K.-U., and Küsel, K.: Carbon flow

- from volcanic CO₂ into soil microbial communities of a wetland mofette, *ISME J.*, 9, 746–759, doi:10.1038/ismej.2014.148, 2014.
- Beulig, F., Urich, T., Nowak, M., Gleixner, G., Trumbore, S. E., and Küsel, K.: Soil carbon accumulation under high levels of CO₂ is associated with reduced foodweb complexity, *Nature Microbiology*, in press, 2016.
- Bol, R. A., Harkness, D. D., Huang, Y., and Howard, D. M.: The influence of soil processes on carbon isotope distribution and turnover in the British uplands, *Eur. J. Soil Sci.*, 50, 41–51, 1999.
- Bräuer, K., Kämpf, H., Koch, U., and Strauch, G.: Monthly monitoring of gas and isotope compositions in the free gas phase at degassing locations close to the Novy Kostel focal zone in the western Eger Rift Czech Republic, *Chem. Geol.*, 290, 163–176, 2011.
- Conrad, R.: Quantification of methanogenic pathways using stable carbon isotopic signatures: a review and a proposal, *Org. Geochem.*, 36, 739–752, 2005.
- Craig, H.: Isotopic standards for carbon and oxygen and correction factors for mass-spectrometric analysis of carbon dioxide, *Geochim. Cosmochim. Ac.*, 12, 133–149, 1957.
- Drake, H., Küsel, K., and Matthies, C.: Acetogenic Prokaryotes, in: *The Prokaryotes*, edited by: Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.-H., and Stackebrandt, E., Springer New York, 2006.
- Ehleringer, J. R., Buchmann, N., and Flanagan, L. B.: Carbon isotope ratios in belowground carbon cycle processes, *Ecol. Appl.*, 10, 412–422, 2000.
- Feisthauer, S., Wick, L. Y., Kastner, M., Kaschabek, S. R., Schloemann, M., and Richnow, H. H.: Differences of heterotrophic (CO₂)-C-13 assimilation by *Pseudomonas knackmussii* strain B13 and *Rhodococcus opacus* ICP and potential impact on biomarker stable isotope probing, *Environ. Microbiol.*, 10, 1641–1651, 2008.
- Fierer, N., Jackson, J. A., Vilgalys, R., and Jackson, R. B.: Assessment of soil microbial community structure by use of taxon-specific quantitative PCR assays, *Appl. Environ. Microb.*, 71, 4117–4120, 2005.
- Fuchs, G.: Alternative pathways of carbon dioxide fixation: insights into the early evolution of life?, in: *Annual Review of Microbiology*, vol. 65, edited by: Gottesman, S. and Harwood, C. S., 631–658, 2011.
- Gelwicks, J. T., Risatti, J. B., and Hayes, J. M.: Carbon isotope effects associated with autotrophic acetogenesis, *Org. Geochem.*, 14, 441–446, 1989.
- Hart, K. M., Kulakova, A. N., Allen, C. C. R., Simpson, A. J., Openheimer, S. F., Masoom, H., Courtier-Murias, D., Soong, R., Kulakov, L. A., Flanagan, P. V., Murphy, B. T., and Kelleher, B. P.: Tracking the fate of microbially sequestered carbon dioxide in soil organic matter, *Environ. Sci. Technol.*, 47, 5128–5137, 2013.
- Hayes, J. M.: Fractionation of carbon and hydrogen isotopes in biosynthetic processes, *Rev. Mineral. Geochem.*, 43, 225–277, 2001.
- Herrmann, M., Hädrich, A., and Küsel, K.: Predominance of thaumarchaeal ammonia oxidizer abundance and transcriptional activity in an acidic fen, *Environ. Microbiol.*, 14, 3013–3025, 2012.
- Hornibrook, E. R. C., Longstaffe, F. J., Fyfe, W. S., and Bloom, Y.: Carbon-isotope ratios and carbon, nitrogen and sulfur abundances in flora and soil organic matter from a temperate-zone bog and marsh, *Geochem. J.*, 34, 237–245, 2000.
- Hughen, K., Lehman, S., Southon, J., Overpeck, J., Marchal, O., Herring, C., and Turnbull, J.: C-14 activity and global carbon cycle changes over the past 50,000 years, *Science*, 303, 202–207, 2004.
- Kämpf, H., Bräuer, K., Schumann, J., Hahne, K., and Strauch, G.: CO₂ discharge in an active, non-volcanic continental rift area (Czech Republic): characterisation ($\delta^{13}\text{C}$, $^3\text{He}/^4\text{He}$) and quantification of diffuse and vent CO₂ emissions, *Chem. Geol.*, 339, 71–83, 2013.
- Kilian, M. R., Van Der Plicht, J., and Van Geel, B.: Dating raised bogs: new aspects of AMS C-14 wiggle matching, a reservoir effect and climatic change, *Quaternary Sci. Rev.*, 14, 959–966, 1995.
- Kramer, C. and Gleixner, G.: Soil organic matter in soil depth profiles: distinct carbon preferences of microbial groups during carbon transformation, *Soil Biol. Biochem.*, 40, 425–433, 2008.
- Küsel, K. and Drake, H. L.: Effects of environmental parameters on the formation and turnover of acetate by forest soils, *Appl. Environ. Microb.*, 61, 3667–3675, 1995.
- Liu, F. H. and Conrad, R.: Chemolithotrophic acetogenic H₂/CO₂ utilization in Italian rice field soil, *ISME J.*, 5, 1526–1539, 2011.
- Lueders, T., Manefield, M., and Friedrich, M. W.: Enhanced sensitivity of DNA- and rRNA-based stable isotope probing by fractionation and quantitative analysis of isopycnic centrifugation gradients, *Environ. Microbiol.*, 6, 73–78, 2004.
- Malik, A., Blagodatskaya, E., and Gleixner, G.: Soil microbial carbon turnover decreases with increasing molecular size, *Soil Biol. Biochem.*, 62, 115–118, 2013.
- Mehlhorn, J., Beulig, F., Küsel, K., and Planer-Friedrich, B.: Carbon dioxide triggered metal(loid) mobilisation in a mofette, *Chem. Geol.*, 382, 54–66, 2014.
- Miltner, A., Kopinke, F. D., Kindler, R., Selesi, D. E., Hartmann, A., and Kastner, M.: Non-phototrophic CO₂ fixation by soil microorganisms, *Plant Soil*, 269, 193–203, 2005.
- Mook, W. G. and van der Plicht, J.: Reporting C-14 activities and concentrations, *Radiocarbon*, 41, 227–239, 1999.
- Namba, K., King, G. M., and Dunfield, K.: Analysis of facultative lithotroph distribution and diversity on volcanic deposits by use of the large subunit of ribulose 1,5-bisphosphate carboxylase/oxygenase, *Appl. Environ. Microb.*, 70, 2245–2253, 2004.
- Pancost, R. D. and Damste, J. S. S.: Carbon isotopic compositions of prokaryotic lipids as tracers of carbon cycling in diverse settings, *Chem. Geol.*, 195, 29–58, 2003.
- Pancost, R. D., van Geel, B., Baas, M., and Damste, J. S. S.: delta C-13 values and radiocarbon dates of microbial biomarkers as tracers for carbon recycling in peat deposits, *Geology*, 28, 663–666, 2000.
- Rennert, T., Eusterhues, K., Pfanz, H., and Totsche, K. U.: Influence of geogenic CO₂ on mineral and organic soil constituents on a mofette site in the NW Czech Republic, *Eur. J. Soil Sci.*, 62, 572–580, 2011.
- Robinson, J. J. and Cavanaugh, C. M.: Expression of Form I and Form II Rubisco in chemoautotrophic symbioses: implications for the interpretation of stable carbon isotope values, *Limnol. Oceanogr.*, 40, 1496–1502, 1995.

- Rumpel, C. and Kogel-Knabner, I.: Deep soil organic matter—a key but poorly understood component of terrestrial C cycle, *Plant Soil*, 338, 143–158, 2011.
- Santruckova, H., Bird, M. I., Elhottova, D., Novak, J., Picek, T., Simek, M., and Tykva, R.: Heterotrophic fixation of CO₂ in soil, *Microb. Ecol.*, 49, 218–225, 2005.
- Scheibe, A., Krantz, L., and Gleixner, G.: Simultaneous determination of the quantity and isotopic signature of dissolved organic matter from soil water using high-performance liquid chromatography/isotope ratio mass spectrometry, *Rapid Commun. Mass Sp.*, 26, 173–180, 2012.
- Selesi, D., Schmid, M., and Hartmann, A.: Diversity of green-like and red-like ribulose-1,5-bisphosphate carboxylase/oxygenase large-subunit genes (cbbL) in differently managed agricultural soils, *Appl. Environ. Microb.*, 71, 175–184, 2005.
- Selesi, D., Pattis, I., Schmid, M., Kandeler, E., and Hartmann, A.: Quantification of bacterial RubisCO genes in soils by cbbL targeted real-time PCR, *J. Microbiol. Meth.*, 69, 497–503, 2007.
- Stein, S., Selesi, D., Schilling, R., Pattis, I., Schmid, M., and Hartmann, A.: Microbial activity and bacterial composition of H₂-treated soils with net CO₂ fixation, *Soil Biol. Biochem.*, 37, 1938–1945, 2005.
- Steinhof, A., Adamiec, G., Gleixner, G., van Klinken, G. J., and Wagner, T.: The new C-14 analysis laboratory in Jena, Germany, *Radiocarbon*, 46, 51–58, 2004.
- Tolli, J. and King, G. M.: Diversity and structure of bacterial chemolithotrophic communities in pine forest and agroecosystem soils, *Appl. Environ. Microb.*, 71, 8411–8418, 2005.
- Trumbore, S. E.: Age of soil organic matter and soil respiration: Radiocarbon constraints on belowground C dynamics, *Ecol. Appl.*, 10, 399–411, 2000.
- Trumbore, S. E.: Carbon respired by terrestrial ecosystems – recent progress and challenges, *Glob. Change Biol.*, 12, 141–153, 2006.
- Vance, E. D., Brookes, P. C., and Jenkinson, D. S.: An extraction method for measuring soil microbial biomass-C, *Soil Biol. Biochem.*, 19, 703–707, 1987.
- Wallander, H., Nilsson, L. O., Hagerberg, D., and Rosengren, U.: Direct estimates of C:N ratios of ectomycorrhizal mycelia collected from Norway spruce forest soils, *Soil Biol. Biochem.*, 35, 997–999, 2003.
- Weisburg, W. G., Barns, S. M., Pelletier, D. A., and Lane, D. J.: 16S ribosomal DNA amplification for phylogenetic study, *J. Bacteriol.*, 173, 697–703, 1991.
- Werth, M. and Kuzyakov, Y.: C-13 fractionation at the root–microorganisms–soil interface: a review and outlook for partitioning studies, *Soil Biol. Biochem.*, 42, 1372–1384, 2010.
- Whiticar, M. J.: Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane, *Chem. Geol.*, 161, 291–314, 1999.
- Wood, H. G., Werkman, C. H., Hemingway, A., and Nier, A. O.: Heavy carbon as a tracer in heterotrophic carbon dioxide assimilation, *J. Biol. Chem.*, 139, 365–376, 1941.
- WRB: World Reference Base for Soil Resources 2006, World Soil Resources Reports No. 103., FAO, Rome, 2007.
- Wu, X. H., Ge, T. D., Yuan, H. Z., Li, B. Z., Zhu, H. H., Zhou, P., Sui, F. G., O'Donnell, A. G., and Wu, J. S.: Changes in bacterial CO₂ fixation with depth in agricultural soils, *Appl. Microbiol. Biot.*, 98, 2309–2319, 2014.
- Wu, X. H., Ge, T. D., Wang, W., Yuan, H. Z., Wegner, C. E., Zhu, Z. K., Whiteley, A. S., and Wu, J. S.: Cropping systems modulate the rate and magnitude of soil microbial autotrophic CO₂ fixation in soil, *Front. Microbiol.*, 6, 379, doi:10.3389/fmicb.2015.00379, 2015.
- Ye, R. Z., Jin, Q. S., Bohannan, B., Keller, J. K., and Bridgham, S. D.: Homoacetogenesis: a potentially underappreciated carbon pathway in peatlands, *Soil Biol. Biochem.*, 68, 385–391, 2014.
- Yuan, H. Z., Ge, T. D., Chen, C. Y., O'Donnell, A. G., and Wu, J. S.: Significant role for microbial autotrophy in the sequestration of soil carbon, *Appl. Environ. Microb.*, 78, 2328–2336, 2012.