



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Genome-wide association analyses identify 18 new loci associated with serum urate concentrations**

**Citation for published version:**

LifeLines Cohort Study, Köttgen, A, Albrecht, E, Teumer, A, Vitart, V, Krumsiek, J, Hundertmark, C, Pistis, G, Ruggiero, D, O'Seaghdha, CM, Haller, T, Yang, Q, Tanaka, T, Johnson, AD, Kutalik, Z, Smith, AV, Shi, J, Struchalin, M, Middelberg, RPS, Brown, MJ, Gaffo, AL, Pirastu, N, Li, G, Hayward, C, Zemunik, T, Huffman, J, Yengo, L, Zhao, JH, Demirkan, A, Feitosa, MF, Liu, X, Malerba, G, Lopez, LM, van der Harst, P, Li, X, Kleber, ME, Wild, SH, Tenesa, A, Navarro, P, Hastie, N, Davies, G, Gow, AJ, Wilson, JF, Farrington, SM, Theodoratou, E, Polasek, O, Campbell, H, Rudan, I, Deary, IJ, Wright, AF & Dunlop, MG 2013, 'Genome-wide association analyses identify 18 new loci associated with serum urate concentrations', *Nature Genetics*, vol. 45, no. 2, pp. 145-154. <https://doi.org/10.1038/ng.2500>

**Digital Object Identifier (DOI):**

[10.1038/ng.2500](https://doi.org/10.1038/ng.2500)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Nature Genetics

**Publisher Rights Statement:**

Published in final edited form as:  
Nat Genet. 2013 February ; 45(2): 145–154. doi:10.1038/ng.2500.

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



Published in final edited form as:

*Nat Genet.* 2013 February ; 45(2): 145–154. doi:10.1038/ng.2500.

## Genome-wide association analyses identify 18 new loci associated with serum urate concentrations

Anna Köttgen<sup>1,2,156</sup>, Eva Albrecht<sup>3,156</sup>, Alexander Teumer<sup>4,156</sup>, Veronique Vitart<sup>5,156</sup>, Jan Krumsiek<sup>6,156</sup>, Claudia Hundertmark<sup>1</sup>, Giorgio Pistis<sup>7</sup>, Daniela Ruggiero<sup>8</sup>, Conall M O'Seaghda<sup>9,10,11</sup>, Toomas Haller<sup>12</sup>, Qiong Yang<sup>9,10,13</sup>, Toshiko Tanaka<sup>14</sup>, Andrew D Johnson<sup>9,10</sup>, Zoltán Kutalik<sup>15,16</sup>, Albert V Smith<sup>17,18</sup>, Julia Shi<sup>19</sup>, Maksim Struchalin<sup>20</sup>, Rita P S Middelberg<sup>21,22</sup>, Morris J Brown<sup>23</sup>, Angelo L Gaffo<sup>24,25</sup>, Nicola Pirastu<sup>26</sup>, Guo Li<sup>27</sup>, Caroline Hayward<sup>5</sup>, Tatijana Zemunik<sup>28</sup>, Jennifer Huffman<sup>5</sup>, Loic Yengo<sup>29,30</sup>, Jing Hua Zhao<sup>31</sup>, Ayse Demirkan<sup>32</sup>, Mary F Feitosa<sup>33</sup>, Xuan Liu<sup>13</sup>, Giovanni Malerba<sup>34</sup>, Lorna M Lopez<sup>35,36</sup>, Pim van der Harst<sup>37</sup>, Xinzhong Li<sup>38</sup>, Marcus E Kleber<sup>39,40</sup>, Andrew A Hicks<sup>41</sup>,

© 2013 Nature America, Inc. All rights reserved.

Correspondence should be addressed to A. Köttgen (anna.koettgen@uniklinik-freiburg.de), V.V. (veronique.vitart@igmm.ed.ac.uk), M. Bochud (murielle.bochud@chuv.ch) or C.G. (christian.gieger@helmholtz-muenchen.de).

<sup>109</sup>A list of contributing members appears in the Supplementary Note.

<sup>156</sup>These authors contributed equally to this work.

<sup>157</sup>These authors jointly directed this work.

Note: Supplementary information is available in the online version of the paper.

### AUTHOR CONTRIBUTIONS

Study design: A. Köttgen, C.G., E.A. and M. Caulfield.

Design and/or management of the individual studies: A.A.H., A. Tenesa, A.F.W., A.L., B.M.P., C.G., D.I.C., D.R., E.G.H., E.O., E. Trabetti, G.C., G.P., H. Campbell, H.-E.W., H. Snieder, I.J.D., J.A., J.C., J.F.W., J.V., L.M.R., M. Ciullo, M. Caulfield, M.F., M. Kubo, M.L., M.V., N.H., N.J.S., N. Kamatani, O.M.W., O.P., O.R., P.B.M., P.D., P.G., P.K., P. Mudgal, P.M.R., P.P.P., P.V., R.M.P., R. Sorice, S.H.W., S.M.F., S.U., T.E., T.L., Toshihiro Tanaka, V.S., W.H.L.K., Y.N., Y.O. and Z.K.

Phenotype collection: A.A.H., A.J.G., A.v.E., B.M.P., E.O., G.C., G.G., G.K.G., G.W., H. Snieder, I.J.D., I.K., I. Persico, J.C., J.F.W., J.V., L. Frogner, M. Ciullo, M. Caulfield, M.G.D., M.G.P., M.J.B., M. Kähönen, M. Kubo, M.L., M. Pirastu, M.V., N.J.S., O.D., O.M.W., O.P., P. Sharma, P.B.M., P.K., P. Mudgal, P.P.P., P.V., S. Schipf, S.H.W., S.M.F., S.T., S.U., T. Zemunik and V.S.

Genotyping: A.A.H., A. Tenesa, A. Teumer, C. Hayward, D.I.C., E.L., F.E., G.C., G.D., G.W.M., I. Persico, J.F.W., M. Ciullo, M. Caulfield, M.E.K., M.G.D., M.G.P., M. Kubo, M.L., M. Putku, M.W., N.J.S., N. Klopp, O.R., P.B.M., P.D., P.M.R., P.P.P., P.v.d.H., R.J.S., S.M.F., T.E., T.L., T. Zeller and T. Zemunik.

Statistical methods and analysis: A. Tenesa, A. Tin, A. Köttgen, A. Teumer, A. Demirkan, C. Hayward, C. Hundertmark, C.G., C. Schurmann, D.C., D.I.C., D.R., E.A., E.G.H., F.M., F.T., G.A.T., G.K.G., G.L., G.M., G.P., I.M.L., I. Prokopenko, J.H., J.K., L.M.L., L.M.R., L.P., M.A.N., M. Steri, M. Bochud, M.E.K., M.F., M. Kähönen, M. Stumvoll, M. Putku, N.P., O.D., P. Mudgal, P.N., P.v.d.H., R.M.P., R.P.S.M., S.C., S.H.W., S. Sanna, T.E., T.H., T.L., V.V., W.H.L.K., X. Li, Y.O. and Z.K.

Interpretation of results: A. Tenesa, A. Tin, A. Köttgen, A.L.G., A. Teumer, B.M.P., C.G., D.R., E.A., G.W.M., H. Campbell, H. Snieder, J.K., M. Ciullo, M.A.N., M. Bochud, M. Caulfield, O.M.W., P.v.d.H., R.M.P., S.H.W., T.H., T.L., Toshihiro Tanaka, V.V., W.H.L.K., Y.O. and Z.K.

Manuscript review: A.A.H., A.B.S., A. Tenesa, A. Dehghan, A.D.J., A. Tin, A. Grotevendt, A. Goel, A.G.U., A.H., A.I., A. Jula, A. Köttgen, A.L., A.L.G., A. Kraja, A.M., A. Döring, A. Tönjes, A.P., A.R.S., A.S., A. Johansson, A. Teumer, A.V.S., B.B., B.H.R.W., B.M.P., B.O.B., B.R.W., B.W.P., C. Hundertmark, C. Hengstenberg, C. Sala, C.L., C.M., C.M.v.D., C.O., C.M.O., C.P.N., C. Schurmann, C.S.F., D.I.C., D.R., D.R.J., D.S.S., D.T., E.B., E.G.H., E. Theodoratou, F.C., F.E., F.R., F.T., G.A.T., G.C., G.G., G.N., G.W.M., H. Campbell, H. Choi, H. Schmidt, H.L.H., H.O., H. Snieder, H.V., H.W., I.B.B., I.K., I.M.L., I.M.N., I. Prokopenko, I.R., J.A., J.B.W., J.C., J.C.C., J.C.M.W., J.E.M., J.F.M., J.F.P., J.F.W., J.H.S., J.H.Z., J.I.R., J.K., J.S., J.S.K., J.V., K.B., K.L., K.S., K.T.K., L.J.L., L. Ferrucci, L.Y., M. Bruinenberg, M.A.N., M. Bochud, M. Caulfield, M. Ciullo, M.F., M.F.F., M.G.D., M.I., M. Burnier, M. Stumvoll, M. Kähönen, M. Kirin, M.M., M.N., M. Perola, M. Struchalin, M. Schallert, M.W., N.B.-N., N.G.M., N.J.W., N. Kamatani, N.M.P.-H., N.S., O.D., O.P., O.R., P. Sharma, P.F., P.K., P. McArdle, P.P.P., P. Salo, P.S.W., P.V., P.v.d.H., Q.Y., Q.Z., R. Schmidt, R.J.F.L., R.J.S., R.N., R.P.S., R. Sorice, S.B., S.H.W., S.J.L.B., S.K., S.L., S.R., S. Sanna, S.-Y.S., T.B.H., T.D.S., T.H., T.L., Toshihiro Tanaka, T. Zemunik, U.G., V.G., V.L., V.V., W.H.L.K., W.H.v.G., W.M., W.Z., X. Liu, Y.N., Y.O. and Z.K.

Analysis group: A. Köttgen, A. Teumer, C.G., C. Hundertmark, C.S.F., D.R., E.A., G.P., J.K., Q.Y., T.H., Toshihiro Tanaka, V.V. and W.H.L.K.

Writing group: A. Köttgen, A. Teumer, C.G., C.M.O., C.S.F., E.A., J.K., M. Bochud, M. Caulfield, M. Ciullo and V.V.

### COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

Reprints and permissions information is available online at <http://www.nature.com/reprints/index.html>.

Ilja M Nolte<sup>42</sup>, Asa Johansson<sup>43,44</sup>, Federico Murgia<sup>45</sup>, Sarah H Wild<sup>46</sup>, Stephan J L Bakker<sup>47</sup>, John F Peden<sup>48</sup>, Abbas Dehghan<sup>20,49</sup>, Maristella Steri<sup>50</sup>, Albert Tenesa<sup>5,51</sup>, Vasiliki Lagou<sup>52,53</sup>, Perttu Salo<sup>54</sup>, Massimo Mangino<sup>55</sup>, Lynda M Rose<sup>56</sup>, Terho Lehtimäki<sup>57</sup>, Owen M Woodward<sup>58</sup>, Yukinori Okada<sup>59,60</sup>, Adrienne Tin<sup>2</sup>, Christian Müller<sup>61</sup>, Christopher Oldmeadow<sup>62</sup>, Margus Putku<sup>63</sup>, Darina Czamara<sup>64</sup>, Peter Kraft<sup>65</sup>, Laura Frogheri<sup>45</sup>, Gian Andri Thun<sup>66,67</sup>, Anne Grotevendt<sup>68</sup>, Gauti Kjartan Gislason<sup>17</sup>, Tamara B Harris<sup>69</sup>, Lenore J Launer<sup>69</sup>, Patrick McArdle<sup>19</sup>, Alan R Shuldiner<sup>19</sup>, Eric Boerwinkle<sup>70</sup>, Josef Coresh<sup>2,71</sup>, Helena Schmidt<sup>72</sup>, Michael Schallert<sup>73</sup>, Nicholas G Martin<sup>22</sup>, Grant W Montgomery<sup>22</sup>, Michiaki Kubo<sup>74</sup>, Yusuke Nakamura<sup>75</sup>, Toshihiro Tanaka<sup>76</sup>, Patricia B Munroe<sup>77</sup>, Nilesh J Samani<sup>78,79</sup>, David R Jacobs Jr<sup>80</sup>, Kiang Liu<sup>81</sup>, Pio D'Adamo<sup>26</sup>, Sheila Ulivi<sup>82</sup>, Jerome I Rotter<sup>83</sup>, Bruce M Psaty<sup>27,84,85,86,87</sup>, Peter Vollenweider<sup>88</sup>, Gerard Waeber<sup>88</sup>, Susan Campbell<sup>5</sup>, Olivier Devuyst<sup>89</sup>, Pau Navarro<sup>5</sup>, Ivana Kolcic<sup>28</sup>, Nicholas Hastie<sup>5</sup>, Beverley Balkau<sup>90</sup>, Philippe Froguel<sup>29,91</sup>, Tõnu Esko<sup>12,63</sup>, Andres Salumets<sup>12,92,93</sup>, Kay Tee Khaw<sup>94</sup>, Claudia Langenberg<sup>31</sup>, Nicholas J Wareham<sup>31</sup>, Aaron Isaacs<sup>32</sup>, Aldi Kraja<sup>33</sup>, Qunyuan Zhang<sup>33</sup>, Philipp S Wild<sup>95,96</sup>, Rodney J Scott<sup>97</sup>, Elizabeth G Holliday<sup>62</sup>, Elin Org<sup>63</sup>, Margus Viigimaa<sup>98,99</sup>, Stefania Bandinelli<sup>100</sup>, Jeffrey E Metter<sup>14</sup>, Antonio Lupo<sup>101</sup>, Elisabetta Trabetti<sup>34</sup>, Rossella Sorice<sup>8</sup>, Angela Döring<sup>102,103</sup>, Eva Lattka<sup>104</sup>, Konstantin Strauch<sup>3,105</sup>, Fabian Theis<sup>6</sup>, Melanie Waldenberger<sup>104</sup>, H-Erich Wichmann<sup>103,106,107</sup>, Gail Davies<sup>35</sup>, Alan J Gow<sup>35,36</sup>, Marcel Bruinenberg<sup>108</sup>, LifeLines Cohort Study<sup>109</sup>, Ronald P Stolk<sup>42</sup>, Jaspal S Kooner<sup>110,111</sup>, Weihua Zhang<sup>111,112</sup>, Bernhard R Winkelmann<sup>113</sup>, Bernhard O Boehm<sup>114</sup>, Susanne Lucae<sup>64</sup>, Brenda W Penninx<sup>115,116,117</sup>, Johannes H Smit<sup>115</sup>, Gary Curhan<sup>118</sup>, Poorva Mudgal<sup>65</sup>, Robert M Plenge<sup>119,120,121</sup>, Laura Portas<sup>45</sup>, Ivana Persico<sup>45</sup>, Mirna Kirin<sup>46</sup>, James F Wilson<sup>46</sup>, Irene Mateo Leach<sup>122</sup>, Wiek H van Gilst<sup>122</sup>, Anuj Goel<sup>48</sup>, Halit Ongen<sup>48</sup>, Albert Hofman<sup>20,49</sup>, Fernando Rivadeneira<sup>20,49,123</sup>, Andre G Uitterlinden<sup>20,49,123</sup>, Medea Imboden<sup>66,67</sup>, Arnold von Eckardstein<sup>124</sup>, Francesco Cucca<sup>50</sup>, Ramaiah Nagaraja<sup>125</sup>, Maria Grazia Piras<sup>50</sup>, Matthias Nauck<sup>68</sup>, Claudia Schurmann<sup>4</sup>, Kathrin Budde<sup>68</sup>, Florian Ernst<sup>4</sup>, Susan M Farrington<sup>5</sup>, Evropi Theodoratou<sup>46</sup>, Inga Prokopenko<sup>52,53</sup>, Michael Stumvoll<sup>126,127</sup>, Antti Jula<sup>128</sup>, Markus Perola<sup>12,54,129</sup>, Veikko Salomaa<sup>54</sup>, So-Youn Shin<sup>130</sup>, Tim D Spector<sup>55</sup>, Cinzia Sala<sup>7</sup>, Paul M Ridker<sup>56,131</sup>, Mika Kähönen<sup>132</sup>, Jorma Viikari<sup>133</sup>, Christian Hengstenberg<sup>134</sup>, Christopher P Nelson<sup>78,79</sup>, CARDIOGRAM Consortium<sup>109</sup>, DIAGRAM Consortium<sup>109</sup>, ICBP Consortium<sup>109</sup>, MAGIC Consortium<sup>109</sup>, James F Meschia<sup>135</sup>, Michael A Nalls<sup>136</sup>, Pankaj Sharma<sup>137</sup>, Andrew B Singleton<sup>136</sup>, Naoyuki Kamatani<sup>138</sup>, Tanja Zeller<sup>61</sup>, Michel Burnier<sup>139</sup>, John Attia<sup>62</sup>, Maris Laan<sup>63</sup>, Norman Klopp<sup>104</sup>, Hans L Hillege<sup>37</sup>, Stefan Kloiber<sup>64</sup>, Hyon Choi<sup>118,140,141</sup>, Mario Pirastu<sup>45</sup>, Silvia Tore<sup>45</sup>, Nicole M Probst-Hensch<sup>66,67</sup>, Henry Völzke<sup>142</sup>, Vilmundur Gudnason<sup>17,18</sup>, Afshin Parsa<sup>19</sup>, Reinhold Schmidt<sup>73</sup>, John B Whitfield<sup>22</sup>, Myriam Fornage<sup>143,144</sup>, Paolo Gasparini<sup>26</sup>, David S Siscovick<sup>27,84,85</sup>, Ozren Poljšek<sup>28</sup>, Harry Campbell<sup>46</sup>, Igor Rudan<sup>28,46</sup>, Nabila Bouatia-Naji<sup>29</sup>, Andres Metspalu<sup>12,63</sup>, Ruth J F Loos<sup>31</sup>, Cornelia M van Duijn<sup>32</sup>, Ingrid B Borecki<sup>33,145</sup>, Luigi Ferrucci<sup>14</sup>, Giovanni Gambaro<sup>146</sup>, Ian J Deary<sup>35,36</sup>, Bruce H R Wolffenbuttel<sup>147</sup>, John C Chambers<sup>111,112</sup>, Winfried März<sup>40,148</sup>, Peter P Pramstaller<sup>41</sup>, Harold Snieder<sup>42</sup>, Ulf Gyllensten<sup>44</sup>, Alan F Wright<sup>5</sup>, Gerjan Navis<sup>149</sup>, Hugh Watkins<sup>48,150</sup>, Jacqueline C M Witteman<sup>20,49</sup>, Serena Sanna<sup>50</sup>, Sabine Schipf<sup>142</sup>, Malcolm G Dunlop<sup>5</sup>, Anke Tönjes<sup>126,127</sup>, Samuli Ripatti<sup>128,129,130</sup>, Nicole Soranzo<sup>130</sup>, Daniela Toniolo<sup>7,151</sup>, Daniel I Chasman<sup>56,131</sup>, Olli Raitakari<sup>152,153</sup>, W H Linda Kao<sup>2,71</sup>, Marina Ciullo<sup>8,156,157</sup>, Caroline S Fox<sup>9,10,154,156,157</sup>, Mark Caulfield<sup>77,156,157</sup>, Murielle Bochud<sup>155,156,157</sup>, and Christian Gieger<sup>3,156,157</sup>

<sup>1</sup>Renal Division, Freiburg University Hospital, Freiburg, Germany <sup>2</sup>Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA <sup>3</sup>Institute of Genetic Epidemiology, Helmholtz Zentrum München—German Research Center for Environmental Health, Neuherberg, Germany <sup>4</sup>Interfaculty Institute for Genetics and Functional Genomics, Ernst-Moritz-Arndt-University Greifswald, Greifswald, Germany <sup>5</sup>Medical Research

Council (MRC) Human Genetics Unit, MRC Institute of Genetics and Molecular Medicine (IGMM), University of Edinburgh, Edinburgh, UK <sup>6</sup>Institute of Bioinformatics and Systems Biology, Helmholtz Zentrum München–German Research Center for Environmental Health, Neuherberg, Germany <sup>7</sup>Division of Genetics and Cell Biology, San Raffaele Scientific Institute, Milan, Italy <sup>8</sup>Institute of Genetics and Biophysics A. Buzzati-Traverso, Consiglio Nazionale delle Ricerche (CNR), Naples, Italy <sup>9</sup>National Heart, Lung, and Blood Institute’s Framingham Heart Study, Framingham, Massachusetts, USA <sup>10</sup>National Heart, Lung, and Blood Institute’s Center for Population Studies, Framingham, Massachusetts, USA <sup>11</sup>Renal Division, Massachusetts General Hospital, Boston, Massachusetts, USA <sup>12</sup>Estonian Genome Center, University of Tartu, Tartu, Estonia <sup>13</sup>Department of Biostatistics, Boston University, Boston, Massachusetts, USA <sup>14</sup>Clinical Research Branch, National Institute on Aging, Baltimore, Maryland, USA <sup>15</sup>Department of Medical Genetics, University of Lausanne, Lausanne, Switzerland <sup>16</sup>Swiss Institute of Bioinformatics, Lausanne, Switzerland <sup>17</sup>Icelandic Heart Association Research Institute, Kopavogur, Iceland <sup>18</sup>Faculty of Medicine, University of Iceland, Reykjavik, Iceland <sup>19</sup>University of Maryland School of Medicine, Baltimore, Maryland, USA <sup>20</sup>Department of Epidemiology, Erasmus Medical Center, Rotterdam, The Netherlands <sup>21</sup>Department of Medicine, Prince Charles Hospital, Chermside, Queensland, Australia <sup>22</sup>Genetic Epidemiology, Queensland Institute of Medical Research, Brisbane, Queensland, Australia <sup>23</sup>Clinical Pharmacology Unit, University of Cambridge, Cambridge, UK <sup>24</sup>Division of Rheumatology, University of Alabama at Birmingham, Birmingham, Alabama, USA <sup>25</sup>Birmingham Veterans Affairs Medical Center, University of Alabama at Birmingham, Birmingham, Alabama, USA <sup>26</sup>Institute for Maternal and Child Health, Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS) Burlo Garofolo–Trieste, University of Trieste, Trieste, Italy <sup>27</sup>Cardiovascular Health Research Unit, University of Washington, Seattle, Washington, USA <sup>28</sup>Faculty of Medicine, University of Split, Split, Croatia <sup>29</sup>Centre National de la Recherche Scientifique (CNRS) Unité Mixte de Recherche (UMR) 8199, Genomics and Molecular Physiology of Metabolic Diseases, Institut Pasteur de Lille, Lille, France <sup>30</sup>CNRS UMR 8524, Laboratory of Mathematics, University Lille 1, Model for Data Analysis and Learning (MODAL) Team, Institut National de Recherche en Informatique et en Automatique (INRIA) Lille Nord–Europe, Lille, France <sup>31</sup>MRC Epidemiology Unit, Institute of Metabolic Science, Addenbrooke’s Hospital, Cambridge, UK <sup>32</sup>Subdivision of Genetic Epidemiology, Department of Epidemiology, Erasmus Medical Center, Rotterdam, The Netherlands <sup>33</sup>Division of Statistical Genomics, Department of Genetics, Washington University School of Medicine, St. Louis, Missouri, USA <sup>34</sup>Biology and Genetics Section, Department of Life and Reproduction Sciences, University of Verona, Verona, Italy <sup>35</sup>Department of Psychology, The University of Edinburgh, Edinburgh, UK <sup>36</sup>Centre for Cognitive Ageing and Cognitive Epidemiology, The University of Edinburgh, Edinburgh, UK <sup>37</sup>Department of Cardiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands <sup>38</sup>Institute of Clinical Science, Faculty of Medicine, Imperial College London, London, UK <sup>39</sup>Ludwigshafen Risk and Cardiovascular Health (LURIC) Study, Freiburg, Germany <sup>40</sup>Mannheim Institute of Public Health, Social and Preventive Medicine, Medical Faculty Mannheim, University of Heidelberg, Mannheim, Germany <sup>41</sup>Center for Biomedicine, European Academy Bozen/Bolzano (EURAC), Bolzano, Italy (affiliated institute of the University of Lübeck) <sup>42</sup>Department of Epidemiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands <sup>43</sup>Uppsala Clinical Research Center, Uppsala University Hospital, Uppsala, Sweden <sup>44</sup>Department of Immunology, Genetics and Pathology, Rudbeck Laboratory, Uppsala University, Uppsala, Sweden <sup>45</sup>Institute of Population Genetics, CNR, Sassari, Italy <sup>46</sup>Centre for Population Health Sciences, The University of Edinburgh Medical School, Edinburgh, UK <sup>47</sup>Department of Internal Medicine, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands <sup>48</sup>Department of Cardiovascular Medicine, Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK <sup>49</sup>Member of Netherlands Consortium for Healthy Aging (NCHA), Leiden, The Netherlands <sup>50</sup>Istituto di Ricerca Genetica e Biomedica, Consiglio Nazionale delle Ricerche,



Monserrato, Italy <sup>51</sup>Roslin Institute, The University of Edinburgh, Edinburgh, UK <sup>52</sup>Oxford Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, Oxford, UK <sup>53</sup>Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK <sup>54</sup>Department of Chronic Disease Prevention, National Institute for Health and Welfare, Helsinki, Finland <sup>55</sup>King's College London, St Thomas' Hospital Campus, London, UK <sup>56</sup>Division of Preventive Medicine, Brigham and Women's Hospital, Boston, Massachusetts, USA <sup>57</sup>Fimlab Laboratories, University of Tampere and Tampere University Hospital, Tampere, Finland <sup>58</sup>Department of Physiology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA <sup>59</sup>Laboratory for Statistical Analysis, Center for Genomic Medicine, RIKEN, Tokyo, Japan <sup>60</sup>Department of Allergy and Rheumatology, Graduate School of Medicine, University of Tokyo, Tokyo, Japan <sup>61</sup>University Heart Center Hamburg, Clinic for General and Interventional Cardiology, Hamburg, Germany <sup>62</sup>School of Medicine and Public Health, University of Newcastle, Newcastle, New South Wales, Australia <sup>63</sup>Institute of Molecular and Cell Biology, University of Tartu, Tartu, Estonia <sup>64</sup>Max Planck Institute of Psychiatry, Munich, Germany <sup>65</sup>Program in Molecular and Genetic Epidemiology, Harvard School of Public Health, Boston, Massachusetts, USA <sup>66</sup>Swiss Tropical and Public Health Institute, Basel, Switzerland <sup>67</sup>University of Basel, Basel, Switzerland <sup>68</sup>Institute of Clinical Chemistry and Laboratory Medicine, University Medicine Greifswald, Ernst-Moritz-Arndt-University Greifswald, Greifswald, Germany <sup>69</sup>Laboratory of Epidemiology, Demography and Biometry, National Institute on Aging, Bethesda, Maryland, USA <sup>70</sup>University of Texas Health Science Center at Houston, Houston, Texas, USA <sup>71</sup>Welch Center for Prevention, Epidemiology and Clinical Research, John Hopkins University, Baltimore, Maryland, USA <sup>72</sup>Institute of Molecular Biology and Biochemistry, Medical University Graz, Graz, Austria <sup>73</sup>Department of Neurology, Section of Special Neurology, Medical University Graz, Graz, Austria <sup>74</sup>Laboratory for Genotyping Development, Center for Genomic Medicine, RIKEN, Yokohama, Japan <sup>75</sup>Laboratory of Molecular Medicine, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, Japan <sup>76</sup>Laboratory for Cardiovascular Diseases, Center for Genomic Medicine, RIKEN, Yokohama, Japan <sup>77</sup>William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK <sup>78</sup>Leicester National Institute for Health Research (NIHR) Biomedical Research Unit in Cardiovascular Disease, Glenfield Hospital, Leicester, UK <sup>79</sup>Department of Cardiovascular Sciences, University of Leicester, Leicester, UK <sup>80</sup>Division of Epidemiology and Community Health, University of Minnesota, Minneapolis, Minnesota, USA <sup>81</sup>Department of Preventive Medicine, Feinberg School of Medicine, Northwestern University, Chicago, Illinois, USA <sup>82</sup>Institute for Maternal and Child Health, IRCCS Burlo Garofolo–Trieste, Trieste, Italy <sup>83</sup>Medical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, California, USA <sup>84</sup>Department of Epidemiology, University of Washington, Seattle, Washington, USA <sup>85</sup>Department of Medicine, University of Washington, Seattle, Washington, USA <sup>86</sup>Group Health Research Institute, Group Health Cooperative, Seattle, Washington, USA <sup>87</sup>Department of Health Services, University of Washington, Seattle, Washington, USA <sup>88</sup>Department of Medicine, Internal Medicine, Lausanne University Hospital, Lausanne, Switzerland <sup>89</sup>Institute of Physiology, Zurich Center for Integrative Human Physiology (ZIHP), University of Zurich, Zurich, Switzerland <sup>90</sup>Unité Mixte de Recherche en Santé (UMRS) 1018, University Paris Sud 11, Villejuif, France <sup>91</sup>Department of Genomics of Common Disease, School of Public Health, Imperial College London, Hammersmith Hospital, London, UK <sup>92</sup>Competence Centre on Reproductive Medicine and Biology, Tartu, Estonia <sup>93</sup>Department of Obstetrics and Gynecology, University of Tartu, Tartu, Estonia <sup>94</sup>Clinical Gerontology Unit, Addenbrooke's Hospital, Cambridge, UK <sup>95</sup>Center for Thrombosis and Hemostasis, University Medical Center Mainz, Johannes Gutenberg University Mainz, Mainz, Germany <sup>96</sup>Department of Medicine II, University Medical Center Mainz, Johannes Gutenberg University Mainz, Mainz, Germany <sup>97</sup>School of Biomedical Sciences and Pharmacy, University of Newcastle, Newcastle, New South Wales, Australia <sup>98</sup>Chair of Medical Physics, Department of Biomedical Engineering, Tallinn University of Technology, Tallinn, Estonia <sup>99</sup>Centre of Cardiology, North Estonia Medical

Centre, Tallinn, Estonia <sup>100</sup>Geriatric Unit, Azienda Sanitaria Firenze (ASF), Florence, Italy <sup>101</sup>Division of Nephrology, Department of Medicine, University of Verona, Verona, Italy <sup>102</sup>Institute of Epidemiology II, Helmholtz Zentrum München–German Research Center for Environmental Health, Neuherberg, Germany <sup>103</sup>Institute of Epidemiology I, Helmholtz Zentrum München–German Research Center for Environmental Health, Neuherberg, Germany <sup>104</sup>Research Unit of Molecular Epidemiology, Helmholtz Zentrum München–German Research Center for Environmental Health, Neuherberg, Germany <sup>105</sup>Chair of Genetic Epidemiology, Institute of Medical Informatics, Biometry and Epidemiology, Ludwig-Maximilians-University, Munich, Germany <sup>106</sup>Chair of Epidemiology, Institute of Medical Informatics, Biometry and Epidemiology, Ludwig-Maximilians-University, Munich, Germany <sup>107</sup>Klinikum Grosshadern, Munich, Germany <sup>108</sup>University Medical Center Groningen, LifeLines Department, University of Groningen, Groningen, The Netherlands <sup>110</sup>Faculty of Medicine, National Heart & Lung Institute, Cardiovascular Science, Hammersmith Hospital, Hammersmith Campus, Imperial College London, London, UK <sup>111</sup>Cath Laboratory, Cardiology, Ealing Hospital, Southall, UK <sup>112</sup>Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK <sup>113</sup>ClinPhenomics Study Center, Frankfurt, Germany <sup>114</sup>Division of Endocrinology and Diabetes, Department of Medicine, University Hospital, Ulm, Germany <sup>115</sup>Department of Psychiatry, EMGO Institute for Health and Care Research, VU University Medical Centre, Amsterdam, The Netherlands <sup>116</sup>Department of Psychiatry, University Medical Centre Groningen, University of Groningen, Groningen, The Netherlands <sup>117</sup>Department of Psychiatry, Leiden University Medical Center, Leiden, The Netherlands <sup>118</sup>Channing Laboratory, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA <sup>119</sup>Division of Rheumatology, Immunology and Allergy, Brigham and Women's Hospital, Boston, Massachusetts, USA <sup>120</sup>Division of Genetics, Brigham and Women's Hospital, Boston, Massachusetts, USA <sup>121</sup>Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA <sup>122</sup>Division of Experimental Cardiology, Department of Cardiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands <sup>123</sup>Department of Internal Medicine, Erasmus Medical Center, Rotterdam, The Netherlands <sup>124</sup>Institute of Clinical Chemistry, University Hospital of Zurich, Zurich, Switzerland <sup>125</sup>Laboratory of Genetics, National Institute on Aging, Baltimore, Maryland, USA <sup>126</sup>Medical Department, University of Leipzig, Leipzig, Germany <sup>127</sup>Integrated Research and Treatment Center (IFB) Adiposity Diseases, University of Leipzig, Leipzig, Germany <sup>128</sup>Department of Chronic Disease Prevention, National Institute for Health and Welfare, Turku, Finland <sup>129</sup>Institute of Molecular Medicine, University of Helsinki, Helsinki, Finland <sup>130</sup>Human Genetics, Wellcome Trust Sanger Institute, Hinxton, Cambridge, UK <sup>131</sup>Harvard Medical School, Boston, Massachusetts, USA <sup>132</sup>Department of Clinical Physiology, University of Tampere and Tampere University Hospital, Tampere, Finland <sup>133</sup>Department of Medicine, University of Turku and Turku University Hospital, Turku, Finland <sup>134</sup>Department of Internal Medicine II, University Hospital Regensburg, Regensburg, Germany <sup>135</sup>Department of Neurology, Mayo Clinic, Jacksonville, Florida, USA <sup>136</sup>Molecular Genetics Section, Laboratory of Neurogenetics, National Institute on Aging, US National Institutes of Health, Bethesda, Maryland, USA <sup>137</sup>Imperial College Cerebrovascular Research Unit, Imperial College London, London, UK <sup>138</sup>Laboratory for International Alliance, Center for Genomic Medicine, RIKEN, Yokohama, Japan <sup>139</sup>Nephrology Division, Department of Medicine, Lausanne University Hospital, Lausanne, Switzerland <sup>140</sup>Section of Rheumatology, Boston University School of Medicine, Boston, Massachusetts, USA <sup>141</sup>Clinical Epidemiology Research and Training Unit, Boston University School of Medicine, Boston, Massachusetts, USA <sup>142</sup>Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany <sup>143</sup>University of Texas–Houston Institute of Molecular Medicine, Houston, Texas, USA <sup>144</sup>University of Texas–Houston School of Public Health, Houston, Texas, USA <sup>145</sup>Division of Biostatistics, Washington University School of Medicine, St. Louis, Missouri, USA <sup>146</sup>Renal Program, Institute of Internal Medicine, Columbus-Gemelli University Hospital, Catholic University, Rome, Italy <sup>147</sup>Department

of Endocrinology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands <sup>148</sup>Synlab Centre of Laboratory Diagnostics, Heidelberg, Germany <sup>149</sup>Division of Nephrology, Department of Internal Medicine, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands <sup>150</sup>On behalf of PROCARDIS <sup>151</sup>Institute of Molecular Genetics, CNR, Pavia, Italy <sup>152</sup>Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, Turku, Finland <sup>153</sup>Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital, Turku, Finland <sup>154</sup>Division of Endocrinology, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA <sup>155</sup>University Institute of Social and Preventive Medicine, Lausanne, Switzerland

## Abstract

Elevated serum urate concentrations can cause gout, a prevalent and painful inflammatory arthritis. By combining data from >140,000 individuals of European ancestry within the Global Urate Genetics Consortium (GUGC), we identified and replicated 28 genome-wide significant loci in association with serum urate concentrations (18 new regions in or near *TRIM46*, *INHBB*, *SFMBT1*, *TMEM171*, *VEGFA*, *BAZ1B*, *PRKAG2*, *STC1*, *HNF4G*, *AICF*, *ATXN2*, *UBE2Q2*, *IGF1R*, *NFAT5*, *MAF*, *HLF*, *ACVR1B-ACVRL1* and *B3GNT4*). Associations for many of the loci were of similar magnitude in individuals of non-European ancestry. We further characterized these loci for associations with gout, transcript expression and the fractional excretion of urate. Network analyses implicate the inhibins-activins signaling pathways and glucose metabolism in systemic urate control. New candidate genes for serum urate concentration highlight the importance of metabolic control of urate production and excretion, which may have implications for the treatment and prevention of gout.

---

Uric acid is a final breakdown product of purine oxidation in humans and is present in the blood as urate. Elevated concentrations of serum urate—hyperuricemia—can cause gout<sup>1</sup>. Gout is the most prevalent inflammatory arthritis in developed countries, with an estimated 8.3 million US adults in 2007–2008 having had at least one of the extremely painful attacks<sup>2</sup>. Prevalence is increasing, owing in part to population aging, dietary and lifestyle factors, and rising levels of obesity and insulin resistance<sup>3–5</sup>. Chronic gout inflicts a considerable social and economic burden resulting from the associated pain and disability<sup>6,7</sup> as well as reduced work-related activity and productivity<sup>8</sup>.

Blood urate concentrations are determined by a balance between uric acid production, primarily in the liver, and its disposal via the kidney and gut. High circulating levels are mainly due to the net reabsorption of 90% of filtered urate in the renal proximal tubules<sup>9</sup>. Understanding the control of uric acid homeostasis is critical to improving the management and treatment of patients with hyperuricemia and gout.

The heritability of serum urate concentrations is estimated at 40–70% (refs. 10–12), which justifies the search for its genetic determinants. Previous genome-wide association studies (GWAS) have so far identified 11 genomic loci associated with urate concentrations and gout<sup>13–18</sup>. Together, SNPs at these loci explained about 5–6% of variance in serum urate concentrations<sup>17,18</sup>, suggesting that additional loci remain to be identified. We therefore aimed to identify and validate variants associated with serum urate concentrations in >140,000 individuals of European ancestry and with gout in approximately 70,000 individuals in the GUGC. These variants may provide a basis for the identification of new potential therapeutic targets for gout<sup>19</sup>.

## RESULTS

### Characteristics of the study samples

An overview of the characteristics of the GUGC study samples is provided in Supplementary Table 1. Our study comprises 110,347 individuals from 48 studies contributing to the discovery GWAS meta-analysis of serum urate concentrations and participants of 14 studies contributing to the meta-analysis of gout (2,115 cases and 67,259 controls). Across studies, mean serum urate concentrations ranged from 3.9 to 6.1 mg/dl (median of 5.2 mg/dl), and the proportion with gout ranged from 0.9% to 6.4% (median of 3.3%). Detailed information about the GUGC studies is provided in Supplementary Table 2. Information about study-specific genotyping, imputation, analysis tools and inflation factors for the individual serum urate concentration GWAS (range of 0.98–1.05) are shown in Supplementary Table 3.

### Loci associated with serum urate concentrations

A quantile-quantile plot for the 2,450,547 investigated autosomal SNPs in the discovery GWAS meta-analysis showed many more SNPs with low observed  $P$  values than expected, even after excluding SNPs in known urate concentration-associated regions (Supplementary Fig. 1). All 2,201 SNPs associated with serum urate concentrations at  $P < 5 \times 10^{-8}$  in the discovery stage are listed in Supplementary Table 4. Overall, 37 different genomic loci were identified that contained SNPs associated with serum urate concentrations at  $P < 1 \times 10^{-6}$ ; 26 of these were associated at genome-wide significance ( $P < 5 \times 10^{-8}$ , 10 known and 16 new loci; Fig. 1). Regional association plots provide a detailed overview of the associated regions (Supplementary Fig. 2). To assess the presence of independently associated SNPs within the regions associated at genome-wide significance and suggestive levels of significance ( $5 \times 10^{-8} < P < 1 \times 10^{-6}$ ), conditional analyses were carried out (Online Methods). This analysis identified only the *SLC22A11-NRXN2* region as containing more than one independent signal (Supplementary Table 5). The association at the previously reported *LRRC16A* locus<sup>17</sup> was not independent from that at the *SLC17A1* locus in our study. For each of the 37 independent loci, the index SNP with the lowest  $P$  value was carried forward for replication.

Replication was attempted in up to 32,813 new study participants. Successful replication was defined as  $q$  value  $< 0.05$  in the independent replication sample and genome-wide significance in the combined sample ( $P < 5 \times 10^{-8}$ ; Online Methods). Summary results for the 26 replicated urate concentration-associated loci are shown (Table 1): there were 16 newly identified regions in or near *TRIM46*, *INHBB*, *SFMBT1*, *TMEM171*, *VEGFA*, *BAZ1B*, *PRKAG2*, *STC1*, *HNF4G*, *A1CF*, *ATXN2*, *UBE2Q2*, *IGF1R*, *NFAT5*, *MAF* and *HLF* in addition to 10 known urate concentration-associated regions in or near *PDZK1*, *GCKR*, *SLC2A9*, *ABCG2*, *RREB1*, *SLC17A1*, *SLC16A9*, *SLC22A11*, *NRXN2* and *INHBC*. The proportion of variance in serum urate concentrations explained by these replicated loci in our data was 7.0%, with 5.2% of this estimate explained by the ten previously known loci and 3.4% explained by *SLC2A9* and *ABCG2* alone. The moderate proportion of trait variance explained is in line with findings from GWAS of other phenotypes<sup>20</sup>. The proportion of the age- and sex-adjusted variance in serum urate concentrations explained by all common SNPs rather than just the genome-wide significant index SNPs ranged from 0.27 (standard error (s.e.) of 0.04, ARIC Study) to 0.41 (s.e. of 0.07, SHIP Study) (Online Methods).

Additionally, regions in or near *ORC4L*, *OVOL1* and *BCAS3* were associated at genome-wide significance but had replication  $q$  values of  $> 0.05$ , whereas the *QRICH2* locus had a replication  $q$  value of  $< 0.05$  but did not reach genome-wide significance in the combined



sample. Effect estimates, allele frequencies and heterogeneity measures are presented separately for the discovery, replication and combined analyses for all 37 loci tested for replication (Supplementary Table 6). Although the functions of genes within the associated genomic regions were frequently connected to urate transport for the known loci, newly associated regions contained a series of transcription and growth factors encoding genes potentially connected to the metabolic control of serum urate production and excretion (Supplementary Table 7). Ten of the replicated loci also showed significant association with other complex traits, as listed in the National Human Genome Research Institute (NHGRI) GWAS Catalog<sup>21</sup>, and *VEGFA* and *ATXN2* showed significant associations with ratios of serum metabolites containing a  $\gamma$ -glutamyl amino acid from a published metabolite-SNP association resource<sup>22</sup>.

## Secondary analyses

Three secondary analyses were conducted to identify additional urate concentration-associated SNPs. First, as rare monogenic syndromes featuring gout can be caused by X-chromosome mutations in *PRPS1* (MIM 300661) and *HPRT1* (MIM 300322), we queried 54,926 X-chromosomal SNPs for association with serum urate concentrations in a meta-analysis of 72,026 participants from 25 of the GUGC studies (Supplementary Note). No genome-wide significant associations were observed, either in the combined or the sex-specific analyses (Supplementary Fig. 3); all X-chromosomal SNPs associated with serum urate concentrations at  $P < 1 \times 10^{-4}$  are listed (Supplementary Table 8).

Second, because of the higher prevalence of gout in men and the known sex-related differences in the effects of urate concentration-associated variants in *SLC2A9* and *ABCG2* (refs. 14,16), we conducted meta-analyses of GWAS separately for 49,825 men and 60,522 women. In addition to SNPs at loci detected in the combined analysis, SNPs were associated with serum urate concentrations at  $P < 1 \times 10^{-6}$  in one region in men and in five regions in women. None of these SNPs was significantly associated with serum urate concentrations in the replication samples (Supplementary Table 6), although the index SNP at *HNF1A* in women can be considered borderline significant with an association at  $P = 8.1 \times 10^{-8}$  in the combined discovery and replication data set. Besides *SLC2A9* and *ABCG2*, no additional regions contained SNPs that differed significantly ( $P < 5 \times 10^{-8}$ ) in their association effect sizes between men and women. Sex-related differences for genome-wide significant and suggestive SNPs of the overall and sex-stratified analyses are shown (Supplementary Table 9).

Third, we conducted a search for associated SNPs in genes that are family members of known urate transporter genes and, to the best of our knowledge, had not yet been connected to urate transport in humans. After correcting for multiple testing (Supplementary Note), SNPs in the *SLC22A7* region (*SLC22A11*-*SLC22A12* family member) showed significant association with serum urate concentrations in the discovery ( $P = 1.9 \times 10^{-5}$ ) and replication ( $q$  value  $< 0.05$ ) samples but did not reach the stringent genome-wide significance level in the combined samples (Table 2). It was recently found that the organic anion transporter 2 (OAT2), encoded by *SLC22A7*, transports urate in HEK293 cells<sup>23</sup>. Our findings show that variation in *SLC22A7* has a measurable effect on serum urate concentrations in humans, supporting the idea that it is a genuine urate concentration-associated GWAS locus of small effect size. Regional association plots for loci implicated through sex-specific analyses or the candidate gene approach are also shown (Supplementary Fig. 2).

## Gout GWAS

We conducted a genome-wide gout discovery meta-analysis including 2,538,056 autosomal SNPs. Two previously reported loci had associations that reached genome-wide significance

in the overall and sex-stratified analyses: *ABCG2* (rs1481012,  $P = 2.0 \times 10^{-32}$ ) and *SLC2A9* (rs4475146,  $P = 4.1 \times 10^{-26}$ ) (Supplementary Fig. 4 and Supplementary Table 10). The quantile-quantile plot of the  $P$  values from the combined analysis showed more observed low  $P$  values than expected by chance, even after excluding the previously known urate concentration–associated regions (Supplementary Fig. 5). The index SNPs in *SLC2A9* and *ABCG2* from the combined analysis of gout were in high linkage disequilibrium (LD;  $r^2 > 0.9$ ) with the index SNPs from the serum urate concentration meta-analysis at these loci.

### Serum urate concentration–associated loci and gout

Because elevated serum urate levels are a key risk factor for the development of gout, we investigated the association of the urate concentration–associated SNPs with gout in the data set described. We observed a positive linear correlation between the genetic effect on serum urate concentrations and the log odds of gout for the replicated loci (Pearson's correlation = 0.93; Fig. 2). In addition, information was available on the associations with incident gout over a period of up to 22 years in two independent studies that were not part of either the gout or serum urate concentration discovery analyses. The 1,036 cases of gout in these studies met the American College of Rheumatology Criteria<sup>24,25</sup>. In all gout samples combined (3,151 cases and 68,350 controls), 17 out of 26 of the replicated urate concentration–associated SNPs showed nominal association with gout ( $P < 0.05$ ; Table 1). SNP associations with prevalent and incident gout were of generally comparable direction and magnitude overall, as well as among men and women separately (Supplementary Table 11).

### Genetic urate risk score is associated with gout

A weighted genetic urate score was constructed on the basis of the number of risk alleles across loci from the main analysis as described previously<sup>26</sup> and scaled to a risk allele count range as outlined (Supplementary Note). The association between urate scores and gout was evaluated in studies with a large number of cases (prevalent gout: ARIC, SHIP and KORA F4; incident gout: NHS and HPFS). Risk scores ranged from 10 to 45 (mean of  $31 \pm 5$ ). Risk scores were significantly associated with increased odds of prevalent gout (odds ratio (OR) = 1.11 per risk score unit increase, 95% confidence interval (CI) = 1.09–1.14;  $P = 2.5 \times 10^{-29}$ ;  $n = 693$  cases) and incident gout over a period of up to 22 years (OR = 1.10, 95% CI = 1.08–1.13;  $P = 3.7 \times 10^{-21}$ ;  $n = 1,036$  cases). Gout prevalence increased from <1% to 18% across risk score categories in the population-based studies, and the increased prevalence of gout with higher genetic urate score categories could be replicated in the independent studies recording incident cases of gout (Supplementary Fig. 6).

### Characterization of serum urate concentration–associated loci

To gain insight into how the identified variants might contribute to altered serum urate levels, we evaluated their association with the fractional excretion of uric acid (FEUA,  $n = 6,799$ ; Supplementary Table 12). FEUA is the proportion of urate filtered by the glomeruli that is eventually excreted in the urine. Physiological values are typically well below 10%. SNPs at 10 replicated loci showed nominal association with FEUA (Table 3); SNPs at the *SLC2A9*, *GCKR* and *IGF1R* loci passed a multiple testing–corrected threshold for 26 replicated SNPs ( $P < 1.9 \times 10^{-3}$ ). In all ten instances, the allele associated with higher serum urate concentrations was associated with lower FEUA.

### Associations in individuals of non-European ancestry

To evaluate whether the observed SNP associations are generalizable to individuals of non-European ancestry, we investigated the replicated and genome-wide significant urate concentration–associated SNPs in 8,340 individuals of Indian ancestry, 5,820 African-

Americans and 15,286 Japanese (Supplementary Table 13). Although allele frequencies at the index SNPs varied considerably across the groups (Fig. 3, right), the effects on serum urate concentrations were of comparable magnitude and identical direction for the majority of SNPs (Fig. 3, left), further supporting the idea that these index SNPs represent true urate concentration–associated loci.

### Association with transcript expression in *cis*

The association of implicated SNPs with transcript expression in *cis* can potentially provide insights into the gene underlying the association signal. We therefore queried existing expression quantitative trait locus (eQTL) databases containing data from various tissues (Supplementary Note). For ten of the replicated loci, there were significant eSNPs that were associated with the expression of one or more transcript in one or more tissue (Supplementary Table 14). These eSNPs were either the index SNP itself or a variant in high LD with the index SNP ( $r^2 > 0.8$ ); we highlight only instances where the index SNP or a perfect proxy was the best eSNP for a given transcript and tissue. These analyses identify the *INHBB* gene rather than any of the neighboring genes as being causative at 2q13 because only *INHBB* transcript levels were significantly associated with the index SNP in the different data sets queried ( $P$  value range of  $3 \times 10^{-5}$  to  $2 \times 10^{-14}$ ).

### Association with correlated traits

The 26 identified and replicated urate concentration–associated loci were evaluated for their association with complex traits that are phenotypically correlated with serum urate concentrations (blood pressure, cardiovascular outcomes and glucose homeostasis; Supplementary Table 15). SNPs at the *GCKR*, *BAZ1B* and *ATXN2* loci showed genome-wide significant associations with one or more of these traits. The direction of association between serum urate and plasma C-reactive protein (CRP) concentrations at *GCKR* and *BAZ1B* was consistent with the epidemiologically observed associations, as was the case for the pleiotropic *ATXN2* locus and blood pressure.

In aggregate, a weighted risk score composed of the 26 replicated urate concentration–increasing alleles was significantly associated ( $P < 4.5 \times 10^{-3}$ ) only with plasma CRP concentrations after correcting for the number of traits investigated (Supplementary Note). This association was driven by the pleiotropic *GCKR* locus rather than being a general effect of urate concentration–associated variants on the inflammatory marker CRP; the association between the score and CRP concentrations was abolished in a sensitivity analysis when rs1260326 in *GCKR* was excluded from the genetic urate score. The absence of significant associations between the genetic urate score and the investigated traits, including blood pressure, might suggest a lack of causal relationships between serum urate concentrations and these traits, given that the present study had adequate statistical power to detect significant associations, as shown previously in a smaller subset<sup>18</sup>. However, the methodology used has limitations, and alternative explanations cannot be ruled out.

### Network analysis identifies additional loci

To identify further urate concentration–associated genomic regions and in order to place these into a biologically interpretable context, we incorporated previous knowledge on molecular interactions in which the gene products of implicated genes operate. First, GRAIL<sup>27</sup> was used to infer the most likely gene underlying the association at each of the 37 loci from the discovery analyses of the sex-combined sample (see Supplementary Note for details and Supplementary Table 16 for an overview of GRAIL results). We then constructed a functional association network using the most likely genes as seed genes (Supplementary Note). The complete networks of one-, two- and three-edge neighborhoods are provided as online supplements (see URLs) and highlight genes that are implicated by

forming functional associations, mostly protein-protein interactions, with one of the seed genes. To systematically identify urate concentration–associated SNPs in these genes, we selected the SNP with the lowest urate concentration–associated  $P$  value in the new candidate genes implicated by the network analyses that passed a multiple testing–corrected threshold ( $P < 6.8 \times 10^{-5}$ , Bonferroni correction for SNPs in all implicated genes). Replication was attempted for the resulting 17 SNPs in up to 23,078 participants (Supplementary Table 6). Replication criteria, defined as in the discovery analysis, were met for two regions: *B3GNT4* and *ACVR1B-ACVRL1* (Fig. 1 and Table 2). Moreover, *ARNT* had a suggestive replication  $q$  value of  $<0.05$  but did not reach genome-wide significance. Regional association plots are shown (Supplementary Fig. 7).

A specific subnetwork from the analysis around the inhibins-activins pathway was highlighted as part of the functional association network analysis, in which three seed genes (*INHBB*, *INHBC-INHBE* and *ACVR2A*) were connected with three genes implicated by the functional association network analysis (*ACVR1B-ACVRL1*, *ACVR1C* and *BMP2*), one of which replicated (Supplementary Fig. 8). The gene products operate in one pathway and are inhibins-activins receptors and their ligands. Detailed information on all replicated genes in the inhibins-activins network is provided in the Supplementary Note. Another pathway linking a large number of GWAS discovery association signals contains genes involved in growth factor signaling through tyrosine kinase receptors implicated in the control of cell growth and differentiation (Supplementary Fig. 9). One of these genes, *PKRL*, is also highlighted in the Supplementary Note, which provides information about identified associated loci that include candidate genes with a role in glucose metabolism. The functional network associations that formed the basis for Supplementary Figures 8 and 9 are provided in Supplementary Table 17. Pathway analyses using Ingenuity Pathway Analysis software (Supplementary Note) showed functional network associations with gene expression, cellular organization, carbohydrate metabolism, molecular transport and endocrine system disorders (lowest  $P = 1 \times 10^{-28}$ ; Supplementary Table 18).

## DISCUSSION

In GUGC, we identified and replicated 28 genome-wide significant urate concentration–associated loci, 18 of which were newly identified, using GWAS ( $n = 26$ ) and pathway ( $n = 2$ ) approaches. The loci from our GWAS meta-analysis were characterized in detail, including analyzing their associations with serum urate concentrations in samples of non-European ancestry and with the fractional excretion of urate. The serum urate concentration–increasing allele at all loci was associated with higher risk of gout. The functional association network analysis supports a new role for inhibins-activins pathways in regulating urate homeostasis.

We confirmed the presence of urate concentration–associated SNPs from previous GWAS for all 11 previously reported loci<sup>13–18</sup>, 6 of which map to genes encoding proteins related to urate transport across membranes. Conversely, none of the genes at the newly associated loci from this study seem to be obvious candidates for factors involved in urate transport. A primary candidate for having an influence on urate generation is *PRPSAP1*, which encodes a protein involved in the regulation of purine synthesis<sup>28</sup>. Many of the remaining index SNPs map to genes coding for transcription or growth factors with broad downstream responses. Some of these may ultimately affect the activity of known urate transporters, such as the transcription factor HNF-1 $\alpha$ , encoded by a suggestive urate concentration–associated locus that has response elements upstream of *ABCG2*, *SLC17A3*, *SLC22A11*, *SLC16A9* and *PDZK1*.



Many of the newly identified loci can be connected into functional or biochemical pathways. All pathway approaches evaluated in our analyses highlight terms related to carbohydrate metabolism, such as regulation of glycolysis, glucose, insulin and pyruvate (Supplementary Tables 16 and 18). In addition to the previously identified *GCKR* locus, the gene products of five of the newly discovered replicated loci (Supplementary Note) directly modulate glucose flux. The association of serum urate concentrations with these glucose metabolism-related loci may be explained by increased flow of glucose-6-phosphate through the pentose phosphate pathway and/or high levels of aerobic glycolysis with lactate production as observed in proliferating cells (Warburg effect)<sup>29–31</sup>. The pentose phosphate pathway generates ribose-5-phosphate, a key precursor of *de novo* purine synthesis and thereby of uric acid production. The amount of lactate generated can influence urate transport across membranes and can therefore alter urate elimination by the kidney. Detailed gene information and an illustration of how several of these genes are connected in regulatory networks targeting the rate-limiting glycolytic enzymes glucokinase and pyruvate kinase are provided (Supplementary Note). Moreover, insulin and insulin-like growth factor 1, the receptors of which showed suggestive and significant association with urate concentrations, induce glucokinase gene (*GCK*) expression in the liver and pancreas, respectively<sup>32</sup>.

In addition to known urate transporters expressed in the renal proximal tubules, three loci linked to glucose metabolism and/or insulin response (*GCKR*, *IGF1R* and *NFAT5*) were nominally associated with FEUA. The urate concentration-increasing alleles at all loci were associated with lower FEUA, which could either be owing to increased urate reabsorption in the kidney by the mechanisms discussed or possibly to the reported effect of insulin on decreasing renal urate clearance and sodium excretion in healthy people<sup>33,34</sup>.

Despite the connection of these new candidate genes for urate regulation in networks related to glucose metabolism, there was no significant aggregate effect of urate concentration-associated loci on glycemic traits or measures of insulin resistance. A possible explanation might be the absence of a sizeable effect of SNPs in heterogeneous pathways on a single related physiological process or the lack of statistical power of current GWAS efforts to detect small effect sizes.

Another major pathway highlighted by our analyses is the inhibins-activins growth factor system (Supplementary Fig. 8 and Supplementary Note). This pathway is thought to mediate a very wide range of processes, which could affect urate levels through a variety of known functions, such as energy balance, insulin release, apoptosis, inflammation and sex hormone regulation. In addition, our approach demonstrates the value of the functional association network analyses in identifying additional loci from conventional GWAS results.

The mainstay of gout therapy is allopurinol, which inhibits the formation of urate and is an inexpensive and effective preventative agent for chronic gout. Although other therapeutic options exist, the overall number of urate concentration-lowering agents approved for clinical use is limited. Data from a clinical trial indicate that only 21% of patients offered the most common dose of allopurinol (300 mg/d) achieve optimal levels of serum urate<sup>35</sup>. In addition, there are rare, serious side-effects, and reduced doses of allopurinol are recommended in individuals with impaired renal function. The identification of additional therapeutic options to lower serum urate levels is therefore an active area of research. Our finding that gene loci implicated in glucose homeostasis influence serum urate concentrations fits with the observation that drugs that decrease insulin resistance, such as thiazolidinediones and metformin, tend to decrease serum urate levels<sup>36–40</sup>. Thus, the identification of new gene loci in the present study may open new avenues for research to improve the treatment and prevention of gout.

The major strengths of our study are the large sample size with >110,000 individuals for the discovery step, the independent replication of associations and the possibility to examine identified loci for their relationship to prevalent and American College of Rheumatology Criteria–confirmed incident gout. The comprehensive evaluation of urate concentration–associated variants provided additional insights by detecting associations with fractional urate excretion and transcript expression.

Limitations of our study include the relatively modest sample size available for the replication step, which compromises power and potentially results in an inability to validate true urate concentration–associated loci such as *ORC4L*, *OVOL1* and *BCAS3*. The definition of gout varied across several cohorts, which may have led to some misclassification but is unlikely to result in false positive results. As with other GWAS, our analysis focused mostly on common SNPs, and we could therefore not assess the contributions of rare variants, such as a recently described risk variant for gout<sup>41</sup>.

We identified 28 genome-wide significant loci associated with serum urate concentrations. We found that alleles associated with increased serum urate concentrations were also associated with increased risk of gout. The modulation of urate production and excretion by signaling processes that influence metabolic pathways, such as glycolysis and the pentose phosphate pathway, seem to be central pathways including the genes from the newly identified associated loci. These findings may have implications for further research into urate concentration–lowering drugs to treat and prevent the common inflammatory arthritis gout.

## URLs

Complete results from network analyses can be found at <http://www.gwas.eu/gugc/>. R, <http://www.r-project.org/>; Quanto, <http://hydra.usc.edu/gxe/>.

## ONLINE METHODS

### Participating studies

The overall study comprises data of >70 study samples as detailed in Supplementary Table 1. The meta-analysis of serum urate levels combined genome-wide scans of 48 studies totaling 110,347 individuals, and the meta-analysis of gout comprised 14 studies totaling 2,115 cases and 67,259 controls.

Selected SNPs from the serum urate analysis were followed up in 12 studies with *in silico* GWAS data totaling 18,821 individuals, as well as 3 studies in which *de novo* genotyping was conducted (HYPEST, KORA S2 and Ogliastra Genetic Park, totaling 13,992 individuals). All individuals were of European descent.

Additional study samples consisted of two nested case-control studies of incident gout (HPFS and NHS, totaling 1,036 cases and 1,091 controls), as well as 8,340 participants of Indian ancestry (LOLIPOP), 5,820 African-Americans (ARIC, CARDIA and JHS) and 15,288 samples of Japanese ancestry (BioBank Japan).

A list of all studies contributing to the different analyses is provided in the Supplementary Note. Information on study design, study-specific exclusions, urate concentration measurement and gout definition for each study is provided in Supplementary Table 2. The individual studies obtained written informed consent from their study participants, and the studies were approved by their respective local ethics committees.

## Power calculations and explained variance

Power calculations were performed a priori, separately for discovery and replication, using Quanto (see URLs). The standard deviation of serum urate concentrations for the calculations was taken from the population-based ARIC Study<sup>42</sup>, one of the largest contributing studies. For a discovery analysis in 110,000 individuals (urate), there was >80% power to detect a SNP explaining 0.04% or more of the variance in serum urate concentration at  $P < 5 \times 10^{-8}$  (corresponding to a ~0.04 mg/dl change in serum urate per allele). Assuming a 2% population prevalence of gout, there was >80% power to detect a risk variant conferring 30% increased odds of gout in 3,000 cases and ~67,000 controls at  $P < 5 \times 10^{-8}$  for a SNP with minor allele frequency (MAF) of 10%.

We further used the method by Park *et al.*<sup>43</sup> to estimate that 16 new genome-wide significant urate concentration-associated loci would be detected, assuming a study of 100,000 individuals and effect, frequency and statistical power of previously published loci<sup>18</sup> as the reference. The proportion of the explained variance in serum urate concentrations for this study was estimated at 7.7%.

For the replication analyses, 40,546 samples were estimated to be required to independently replicate a SNP with MAF of 10% and effect of 0.06 mg/dl per allele with 80% power (or 15,201 samples for a SNP with MAF of 0.4). For this analysis, we assumed 20 SNPs carried forward to replication, with statistical significance defined as a multiple testing-corrected one-sided  $P$  value.

The proportion of variance in serum urate concentrations explained by all 26 independent replicated SNPs from the overall analysis was calculated as the sum of the mean proportion of variance explained by each SNP separately under an additive model adjusted for sex and age, calculated within 31 cohorts. The proportion of the age- and sex-adjusted variance in serum urate concentration explained by all common (MAF > 0.01) genotyped SNPs that met quality control criteria was calculated in three large population-based studies (ARIC,  $n = 9,049$ ; CoLaus,  $n = 5,409$ ; SHIP,  $n = 4,067$ ) using the REML method in GCTA software<sup>44</sup>.

## Statistical analyses in the individual studies

In each study of the discovery GWAS, genotyping was performed on genome-wide chips, and imputation was conducted using HapMap 2 data as the reference. Quality control before imputation was applied in each study separately. Detailed genotyping and imputation information for each study is provided (Supplementary Table 3).

In the individual-study GWAS, outcomes were related to the SNP dosages using linear (urate concentration, mg/dl) or logistic (gout) regression adjusted for age and sex as well as study-specific covariates, if applicable (for example, principal components and study center). Secondary sex-stratified analyses were also conducted. An additive genetic model was used.

## Quality control

Each file of genome-wide per-SNP summary statistics underwent extensive quality control before meta-analysis. Examination of file formatting, data plausibility and distributions of test statistics and quality measurements was facilitated by the `gwasqc()` function of the GWAToolbox package v1.0.0 in R<sup>45</sup>. Additionally, the direction and magnitude of effect at the known urate concentration-associated SNP rs16890979 in *SLC2A9* was investigated, and the minor allele was consistent in frequency and associated with lower serum urate concentrations in all studies.

Before all meta-analyses, monomorphic SNPs were excluded, and all study-specific results were corrected by the genomic inflation factor of the study if it was  $>1$ , calculated by dividing the median of the observed  $\chi$ -square distribution of the GWAS by the median of the expected  $\chi$ -square distribution under the null hypothesis of no association. Study-specific inflation factors for urate analyses are shown in Supplementary Table 3. For meta-analyses of gout, only cohorts with  $>50$  gout cases were included, and SNPs with an effect size of  $|\beta| > 1,000$  were excluded to remove only a minimum number of SNPs with implausibly large effects that could systematically influence the results.

### Meta-analyses

All meta-analyses were carried out in duplicate by two independent analysts. Meta-analysis was performed on the results of all genome-wide scans using a fixed-effects model applying inverse variance weighting as implemented in METAL<sup>46</sup>. Results were confirmed by comparing the results to those from a  $z$  score-based meta-analysis. SNPs that were present in  $<75\%$  of all samples contributing to the respective meta-analysis were excluded, and the remaining SNPs were used as the basis for all subsequent analyses.

Consequently, data for 2,450,547 genotyped or imputed autosomal SNPs were available for the primary analysis of serum urate concentrations, and data for 2,538,056 SNPs were available for the primary gout analysis. After all meta-analyses, a second genomic control correction was applied to primary analyses (genomic inflation factor = 1.12 for urate and 1.03 for gout) as well as secondary sex-stratified analyses (genomic inflation factor for urate = 1.07 (men), 1.08 (women) and gout = 1.04 (men), 1.03 (women)). The top three SNPs from the gout meta-analysis results among women were removed because they were thought to represent false positive results on the basis of low MAF (0.02), very low  $P$  values and an  $I^2$  heterogeneity measure of  $>99\%$ .

Genome-wide significance was defined as a  $P$  value of  $<5 \times 10^{-8}$  after the second correction for genomic control, corresponding to a Bonferroni correction of 1 million independent tests<sup>47</sup>. All loci containing SNPs of suggestive significance ( $P < 1 \times 10^{-6}$ ) in the discovery-stage meta-analyses were considered for further analysis and assessed for the presence of independent SNPs within each locus.

### Assessment of the presence of independent signals at each locus

Two steps were performed to identify independent SNPs in the same region that associated with serum urate concentration in the overall analyses. First, SNPs with  $P$  values of  $<1 \times 10^{-5}$  were aggregated on the basis of the LD structure from the HapMap release 28 Utah residents of Northern and Western European ancestry (CEU) data set using PLINK<sup>48</sup> (settings  $r^2 > 0.01$ , 1-Mb distance). Second, to verify the potential independent associations from the first step, a multiple-regression model was calculated in 32 of the studies, adjusting for all lead SNPs at once, and meta-analysis was performed, followed by a comparison of the effect estimate for each SNP with those from the single-SNP association model using a  $t$  test (Supplementary Table 5). Independent SNPs were defined as those with (i) a  $t$ -test  $P$  value of  $>0.05$  and (ii) a difference of the effect estimates between the single- and the multiple-SNP models of  $\geq 20\%$  compared to the single-SNP model. rs1178977 in *BAZ1B* did not fulfill the criteria for independence (20.7% change in effect estimate). Yet, it was treated as an independent SNP, as the adjacent associated genomic region was located  $>75$  Mb away, suggesting that the change in  $\beta$  estimate might be due to other factors than a nearby SNP in LD. Analyses were performed in R.

Subsequently, 37 independent loci of suggestive significance ( $P < 1 \times 10^{-6}$ ) were carried forward for replication.



## Secondary analyses

To further identify genetic variants associated with serum urate concentrations and gout, several additional analyses were carried out. These included X-chromosome analyses, sex-stratified analyses, a gene-based test and urate transporter candidate analyses, as well as secondary analyses to improve the characterization of associated SNPs. To this end, we investigated associations with FEUA, associations in individuals of non-European ancestry, associations with transcript expression, associations with serum metabolite concentrations, risk score analyses and associations with other urate-related phenotypes. Secondary analyses are described in detail in the Supplementary Note.

As a result of the secondary analyses, seven additional SNPs were identified for further replication testing: one from the analysis in men only, five from the analysis in women and one from the urate transporter candidate analysis.

A new part of the secondary analyses was the implementation of functional association networks as described in the Supplementary Note. As detailed there, the approach based on functional associations among the urate genes, mostly protein-protein interactions, led to the identification of 17 additional independent SNPs in newly identified genomic regions that were subjected to replication testing. Altogether, we therefore tested 61 SNPs (37 overall, 6 sex specific, 1 urate transporter candidate and 17 network) for replication in additional study samples.

## Replication analyses

Up to 18,821 participants contributed information to the *in silico* replication analyses of the 61 independent variants tested, and additional *de novo* genotyping was conducted in up to 13,992 individuals. The cohorts are described in Supplementary Table 1. We excluded SNPs with call rate of <0.9 or imputation quality of <0.3 (MACH) or <0.4 (Impute, BEAGLE). The statistical methods used to estimate the associations were identical to the ones used in the discovery cohorts.

Details about *de novo* genotyping, including platforms, array design and quality control, are provided in the Supplementary Note.

To assess evidence for replication, test statistics of all *in silico* and *de novo* replication cohorts were combined using fixed-effects and inverse variance weighting using METAL<sup>46</sup>. The false discovery rate was estimated in the independent replication samples as a *q* value on the basis of the *P*-value distribution of all SNPs tested for replication using the *q*value package in R<sup>49</sup>. Replication was defined, for each SNP, as *q* value < 0.05 in the replication samples alone, indicating that <5% false positives would be expected among these variants. To place an additional filter on these results, only SNPs that were associated with genome-wide significance ( $P < 5 \times 10^{-8}$ ) in the combined analysis were considered to have been successfully replicated; combined estimates were obtained by meta-analysis of the double-corrected test statistics of the discovery GWAS with the replication estimates.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

A detailed list of acknowledgments is provided in the Supplementary Note.


## References

1. So A, Thorens B. Uric acid transport and disease. *J Clin Invest*. 2010; 120:1791–1799. [PubMed: 20516647]
2. Zhu Y, Pandya BJ, Choi HK. Prevalence of gout and hyperuricemia in the US general population: the National Health and Nutrition Examination Survey 2007–2008. *Arthritis Rheum*. 2011; 63:3136–3141. [PubMed: 21800283]
3. Arromdee E, Michet CJ, Crowson CS, O’Fallon WM, Gabriel SE. Epidemiology of gout: is the incidence rising? *J Rheumatol*. 2002; 29:2403–2406. [PubMed: 12415600]
4. Wallace KL, Riedel AA, Joseph-Ridge N, Wortmann R. Increasing prevalence of gout and hyperuricemia over 10 years among older adults in a managed care population. *J Rheumatol*. 2004; 31:1582–1587. [PubMed: 15290739]
5. Yoo HG, et al. Prevalence of insulin resistance and metabolic syndrome in patients with gouty arthritis. *Rheumatol Int*. 2011; 31:485–491. [PubMed: 20091036]
6. Singh JA. Quality of life and quality of care for patients with gout. *Curr Rheumatol Rep*. 2009; 11:154–160. [PubMed: 19296889]
7. Becker MA, et al. Quality of life and disability in patients with treatment-failure gout. *J Rheumatol*. 2009; 36:1041–1048. [PubMed: 19332629]
8. Brook RA, et al. The economic burden of gout on an employed population. *Curr Med Res Opin*. 2006; 22:1381–1389. [PubMed: 16834837]
9. Hediger MA, Johnson RJ, Miyazaki H, Endou H. Molecular physiology of urate transport. *Physiology (Bethesda)*. 2005; 20:125–133. [PubMed: 15772301]
10. Whitfield JB, Martin NG. Inheritance and alcohol as factors influencing plasma uric acid levels. *Acta Genet Med Gemellol (Roma)*. 1983; 32:117–126. [PubMed: 6685961]
11. Yang Q, et al. Genome-wide search for genes affecting serum uric acid levels: the Framingham Heart Study. *Metabolism*. 2005; 54:1435–1441. [PubMed: 16253630]
12. Nath SD, et al. Genome scan for determinants of serum uric acid variability. *J Am Soc Nephrol*. 2007; 18:3156–3163. [PubMed: 17978310]
13. Li S, et al. The *GLUT9* gene is associated with serum uric acid levels in Sardinia and Chianti cohorts. *PLoS Genet*. 2007; 3:e194. [PubMed: 17997608]
14. Doring A, et al. *SLC2A9* influences uric acid concentrations with pronounced sex-specific effects. *Nat Genet*. 2008; 40:430–436. [PubMed: 18327256]
15. Vitart V, et al. *SLC2A9* is a newly identified urate transporter influencing serum urate concentration, urate excretion and gout. *Nat Genet*. 2008; 40:437–442. [PubMed: 18327257]
16. Dehghan A, et al. Association of three genetic loci with uric acid concentration and risk of gout: a genome-wide association study. *Lancet*. 2008; 372:1953–1961. [PubMed: 18834626]
17. Kolz M, et al. Meta-analysis of 28,141 individuals identifies common variants within five new loci that influence uric acid concentrations. *PLoS Genet*. 2009; 5:e1000504. [PubMed: 19503597]
18. Yang Q, et al. Multiple genetic loci influence serum urate levels and their relationship with gout and cardiovascular disease risk factors. *Circ Cardiovasc Genet*. 2010; 3:523–530. [PubMed: 20884846]
19. Woodward OM, et al. Identification of a urate transporter, *ABCG2*, with a common functional polymorphism causing gout. *Proc Natl Acad Sci USA*. 2009; 106:10338–10342. [PubMed: 19506252]
20. Visscher PM, Brown MA, McCarthy MI, Yang J. Five years of GWAS discovery. *Am J Hum Genet*. 2012; 90:7–24. [PubMed: 22243964]
21. Hindorf LA, et al. Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. *Proc Natl Acad Sci USA*. 2009; 106:9362–9367. [PubMed: 19474294]
22. Suhre K, et al. Human metabolic individuality in biomedical and pharmaceutical research. *Nature*. 2011; 477:54–60. [PubMed: 21886157]
23. Sato M, et al. Renal secretion of uric acid by organic anion transporter 2 (*OAT2/SLC22A7*) in human. *Biol Pharm Bull*. 2010; 33:498–503. [PubMed: 20190416]


24. Wallace SL, et al. Preliminary criteria for the classification of the acute arthritis of primary gout. *Arthritis Rheum.* 1977; 20:895–900. [PubMed: 856219]
25. Choi HK, Atkinson K, Karlson EW, Willett W, Curhan G. Purine-rich foods, dairy and protein intake, and the risk of gout in men. *N Engl J Med.* 2004; 350:1093–1103. [PubMed: 15014182]
26. Ehret GB, et al. Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature.* 2011; 478:103–109. [PubMed: 21909115]
27. Raychaudhuri S, et al. Identifying relationships among genomic disease regions: predicting genes at pathogenic SNP associations and rare deletions. *PLoS Genet.* 2009; 5:e1000534. [PubMed: 19557189]
28. Ishizuka T, et al. The human phosphoribosylpyrophosphate synthetase-associated protein 39 gene (*PRPSAP1*) is located in the chromosome region 17q24-q25. *Genomics.* 1996; 33:332–334. [PubMed: 8660991]
29. Luo Z, Saha AK, Xiang X, Ruderman NB. AMPK, the metabolic syndrome and cancer. *Trends Pharmacol Sci.* 2005; 26:69–76. [PubMed: 15681023]
30. Tong X, Zhao F, Mancuso A, Gruber JJ, Thompson CB. The glucose-responsive transcription factor ChREBP contributes to glucose-dependent anabolic synthesis and cell proliferation. *Proc Natl Acad Sci USA.* 2009; 106:21660–21665. [PubMed: 19995986]
31. Levine AJ, Puzio-Kuter AM. The control of the metabolic switch in cancers by oncogenes and tumor suppressor genes. *Science.* 2010; 330:1340–1344. [PubMed: 21127244]
32. Iynedjian PB. Molecular physiology of mammalian glucokinase. *Cell Mol Life Sci.* 2009; 66:27–42. [PubMed: 18726182]
33. Quinones Galvan A, et al. Effect of insulin on uric acid excretion in humans. *Am J Physiol.* 1995; 268:E1–E5. [PubMed: 7840165]
34. Ter Maaten JC, et al. Renal handling of urate and sodium during acute physiological hyperinsulinaemia in healthy subjects. *Clin Sci (Lond).* 1997; 92:51–58. [PubMed: 9038591]
35. Becker MA, et al. Febuxostat compared with allopurinol in patients with hyperuricemia and gout. *N Engl J Med.* 2005; 353:2450–2461. [PubMed: 16339094]
36. Iwatani M, et al. Troglitazone decreases serum uric acid concentrations in type II diabetic patients and non-diabetics. *Diabetologia.* 2000; 43:814–815. [PubMed: 10907128]
37. Gokcel A, et al. Evaluation of the safety and efficacy of sibutramine, orlistat and metformin in the treatment of obesity. *Diabetes Obes Metab.* 2002; 4:49–55. [PubMed: 11874442]
38. Seber S, Ucak S, Basat O, Altuntas Y. The effect of dual PPAR $\alpha$ / $\gamma$  stimulation with combination of rosiglitazone and fenofibrate on metabolic parameters in type 2 diabetic patients. *Diabetes Res Clin Pract.* 2006; 71:52–58. [PubMed: 16009445]
39. Tsouli SG, Liberopoulos EN, Mikhailidis DP, Athyros VG, Elisaf MS. Elevated serum uric acid levels in metabolic syndrome: an active component or an innocent bystander? *Metabolism.* 2006; 55:1293–1301. [PubMed: 16979398]
40. Rizos CV, Liberopoulos EN, Mikhailidis DP, Elisaf MS. Pleiotropic effects of thiazolidinediones. *Expert Opin Pharmacother.* 2008; 9:1087–1108. [PubMed: 18422468]
41. Sulem P, et al. Identification of low-frequency variants associated with gout and serum uric acid levels. *Nat Genet.* 2011; 43:1127–1130. [PubMed: 21983786]
42. The ARIC Investigators. The Atherosclerosis Risk in Communities (ARIC) Study: design and objectives. *Am J Epidemiol.* 1989; 129:687–702. [PubMed: 2646917]
43. Park JH, et al. Estimation of effect size distribution from genome-wide association studies and implications for future discoveries. *Nat Genet.* 2010; 42:570–575. [PubMed: 20562874]
44. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet.* 2011; 88:76–82. [PubMed: 21167468]
45. Fuchsberger C, Taliun D, Pramstaller PP, Pattaro C. GWAToolbox: an R package for fast quality control and handling of genome-wide association studies meta-analysis data. *Bioinformatics.* 2012; 28:444–445. [PubMed: 22155946]
46. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics.* 2010; 26:2190–2191. [PubMed: 20616382]

47. Pe'er I, Yelensky R, Altshuler D, Daly MJ. Estimation of the multiple testing burden for genomewide association studies of nearly all common variants. *Genet Epidemiol.* 2008; 32:381–385. [PubMed: 18348202]
48. Purcell S, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet.* 2007; 81:559–575. [PubMed: 17701901]
49. Storey JD, Tibshirani R. Statistical significance for genomewide studies. *Proc Natl Acad Sci USA.* 2003; 100:9440–9445. [PubMed: 12883005]






**Figure 1.** Multiple genomic loci contain SNPs associated with serum urate concentrations. Truncated Manhattan plot showing  $-\log_{10}(P)$  values for all SNPs of the urate discovery GWAS ordered by chromosomal position. The gene closest to the SNP with the lowest  $P$  value at each locus (index SNP) is listed. Loci in gray met one but not both replication criteria. Blue triangles represent loci containing SNPs with  $P$  values below  $1 \times 10^{-25}$ .



**Figure 2.** Minor alleles of all replicated GWAS loci show direction-consistent association with serum urate concentrations and the odds of gout. Loci with significant sex-specific effects are shown in red. Lines correspond to 95% confidence intervals. The Pearson's correlation coefficient was calculated using the log odds ratio ( $\ln(\text{OR})$ ) for gout.



**Figure 3.** SNP effects on urate concentrations (mg/dl) are similar among individuals of European ancestry, African-Americans, Indians and Japanese, whereas allele frequencies vary. Comparison of SNP effects at the replicated loci and at four additional loci meeting one of the replication criteria. Left, effects on serum urate concentrations per minor allele (defined in individuals of European ancestry) sorted by effect size. Right, comparison of allele frequencies across the four different samples. Symbols are absent if the data could not be provided for reasons related to allele frequency (not polymorphic, low minor allele frequency) or poor imputation quality.

Table 1

SNPs associated with serum urate concentrations in individuals of European ancestry

SNP	Chr.	bp (Build 36)	Closest gene	GRAIL gene	AI	A2	Freq. A1	Serum urate (mg/dl)			Gout	
								Effect	s.e.	P value	OR	P value
<b>Discovered loci described previously</b>												
rs1471633	1	144435096	<i>PDZK1</i>	<i>PDZK1</i>	A	C	0.46	0.059	0.005	$1.2 \times 10^{-29}$	1.03	$2.8 \times 10^{-1}$
rs1260326	2	27584444	<i>GCKR</i>	<i>GCKR</i>	T	C	0.41	0.074	0.005	$1.2 \times 10^{-44}$	1.14	$8.2 \times 10^{-6}$
rs12498742	4	9553150	<i>SLC2A9</i>	<i>SLC2A9</i>	A	G	0.77	0.373	0.006	0 <sup>b</sup>	1.56	$1.9 \times 10^{-31}$
rs2231142	4	89271347	<i>ABCG2</i>	<i>ABCG2</i>	T	G	0.11	0.217	0.009	$1.0 \times 10^{-134}$	1.73	$1.7 \times 10^{-39}$
rs675209	6	7047083	<i>RREB1</i>	<i>RREB1</i>	T	C	0.27	0.061	0.006	$1.3 \times 10^{-23}$	1.09	$1.1 \times 10^{-2}$
rs1165151	6	25929595	<i>SLC17A1</i>	<i>SLC17A3</i>	T	G	0.47	-0.091	0.005	$7.0 \times 10^{-70}$	0.86	$5.3 \times 10^{-7}$
rs1171614	10	61139544	<i>SLC16A9</i>	<i>SLC16A9</i>	T	C	0.22	-0.079	0.007	$2.3 \times 10^{-28}$	0.91	$1.7 \times 10^{-2}$
rs2078267	11	64090690	<i>SLC22A11</i>	<i>SLC22A11</i>	T	C	0.51	-0.073	0.006	$9.4 \times 10^{-38}$	0.88	$2.3 \times 10^{-5}$
rs478607	11	64234639	<i>NRXN2</i>	<i>SLC22A12</i>	A	G	0.84	-0.047	0.007	$4.4 \times 10^{-11}$	0.97	$4.1 \times 10^{-1}$
rs3741414	12	56130316	<i>INHBC</i>	<i>INHBE</i>	T	C	0.24	-0.072	0.007	$2.2 \times 10^{-25}$	0.87	$2.7 \times 10^{-4}$
<b>New loci</b>												
rs11264341	1	153418117	<i>TRIM46</i>	<i>PKLR</i>	T	C	0.43	-0.050	0.006	$6.2 \times 10^{-19}$	0.92	$7.4 \times 10^{-3}$
rs17050272	2	121022910	<i>INHBB</i>	<i>INHBB</i>	A	G	0.43	0.035	0.006	$1.6 \times 10^{-10}$	1.03	$3.9 \times 10^{-1}$
rs2307394 <sup>a</sup>	2	148432898	<i>ORC4L</i>	<i>ACVR2A</i>	T	C	0.68	-0.029	0.005	$2.2 \times 10^{-8}$	0.94	$6.3 \times 10^{-2}$
rs6770152	3	53075254	<i>SFMBT1</i>	<i>MUSTN1</i>	T	G	0.58	-0.044	0.005	$2.6 \times 10^{-16}$	0.90	$3.0 \times 10^{-4}$
rs17632159	5	72467238	<i>TMEM171</i>	<i>TMEM171</i>	C	G	0.31	-0.039	0.006	$3.5 \times 10^{-11}$	0.91	$6.0 \times 10^{-3}$
rs729761	6	43912549	<i>VEGFA</i>	<i>VEGFA</i>	T	G	0.30	-0.047	0.006	$8.0 \times 10^{-16}$	0.87	$4.1 \times 10^{-5}$
rs1178977	7	72494985	<i>BAZ1B</i>	<i>MLX1PL</i>	A	G	0.81	0.047	0.007	$1.2 \times 10^{-12}$	1.14	$6.7 \times 10^{-4}$
rs10480300	7	151036938	<i>PRKAG2</i>	<i>PRKAG2</i>	T	C	0.28	0.035	0.006	$4.1 \times 10^{-9}$	1.09	$6.5 \times 10^{-3}$
rs17786744	8	23832951	<i>STC1</i>	<i>STC1</i>	A	G	0.58	-0.029	0.005	$1.4 \times 10^{-8}$	0.93	$2.1 \times 10^{-2}$
rs2941484	8	76641323	<i>HNF4G</i>	<i>HNF4G</i>	T	C	0.44	0.044	0.005	$4.4 \times 10^{-17}$	1.04	$1.7 \times 10^{-1}$
rs10821905	10	52316099	<i>A1CF</i>	<i>ASAH2</i>	A	G	0.18	0.057	0.007	$7.4 \times 10^{-17}$	1.09	$2.6 \times 10^{-2}$
rs642803 <sup>a</sup>	11	65317196	<i>OVOL1</i>	<i>LTBP3</i>	T	C	0.46	-0.036	0.005	$2.9 \times 10^{-13}$	0.90	$2.2 \times 10^{-4}$
rs653178	12	110492139	<i>ATXN2</i>	<i>PTPN11</i>	T	C	0.51	-0.035	0.005	$7.2 \times 10^{-12}$	0.95	$7.3 \times 10^{-2}$
rs1394125	15	73946038	<i>UBE2Q2</i>	<i>NRG4</i>	A	G	0.34	0.043	0.006	$2.5 \times 10^{-13}$	1.03	$3.6 \times 10^{-1}$



SNP	Chr.	bp (Build 36)	Closest gene	GRAIL gene	A1	A2	Freq. A1	Serum urate (mg/dl)			Gout	
								Effect	s.e.	P value	OR	P value
rs6598541	15	97088658	<i>IGF1R</i>	<i>IGF1R</i>	A	G	0.36	0.043	0.006	$4.8 \times 10^{-15}$	1.04	$2.5 \times 10^{-1}$
rs7193778	16	68121391	<i>NFAT5</i>	<i>NFAT5</i>	T	C	0.86	-0.046	0.008	$8.2 \times 10^{-10}$	0.92	$3.4 \times 10^{-2}$
rs7188445	16	78292488	<i>MAF</i>	<i>MAF</i>	A	G	0.33	-0.032	0.005	$1.6 \times 10^{-9}$	0.95	$1.1 \times 10^{-1}$
rs7224610	17	50719787	<i>HLF</i>	<i>HLF</i>	A	C	0.58	-0.042	0.005	$5.4 \times 10^{-17}$	0.96	$1.6 \times 10^{-1}$
rs2079742 <sup>a</sup>	17	56820479	<i>BCAS3</i>	<i>C17orf82</i>	T	C	0.85	0.043	0.008	$1.2 \times 10^{-8}$	1.04	$3.5 \times 10^{-1}$
rs164009 <sup>a</sup>	17	71795264	<i>QRICH2</i>	<i>PRPSAP1</i>	A	G	0.61	0.028	0.005	$1.6 \times 10^{-7}$	1.08	$8.6 \times 10^{-3}$

P values are corrected for inflation using genomic control. Median imputation quality was calculated across all cohorts for urate concentration-associated SNPs in the overall sample and ranged from 0.83 to 1 (median of 0.98). Allele frequencies are presented for the discovery sample. Gout estimates are provided for the combined discovery and validation samples. Sample sizes available for the individual SNPs are listed in Supplementary Table 6. Chr., chromosome; A1, allele 1, effect allele; freq., frequency; s.e., standard error.

<sup>a</sup>These SNPs met one but not both criteria required for replication.

<sup>b</sup>P value  $< 1 \times 10^{-700}$ .

**Table 2**

Urate concentration-associated SNPs from secondary analyses

SNP	Chr.	bp (Build 36)	Gene	A1	A2	Freq. A1	Serum urate (mg/dl)			Gout	
							Effect	s.e.	P value	OR	P value
<b>Urate transporter candidate analysis</b>											
rs4149178 <sup>a</sup>	6	43380166	<i>SLC22A7</i>	A	G	0.84	-0.034	0.007	$1.2 \times 10^{-6}$	0.95	$2.2 \times 10^{-1}$
<b>Functional association network analysis</b>											
rs4970988 <sup>a</sup>	1	149216686	<i>ARNT</i>	A	G	0.36	-0.028	0.005	$1.0 \times 10^{-7}$	0.96	$2.1 \times 10^{-1}$
rs7976059	12	50537539	<i>ACVRL1-ACVRL1</i>	T	G	0.35	0.032	0.005	$1.9 \times 10^{-9}$	0.95	$1.8 \times 10^{-1}$
rs7953704	12	121191945	<i>B3GNT4</i>	A	G	0.47	-0.029	0.005	$2.6 \times 10^{-8}$	0.95	$1.6 \times 10^{-1}$

Allele frequencies are presented for the discovery sample. Chr., chromosome; A1, allele 1, effect allele; freq., frequency; s.e., standard error.

<sup>a</sup>These SNPs met one but not both criteria required for replication.

**Table 3**  
Nominally significant associations between urate concentration–associated SNPs and FEUA

SNP	Closest gene	A1	A2	Serum urate (mg/dl)			FEUA (%)	
				Effect	P value	P value	Effect	s.e.
rs1260326	<i>GCKR</i>	T	C	0.074	$1.2 \times 10^{-44}$	–0.073	0.018	$3.0 \times 10^{-5}$
rs12498742	<i>SLC2A9</i>	A	G	0.373	0 <sup>a</sup>	–0.184	0.022	$1.2 \times 10^{-16}$
rs2231142	<i>ABCG2</i>	T	G	0.217	$1.0 \times 10^{-134}$	–0.076	0.030	$9.8 \times 10^{-3}$
rs675209	<i>RREB1</i>	T	C	0.061	$1.3 \times 10^{-23}$	–0.064	0.021	$2.2 \times 10^{-3}$
rs2078267	<i>SLC22A11</i>	T	C	–0.073	$9.4 \times 10^{-38}$	0.042	0.020	$3.7 \times 10^{-2}$
rs478607	<i>NRXN2</i>	A	G	–0.047	$4.4 \times 10^{-11}$	0.046	0.023	$4.6 \times 10^{-2}$
rs1394125	<i>UBE2Q2</i>	A	G	0.043	$2.5 \times 10^{-13}$	–0.059	0.023	$1.1 \times 10^{-2}$
rs6598541	<i>IGF1R</i>	A	G	0.043	$4.8 \times 10^{-15}$	–0.059	0.019	$1.5 \times 10^{-3}$
rs7193778	<i>NFAT5</i>	T	C	–0.046	$8.2 \times 10^{-10}$	0.086	0.032	$7.2 \times 10^{-3}$
rs7224610	<i>HLF</i>	A	C	–0.042	$5.4 \times 10^{-17}$	0.045	0.017	$9.3 \times 10^{-3}$

Serum urate concentration estimates are provided for the combined discovery and replication sample. Sample size for the SNPs for FEUA ranged from 6,798 to 6,799. A1, allele 1, effect allele; s.e., standard error.

<sup>a</sup>  $P < 1 \times 10^{-700}$ .