


Three Lemmas in Geometry


Yufei Zhao

Massachusetts Institute of Technology

1 Diameter of incircle



Lemma 1. Let the incircle of triangle ABC touch side BC at D , and let DT be a diameter of the circle. If line AT meets BC at X , then $BD = CX$.



Proof. Assume wlog that $AB \geq AC$. Consider the dilation with center A that carries the incircle to an excircle. The line segment DT is the diameter of the incircle that is perpendicular to BC , and therefore its image under the dilation must be the diameter of the excircle that is perpendicular to BC . It follows that T must get mapped to the point of tangency between the excircle and BC . In addition, the image of T must lie on the line AT , and hence T gets mapped to X . Thus, the excircle is tangent to BC at X .

It remains to prove that $BD = CX$. Let the incircle of ABC touch sides AB and AC at F and E , respectively. Let the excircle of ABC opposite to A touch rays AB and AC at Z and Y , respectively,

then using equal tangents, we have

$$\begin{aligned} 2BD &= BF + BX + XD = BF + BZ + XD = FZ + XD \\ &= EY + XD = EC + CY + XD = DC + XC + XD = 2CX. \end{aligned}$$

Thus $BD = CX$. □

Problems

1. (IMO 1992) In the plane let \mathcal{C} be a circle, ℓ a line tangent to the circle \mathcal{C} , and M a point on ℓ . Find the locus of all points P with the following property: there exists two points Q, R on ℓ such that M is the midpoint of QR and \mathcal{C} is the inscribed circle of triangle PQR .
2. (USAMO 1999) Let $ABCD$ be an isosceles trapezoid with $AB \parallel CD$. The inscribed circle ω of triangle BCD meets CD at E . Let F be a point on the (internal) angle bisector of $\angle DAC$ such that $EF \perp CD$. Let the circumscribed circle of triangle ACF meet line CD at C and G . Prove that the triangle AFG is isosceles.
3. (IMO Shortlist 2005) In a triangle ABC satisfying $AB + BC = 3AC$ the incircle has centre I and touches the sides AB and BC at D and E , respectively. Let K and L be the symmetric points of D and E with respect to I . Prove that the quadrilateral $ACKL$ is cyclic.
4. (Nagel line) Let ABC be a triangle. Let the excircle of ABC opposite to A touch side BC at D . Similarly define E on AC and F on AB . Then AD, BE, CF concur (why?) at a point N known as the *Nagel point*.

Let G be the centroid of ABC and I the incenter of ABC . Show that I, G, N lie in that order on a line (known as the *Nagel line*, and $GN = 2IG$).

5. (USAMO 2001) Let ABC be a triangle and let ω be its incircle. Denote by D_1 and E_1 the points where ω is tangent to sides BC and AC , respectively. Denote by D_2 and E_2 the points on sides BC and AC , respectively, such that $CD_2 = BD_1$ and $CE_2 = AE_1$, and denote by P the point of intersection of segments AD_2 and BE_2 . Circle ω intersects segment AD_2 at two points, the closer of which to the vertex A is denoted by Q . Prove that $AQ = D_2P$.
6. (Tournament of Towns 2003 Fall) Triangle ABC has orthocenter H , incenter I and circumcenter O . Let K be the point where the incircle touches BC . If IO is parallel to BC , then prove that AO is parallel to HK .
7. (IMO 2008) Let $ABCD$ be a convex quadrilateral with $|BA| \neq |BC|$. Denote the incircles of triangles ABC and ADC by ω_1 and ω_2 respectively. Suppose that there exists a circle ω tangent to the ray BA beyond A and to the ray BC beyond C , which is also tangent to the lines AD and CD . Prove that the common external tangents of ω_1 and ω_2 intersect on ω .

(Hint: show that $AB + AD = CB + CD$. What does this say about the lengths along AC ?)

2 Center of spiral similarity

A *spiral similarity*¹ about a point O (known as the center of the spiral similarity) is a composition of a rotation and a dilation, both centered at O . (See diagram)




Figure 1: An example of a spiral similarity.

For instance, in the complex plane, if $O = 0$, then spiral similarities are described by multiplication by a nonzero complex number. That is, spiral similarities have the form $z \mapsto \alpha z$, where $\alpha \in \mathbb{C} \setminus \{0\}$. Here $|\alpha|$ is the dilation factor, and $\arg \alpha$ is the angle of rotation. It is easy to deduce from here that if the center of the spiral similarity is some other point, say z_0 , then the transformation is given by $z \mapsto z_0 + \alpha(z - z_0)$ (why?).

Fact. Let A, B, C, D be four distinct point in the plane such that $ABCD$ is not a parallelogram. Then there exists a unique spiral similarity that sends A to B , and C to D .

Proof. Let a, b, c, d be the corresponding complex numbers for the points A, B, C, D . We know that a spiral similarity has the form $\mathbf{T}(z) = z_0 + \alpha(z - z_0)$, where z_0 is the center of the spiral similarity, and α is data on the rotation and dilation. So we would like to find α and z_0 such that $\mathbf{T}(a) = c$ and $\mathbf{T}(b) = d$. This amount to solving the system

$$z_0 + \alpha(a - z_0) = c, \quad z_0 + \alpha(b - z_0) = d.$$

Solving it, we see that the unique solution is

$$\alpha = \frac{c - d}{a - b}, \quad z_0 = \frac{ad - bc}{a - b - c + d}.$$

Since $ABCD$ is not a parallelogram, $a - b - c + d \neq 0$, so that this is the unique solution to the system. Hence there exists a unique spiral similarity that carries A to B and C to D . \square


Exercise. How can you quickly determine the value of α in the above proof without even needing to set up the system of equations?

Exercise. Give a geometric argument why the spiral similarity, if it exists, must be unique. (Hint: suppose that \mathbf{T}_1 and \mathbf{T}_2 are two such spiral similarities, then what can you say about $\mathbf{T}_1 \circ \mathbf{T}_2^{-1}$?)

So we know that a spiral similarity exists, but where is its center? The following lemma tells us how to locate it.

¹If you want to impress your friends with your mathematical vocabulary, a spiral similarity is sometimes called a *similitude*, and a dilation is sometimes called a *homothety*. (Actually, they are not quite exactly the same thing, but shhh!)

Lemma 2. Let A, B, C, D be four distinct point in the plane, such that AC is not parallel to BD . Let lines AC and BD meet at X . Let the circumcircles of ABX and CDX meet again at O . Then O is the center of the unique spiral similarity that carries A to C and B to D .



Proof. We use directed angles mod π (i.e., directed angles between *lines*, as opposed to rays) in order to produce a single proof that works in all configurations. Let $\angle(\ell_1, \ell_2)$ denote the angle of rotation that takes line ℓ_1 to ℓ_2 . A useful fact is that points P, Q, R, S are concyclic if and only if $\angle(PQ, QR) = \angle(PS, SR)$.

We have

$$\angle(OA, AC) = \angle(OA, AX) = \angle(OB, BX) = \angle(OB, BD),$$

and

$$\angle(OC, CA) = \angle(OC, CX) = \angle(OD, DX) = \angle(OD, DB).$$

It follows that triangles AOC and BOD are similar and have the same orientation. Therefore, the spiral similarity centered at O that carries A to C must also carry B to D . \square

Finally, it is worth mentioning that spiral similarities come in pairs. If we can send AB to CD , then we can just as easily send AC to BD using the same center.

Fact. If O is the center of the spiral similarity that sends A to C and B to D , then O is also the center of the spiral similarity that sends A to B and C to D .

Proof. Since spiral similarity preserves angles at O , we have $\angle AOB = \angle COD$. Also, the dilation ratio of the first spiral similarity is $OC/OA = OD/OB$. So the rotation about O with angle $\angle AOB = \angle COD$ followed by a dilation with ratio $OB/OA = OD/OC$ sends A to B , and C to D , as desired. \square

Problems

- (IMO Shortlist 2006) Let $ABCDE$ be a convex pentagon such that

$$\angle BAC = \angle CAD = \angle DAE \quad \text{and} \quad \angle CBA = \angle DCA = \angle EDA.$$


Diagonals BD and CE meet at P . Prove that line AP bisects side CD .

- (USAMO 2006) Let $ABCD$ be a quadrilateral, and let E and F be points on sides AD and BC , respectively, such that $AE/ED = BF/FC$. Ray FE meets rays BA and CD at S and T , respectively. Prove that the circumcircles of triangles SAE , SBF , TCF , and TDE pass through a common point.
- (China 1992) Convex quadrilateral $ABCD$ is inscribed in circle ω with center O . Diagonals AC and BD meet at P . The circumcircles of triangles ABP and CDP meet at P and Q . Assume that points O, P , and Q are distinct. Prove that $\angle OQP = 90^\circ$.

4. Let $ABCD$ be a quadrilateral. Let diagonals AC and BD meet at P . Let O_1 and O_2 be the circumcenters of APD and BPC . Let M, N and O be the midpoints of AC, BD and O_1O_2 . Show that O is the circumcenter of MPN .
5. (Miquel point of a quadrilateral) Let $\ell_1, \ell_2, \ell_3, \ell_4$ be four lines in the plane, no two parallel. Let C_{ijk} denote the circumcircle of the triangle formed by the lines ℓ_i, ℓ_j, ℓ_k (these circles are called *Miquel circles*). Then $C_{123}, C_{124}, C_{134}, C_{234}$ pass through a common point (called the *Miquel point*).
(It's not too hard to prove this result using angle chasing, but can you see why it's almost an immediate consequence of the lemma?)
6. (IMO 2005) Let $ABCD$ be a given convex quadrilateral with sides BC and AD equal in length and not parallel. Let E and F be interior points of the sides BC and AD respectively such that $BE = DF$. The lines AC and BD meet at P , the lines BD and EF meet at Q , the lines EF and AC meet at R . Consider all the triangles PQR as E and F vary. Show that the circumcircles of these triangles have a common point other than P .
7. (IMO Shortlist 2006) Points A_1, B_1 and C_1 are chosen on sides BC, CA , and AB of a triangle ABC , respectively. The circumcircles of triangles AB_1C_1, BC_1A_1 , and CA_1B_1 intersect the circumcircle of triangle ABC again at points A_2, B_2 , and C_2 , respectively ($A_2 \neq A, B_2 \neq B$, and $C_2 \neq C$). Points A_3, B_3 , and C_3 are symmetric to A_1, B_1, C_1 with respect to the midpoints of sides BC, CA , and AB , respectively. Prove that triangles $A_2B_2C_2$ and $A_3B_3C_3$ are similar.


3 Symmedian

Let ABC be a triangle. Let M be the midpoint of BC , so that AM is a *median* of ABC . Let N be a point on side BC so that $\angle BAM = \angle CAN$. Then AN is a *symmedian* of ABC . In other words, *the symmedian is the reflection of the median across the angle bisector*. The following lemma gives an important property of the symmedian.



Lemma 3. Let ABC be a triangle and Γ its circumcircle. Let the tangent to Γ at B and C meet at D . Then AD coincides with a symmedian of $\triangle ABC$.

We give three proofs. (The three diagram each correspond to a separate proof.) The first proof is a “sine law chase.”



First proof. Let the reflection of AD across the angle bisector of $\angle BAC$ meet BC at M' . Then

$$\begin{aligned}
 \frac{BM'}{M'C} &= \frac{AM' \frac{\sin \angle BAM'}{\sin \angle ABC}}{AM' \frac{\sin \angle CAM'}{\sin \angle ACB}} && \text{[Sine law on } ABM' \text{ and } ACM'] \\
 &= \frac{\sin \angle BAM' \sin \angle ABD}{\sin \angle ACD \sin \angle CAM'} && \text{[Using the tangent-chord angles]} \\
 &= \frac{\sin \angle CAD \sin \angle ABD}{\sin \angle ACD \sin \angle BAD} && \text{[From construction of } M'] \\
 &= \frac{CD}{AD} \frac{AD}{BD} && \text{[Sine law on } ACD \text{ and } ABD] \\
 &= 1.
 \end{aligned}$$

Therefore, AM' is the median, and thus AD is the symmedian. □

Remark. Some people like to start this proof by setting M to be the midpoint of BC , and then using sine law to show that $\frac{\sin \angle BAD}{\sin \angle CAD} = \frac{\sin \angle CAM}{\sin \angle BAM}$. I do not recommend this variation, since it's not immediately clear that $\angle CAM = \angle BAD$ follows, especially when $\angle BAD$ is obtuse.

Next we give a synthetic proof that highlights some additional features in the configuration.

Second proof. Let O be the circumcenter of ABC and let ω be the circle centered at D with radius DB . Let lines AB and AC meet ω at P and Q , respectively. Since $\angle ABC = \angle AQP$, triangles ABC and AQP are similar. The idea is use the fact that, up to dilation, triangles ABC and AQP are reflections of each other across the angle bisector of $\angle A$.

Since

$$\angle PBQ = \angle BQC + \angle BAC = \frac{1}{2}(\angle BDC + \angle BOC) = 90^\circ,$$

we see that PQ is a diameter of ω and hence passes through D . Let M be the midpoint of BC . Since D is the midpoint of QP , the similarity implies that $\angle BAM = \angle QAD$, from which the result follows. □

The third proof uses facts from projective geometry. Feel free to skip it if you are not comfortable with projective geometry.

Third proof. Let the tangent of Γ at A meet line BC at E . Then E is the pole of AD (since the polar of A is AE and the pole of D is BC). Let BC meet AD at F . Then point B, C, E, F are harmonic. This means that line AB, AC, AE, AF are harmonic. Consider the reflections of the four line across the angle bisector of $\angle BAC$. Their images must be harmonic too. It's easy to check that AE maps onto a line

parallel to BC . Since BC must meet these four lines at harmonic points, it follows that the reflection of AF must pass through the midpoint of BC . Therefore, AF is a symmedian. \square

Problems

1. (Poland 2000) Let ABC be a triangle with $AC = BC$, and P a point inside the triangle such that $\angle PAB = \angle PBC$. If M is the midpoint of AB , then show that $\angle APM + \angle BPC = 180^\circ$.
2. (IMO Shortlist 2003) Three distinct points A, B, C are fixed on a line in this order. Let Γ be a circle passing through A and C whose center does not lie on the line AC . Denote by P the intersection of the tangents to Γ at A and C . Suppose Γ meets the segment PB at Q . Prove that the intersection of the bisector of $\angle AQC$ and the line AC does not depend on the choice of Γ .
3. (Vietnam TST 2001) In the plane, two circles intersect at A and B , and a common tangent intersects the circles at P and Q . Let the tangents at P and Q to the circumcircle of triangle APQ intersect at S , and let H be the reflection of B across the line PQ . Prove that the points A, S , and H are collinear.
4. (USA TST 2007) Triangle ABC is inscribed in circle ω . The tangent lines to ω at B and C meet at T . Point S lies on ray BC such that $AS \perp AT$. Points B_1 and C_1 lie on ray ST (with C_1 in between B_1 and S) such that $B_1T = BT = C_1T$. Prove that triangles ABC and AB_1C_1 are similar to each other.
5. Let ABC be a triangle. Let X be the center of spiral similarity that takes B to A and A to C . Show that AX coincides with a symmedian of ABC .
6. (USA TST 2008) Let ABC be a triangle with G as its centroid. Let P be a variable point on segment BC . Points Q and R lie on sides AC and AB respectively, such that $PQ \parallel AB$ and $PR \parallel AC$. Prove that, as P varies along segment BC , the circumcircle of triangle AQR passes through a fixed point X such that $\angle BAG = \angle CAX$.
7. (USA 2008) Let ABC be an acute, scalene triangle, and let M, N , and P be the midpoints of BC, CA , and AB , respectively. Let the perpendicular bisectors of AB and AC intersect ray AM in points D and E respectively, and let lines BD and CE intersect in point F , inside of triangle ABC . Prove that points A, N, F , and P all lie on one circle.
8. Let A be one of the intersection points of circles ω_1, ω_2 with centers O_1, O_2 . The line ℓ is tangent to ω_1, ω_2 at B, C respectively. Let O_3 be the circumcenter of triangle ABC . Let D be a point such that A is the midpoint of O_3D . Let M be the midpoint of O_1O_2 . Prove that $\angle O_1DM = \angle O_2DA$.
(Hint: use Problem 5.)